

RED-TOP

BEN-SUL 60 DUST

DIRECTIONS FOR USE: This material may be applied with ground equipment or aircraft.

Where recommended for prevention of Powdery Mildew, apply when disease first appears and repeat at 7 to 10 day intervals throughout the growing season. Shorten the interval during humid, rainy weather.

Where recommended for prevention of Brown Rot, apply at pink bud or popcorn stage during bloom and at petal fall, shuck split and shuck fall. Repeat at 10 to 14 day intervals in cover sprays as long as necessary.

RECOMMENDATIONS FOR USE:

ALFALFA: For control of Lygus Atlantic Mite, Spider Mite, and Clover Mite use 30 to 50 lbs. per acre.

BEANS: For the control of Pacific Mite (Red Spider) use 25 lbs. per acre. For control of Powdery Mildew use 40 to 50 lbs. per acre. Certain varieties of beans may be injured by sulfur.

CORN: For the control of Spider Mites use 30 to 50 lbs. per acre.

COTTON: For control of Mites, Lygus and Flea Hopper apply 10-50 lbs. per acre.

GRAPES: For the control of Powdery Mildew use 8 to 20 lbs. per acre. Make first application when shoots are 6 inches to 8 inches long. Make a second application when shoots are 12 inches to 15 inches long and the third about 14 days later. Repeat applications at 10 to 14 day intervals to protect new growth. Avoid applications during hot weather as injury may occur. Concord and other labrusca type grapes may be injured by sulfur.

HOES: For control of Red Spider Mites use 30 to 50 lbs. per acre.


PEACHES: For the control of Peach Silver Mite, Powdery Mildew, and prevention of Brown Rot of fruit apply 40 to 50 lbs. per acre. For prevention of Brown Rot of fruit apply when fruit begins to ripen.

ACTIVE INGREDIENT

Sulfur

INERT INGREDIENTS

EPA Reg. No. 2935-2



CAUTION: KEEP OUT OF REACH OF CHILDREN

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (90° or above). Do not make sulfur applications at such times.

Do not use this product for any purposes other than recommended on this label.

Do not use within 2 weeks of an oil spray treatment. When growing crops for processing, consult the processor before applying sulfur.

Consult your State Agricultural Experimental Station or your State Agricultural Extension Service for additional information consistent with label cautions and directions.

Do not reuse container. Destroy when empty.

RECOMMENDATIONS FOR USE: (Continued)

PLUMS and PRUNES: For prevention of Brown Rot of fruit apply 40 to 50 lbs. per acre. Apply when fruit starts to ripen.

SUGAR BEETS: For control of Spider Mites and Powdery Mildew apply 60 to 80 lbs. per acre.

IN CASE OF EMERGENCY, CALL CHEMTREC (800) 424-9300