

RECOMMENDATIONS FOR USE:

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION: Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—60 to 800 gals. Citrus—400 to 3,000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre).

Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals.

MIXING DIRECTIONS: Add to spray tank when 1/4 to 1/2 full with agitator running and maintain agitation until spraying is completed.

ALFALFA: Use 1/2 pt. per acre to control GRASSHOPPER NYMPHS when crop is no more than 12" high. Use 1 pt. per acre to control APHIDS, LOOPERS, LEAFHOPPERS, LYGUS BUGS.

ALFALFA WEBWORM AND ARMYWORMS. Use 1/2 pt. per acre to control [redacted] LYGUS BUG on SEED ALFALFA. Combine with recommended chemicals for residual control. Do not apply within 4 days of harvest.

BEANS (dry, lima & green) & PEAS for processing: Use 1 pt. per acre to control ALFALFA LOOPER, APHIDS, RED SPIDER MITES, LYGUS BUG, LEAF MINER, NITIDULID POLLEN BEETLES & LEAFHOPPERS. For control of Large LOOPERS, use 1 1/2 pts. per acre. Do not apply within 4 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE & COLLARDS: Use 1 pt. per acre to control IMPORTED CABBAGEWORM, DIAMOND BACK CATERPILLARS & APHIDS. Use 2 pts. per acre to control CABBAGE LOOPERS. Do not apply within 4 days of harvest on Kale and Collards. Do not use within 1 day of harvest on Broccoli, Cabbage, Cauliflower & Brussels Sprouts.

COTTON: Use 1 pt. per acre to control LYGUS BUG, COTTON LEAF PERFORATOR, FLEAHOPPERS, ARMYWORMS, BOLLWORMS, LOOPERS & to SUPPRESS MITES. Apply at 7 to 10 day intervals while pests persist. For heavy insect or egg population of any listed pest, use 1/2 pt. per acre plus residual materials for best results. Do not apply within 4 days of hand-picking.

CUCUMBERS & SUMMER SQUASH: Use 1 pt. per acre to control THRIPS, SPIDER MITES, LOOPERS, ARMYWORM, CUCUMBER BEETLES & APHIDS. Use 1 1/2 pts. per acre to control LEAF MINERS & PICKLEWORMS. May be used up to harvest.

EGGPLANT & PEPPERS: Use 1 pt. per acre to control APHIDS, RED SPIDER MITES, BLISTER BEETLES, FLEA BEETLES, WHITEFLY & LEAF MINERS. Do not apply within 1 day of harvest.

WALNUTS: Use 1/2 to 1 pt. per 100 gals. water in a full coverage spray to control APHIDS. Do not apply within 10 days of harvest.

GRAPES: Use 1/2 to 2/3 pt. per 100 gals. water to control DROSOPHILA & LEAFHOPPERS. Apply 100 to 150 gals. diluted spray per acre depending on size of vines. Use only 1/2 pt. per 100 gals. water on Muscat varieties. Do not use on Italia varieties as fruit injury may occur. Do not apply within 4 days of harvest.

HOPS: Use 1 pt. per acre to control APHIDS, RED SPIDER MITES, CORN EARWORM, HOP LOOPER & CUTWORMS. Do not apply within 4 days of harvest.

red-top

DIBROM® 8 SPRAY

NALED INSECTICIDE

COMPOSITION:		By Wt.
ACTIVE INGREDIENTS:		
*Naled	58%
Xylene	20%
INERT INGREDIENTS:	22%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate		100%
Contains 8 pounds technical NALED. Equivalent to 7.2 pounds NALED per gallon.		
USDA Reg. No. 2935-284		

CAUTION: KEEP OUT OF REACH OF CHILDREN!

See side panel for additional cautions.

STORE IN COOL, DRY PLACE. WHEN CONTAINER IS EMPTY, IMMEDIATELY WASH THOROUGHLY AND DESTROY. NEVER RE-USE.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

WILBUR-ELLIS COMPANY

FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

2935-284

9-3-71

1043

RECOMMENDATIONS FOR USE: (Continued)

LETTUCE, CELERY, SPINACH & CHARD: Use 1 pint in 50 to 75 gals. water per acre on plants less than $\frac{1}{4}$ grown to control APHIDS, LOOPERS, ARMYWORMS, SALT MARSH CATERPILLARS & LEAF MINERS. On larger plants use $1\frac{1}{2}$ pints in 100 to 150 gals. water per acre. Use ground application only with fine mist spray. Do not apply within 4 days of harvest on Celery or within 1 day on Lettuce, Spinach and Chard.

MELONS (Cantaloupes, Honeydews, Muskmelons,) Pumpkins & Winter Squash: Use 1 to 2 pts. per acre (200 gals. diluted spray per acre) to control LEAFROLLER, ~~CATERPILLARS~~, CABBAGE LOOPERS, ARMYWORMS, LEAFHOPPERS, TWO-SPOTTED MITES & DIPTEROUS LEAFMINERS. Do not apply when temperature is over 90°F. ~~Do not apply when temperature is over 90°F.~~ as injury may result. Do not apply within 1 day of harvest.

ORANGES, LEMONS & GRAPEFRUIT: Use 1 pt. per 100 gals. water to control APHIDS, SOFT BROWN SCALE CRAWLERS, MITES, CITRUS CUTWORMS & LEAF ROLLER (Forrix). Use $\frac{1}{2}$ pt. per 100 gals. to control THRIPS. Do not apply within 7 days of harvest. (In Calif. and Ariz. - do not apply within 1 day of harvest.)

PEACHES (Western States Only): Use $\frac{1}{2}$ to $\frac{3}{4}$ pt. per 100 gals. water to control LYGUS, PEACH TWIG BORER, ORIENTAL FRUIT MOTH & to suppress MITES. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90°F. Do not apply under any circumstances, within 4 days of harvest.

STRAWBERRIES: Use 1 pt. per acre to control RED SPIDER MITES, APHIDS, OMNIVOROUS LEAF TIER & SPITTLEBUGS. Do not apply within 1 day of harvest.

TOMATOES: (Field) Use 1 pt. per acre to control TOMATO FRUIT WORM, HORNWORMS & LEAFMINERS. Do not apply when temperature is over 90°F. Do not apply within 1 day of harvest.

CAUTION: Concentrate may cause skin damage. Do not get on skin, eyes or clothing. Use water-proof gloves and face shield or goggles when handling concentrate. Harmful if swallowed. Avoid breathing spray mist. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Avoid contamination of feed, foodstuffs and drinking water. Keep children and animals off treated areas. Do not use on ornamental Cherries, Nectarines, Liquidamber or Chrysanthemums as injury may occur. Do not store diluted spray. Make new dilution for each use. Wash equipment thoroughly after use.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Containers should be disposed of by breaking and burying with wastes.

R DIBROM is Chevron Chemical Company's Reg. T.M.
F-571-284

U. S. Pat. No. 2,973,882
NET CONTENTS: 1 GALLON

2013

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

ACCEPTED

SEP 3 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON R GISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

2935-284