



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 14 1994

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Joanne Pratt
WILBUR ELLIS CO.
191 W. Shaw Avenue
Fresno, CA 93704

Subject: Label Amendment Submission of 09/10/93 in Response to PR Notice 93-7
EPA Reg. No. 2935-284
DIBROM 8 SPRAY

IMPORTANT NOTICE: This Letter Requires You to Submit a Notification to the Office of Compliance Monitoring within 20 Days.

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

SERIOUS ERROR(S) ON SUBMITTED LABELING:

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons, or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING. Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.



Recycled/Recyclable
Printed with Soy/Canola ink or paper that
contains at least 50% recycled fiber

Each notification must include the following information:

1. Registrant's name and address.
2. EPA Registration number of the product for which the labeling was submitted.
3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

1. Quantity of the product that has been sold or distributed.
2. EPA Producing Establishment number.
3. Location of the product, if known.
4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director
 Compliance Division, (7203)
 Worker Protection Notification
 Office of Compliance Monitoring
 U.S. Environmental Protection Agency
 401 M Street SW
 Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- **BEFORE** selling or distributing any product bearing the final printed labeling as amended

AND

- **WITHIN** one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

**Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001**

Hand or courier deliveries of final labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

cc: Office of Compliance Monitoring

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMAN AND DOMESTIC ANIMALS**

DANGER

Keep children and animals off treated areas. Do not permit children to go on grass until spray has completely dried.

DANGER: Corrosive, causes eye and skin damage. May be fatal if swallowed. Avoid breathing spray mist. Wear a respirator fully approved by the Mining, Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11. Do not get on skin, eyes or clothing. Use waterproof gloves and face shield or goggles when handling concentrate.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

*Add Copy F
Add Copy B*

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not spray on Nectarines, ornamental Cherries, Liquidamber or Chrysanthemums as injury may occur.

Do not store diluted spray. Make new dilution for each use.

PHYSICAL OR CHEMICAL HAZARDS

Wash metal equipment thoroughly after use to avoid corrosion. Do not contaminate water by cleaning of equipment or disposal of wastes.

Replace with Copy D

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY TREATMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

45964



1145 11 5119 1111

DIBROM 8 SPRAY

**NALED INSECTICIDE
ORGANO PHOSPHATE**

ACTIVE INGREDIENTS:	By Wt.
*Naled.....	53%
INERT INGREDIENTS:	42%
Xylene Range Aromatic Solvent.....	20%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	100%
Contains 8 pounds technical NALED. Equivalent to 7.2 pounds NALED per gallon.	

EPA Reg. No. 2935-284-ZA

EPA Est. No. 5481-CA-1

**KEEP OUT OF REACH OF CHILDREN
DANGER — PELIGRO**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le sea puesta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

In Case of Contact — Immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention.

If Swallowed — Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

NOTE TO PHYSICIAN: Naled is a cholinesterase inhibitor. Atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine. Mucosal damage may occur and if gastric lavage is indicated it should be performed cautiously.

REFER TO SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS: 4 x 1 GALLON

WILBUR-ELLIS COMPANY

P.O. Box 16458 • Fresno, CA 93755

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 14 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2935-284

BEST AVAILABLE COPY

RE-ENTRY TREATMENT (cont.)

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing (see above), areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment (see right) in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Danger. Area treated with Dibrom on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here action to take in case of accidental exposure — see left.)"

STORAGE AND DISPOSAL


1. **Prohibitions:** Do not contaminate water, food or feed by storage or disposal.
2. **Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.
3. **Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. **Other:** Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

USE ONLY ACCORDING TO ATTACHED DIRECTIONS

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

DIBROM is a registered trademark of Valent U.S.A. for naled insecticide.
Logo TM of WILBUR-ELLIS Company
IDEAS TO GROW WITH is a registered trademark of WILBUR-ELLIS Company

IN CASE OF EMERGENCY, CAL



RQ, CORROSIVE LIQ

UN 1

F-793

BEST AVAILABLE COPY

NOT FOR USE OR STORAGE IN OR AROUND HOME.

Observe Precautions on Container Label.

DIRECTIONS FOR USE

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION: Vegetable and Field Crops — 30 to 250 gals. Cotton — 3 to 20 Deciduous Fruit Crops — 80 to 800 gals. Citrus — 400 to 3,000 gals. (On Young Fruit Plants — Use a minimum of 15 gals. water per acre.)

Where Recommended for AIR APPLICATION: Field Crops — 3 to 10 gallons.

A short residual contact insecticide for kill of insect and mite adults and larvae present at the time of application.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when $\frac{1}{4}$ to $\frac{1}{2}$ full with agitator running and make agitation until spraying is completed.

Do not apply this product through any type of irrigation system.

RECOMMENDATIONS FOR USE

ALFALFA (Hay and Seed): Grasshoppers — $\frac{1}{2}$ pt. per acre when crop is not more than 12" high. Aphids, Leafhoppers, Lygus Bugs, Alfalfa Weevils, Armyworms — 1 pt. per acre. **ALFALFA (Seed Only):** Lygus, Mites — 2 applications at 7-day intervals. Thrips — $\frac{1}{2}$ pt. per acre. Combine with recommended chemicals for residual control. By Air — Use at least 10 gallons water per acre.

BEANS (Dry or Field), SOYBEANS, LIMA BEANS and PEAS for Processing: Alfalfa Leafhopper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites, Leafhoppers, Mites, Leafhoppers — 1 pt. per acre in water to cover. 100 gals. per acre is preferable. Leafhoppers, Stink Bugs — $\frac{1}{2}$ pt. per acre in water to cover.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE, COLLARD: Imported Cabbageworm, Diamond-Back Caterpillars, Aphids — 1 pt. per acre. Looper — 2 pt. per acre. May be used 2 days before harvest, except do not apply to kale and collards within 4 days of harvest.

CELERY: Aphids, Leafhoppers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals. water per acre on plants less than 12" high. On larger plants use $\frac{1}{2}$ pt. in 100 to 150 gals. water per acre.

COTTON: Lygus — $\frac{1}{2}$ to 1 pt. per acre by air or ground. Leaf Perforator, Fleahoppers, Armyworms, Leafhoppers, Looper and For Suppression of Mites — 1 pt. per acre by air or ground. Apply at 7 to 10 day intervals while pests persist. **FOR HEAVY POPULATION OF LISTED INSECTS:** — Use $\frac{1}{2}$ pt. per acre by air or ground in control programs with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Looper, Thrips, Armyworms — 1 pt. per acre in water to cover. Pickleworms, Leaf Miners — $\frac{1}{2}$ pt. per acre in water to cover. By Air — Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miner, Whitefly — 1 pt. per acre. May be used up to 1 day before harvest.

GRAPES: Drosophila, Leafhoppers — $\frac{1}{2}$ to $\frac{3}{4}$ pt. per 100 gals. water (100 to 230 gals. diluted spray per acre depending on size of vines). Use only $\frac{1}{2}$ pt. per 100 gals. water on Muscat varieties.

CAUTION: Postbloom applications can cause fruit russetting. Do not use on Itaka varieties as injury may occur. Do not apply within 24 hours following harvest. Do not apply within 1 day of harvest.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm — 1 pt. per acre.

MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH: Looper, Armyworms, Leafhoppers, Spider Mites, Leaf Miners — 1 to 2 pts. per acre (200 gals. diluted spray per acre). May be used 1 day before harvest. Do not apply when temperature is over 80°F.

33252

BEST AVAILABLE COPY

8 of 12

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale Crawlers, Mites, Citrus Curculionids, Leaf Roller (Tortrix) — 1 pt. to 100 gals. water. Thrips (Bloom) — 1/2 pt. per 100 gals. water. Use up to 7 days of harvest. (Calif. and Ariz. — Do not apply within 1 day of harvest.) Bees and other bees present for 24 hours following treatment.

PEACHES (Western States Only): Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only) — 1/2 pt. to 100 gals. water. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 80°F. Do not use outside of 24-hour following treatment.

RICE: Grasshoppers — 1/2 to 3/4 pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. May be used 2 days before harvest.

SPINACH, CHARD, TURNIP GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals. water per acre on plants less than 1/2 grown. On larger plants, use 1 1/2 pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest) — 1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider Mites, Leafhoppers — 1 pt. per acre by air or ground. By Air — Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworms — 1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre. Flea Beetles — 1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as necessary.

TOMATOES (Field): Fruit Flies (Drosophila) — 1 pt. in no less than 40 gals. water per acre with ground sprayer. 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. (Field and Greenhouse): Tomato Fruit Worm, Hornworms, Leaf Miners, Flea Beetles, Mites — 1 pt. in 100 gals. water. Spray plants thoroughly. For Field — Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 80°F. May be used 1 day before harvest.

WALNUTS: Aphids — 1/2 to 1 pt. to 100 gals. water. Use up to 10 days of harvest.

GRASSHOPPERS: Rangeland, Field Areas (crops listed on label) and Pastures including dairy pastures; Range Caterpillars on Rangeland and Pastures — 1/2 to 3/4 pt. per acre by air or ground. Young Grasshoppers — 1/2 pt. per acre by air or ground. Armyworms (except rice) — 1 pt. per acre by air or ground. By Air — Use in 1 to 5 gals. water per acre. Apply when hoppers hatch is completed. Repeat as necessary. Animals may be present during treatment. See crop recommendations for use limitations near harvest. Forage plants on pastures or field areas should not be cut for hay to be sold or shipped interstate within 4 days of application.

GREENHOUSE, VAPOR TREATMENT OR ROSES: OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs — 1 fluid ounce per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes (or in open pans on hot plates). Apply continuously or to intermittent sections of pipe on each side and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 180°F. In steam houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes, wear goggles and avoid breathing vapor. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment before harvest. Spider Mites (Resistant and Non-resistant) — Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Adult Whiteflies — Make 2 applications at 7-day intervals. Repeat as necessary. Aphids — Apply when aphids appear. Repeat as necessary. WARNING — Avoid over-treatment and direct application to plants as injury may result. White Butterfly and Golden Pheasant roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor.

FRUIT FLIES (Drosophila sp.) In and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills — Use 2 ounces in 2 1/2 gals. water (5 pints to 100 gals.) as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as necessary. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying.

ADULT MOSQUITO, GNAT, STABLE FLY (Dog Flies) and HOUSE FLY CONTROL: Residential Areas, Crops Listed on Label, Tidal Marshes, Swamps, Woodland, Livestock Pastures, Feed Lots and Pastures including Dairy Cattle — Consult your State Fish & Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to tidal or marsh waters which are important shrimp producing areas. AIRCRAFT: East — Apply 0.8 to 4 fluid ozs. per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. West — Apply 1.8 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 qts. of diluted spray per acre. MIST OR COLD FOG — Use 3 to 5 qts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

CORRALS, ADJACENT PASTURES, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitoes — Airplane Application — Use 1.6 to 4 fl. ozs. per acre. Dilute 1 to 2 1/2 pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

CLOVER MITES, ROACHES, EARWIGS and LEAFHOPPERS: Outside Dwellings and in Latrines — Use 1 fl. oz. in 3 gals. water. Apply to turf and to soil surfaces around flowers, shrubs and trees for general pest cleanup.

BEST AVAILABLE COPY

COPY B**Personal Protective Equipment****Applicators and other handlers must wear:**

- A) Coveralls over long sleeved shirt and long pants.
- B) Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber \geq 14 mils, or Nitrile Rubber \geq 14 mils. + VITON
- C) Chemical-resistant footwear plus socks.
- D) Protective eyewear.
- E) Chemical-resistant headgear for overhead exposure.
- F) Chemical-resistant apron when cleaning equipment, mixing or loading.
- G) A respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

BEST AVAILABLE COPY

COPY D

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURE USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontaminations, notification, and emergency assistance. It also contains specific instructions, and exceptions pertaining to ~~X~~ protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

missing language

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPF required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- A) Coveralls over long sleeved shirt and long pants
- B) Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber ≥ 14 mils, or Nitrile Rubber ≥ 14 mils. + VITON
- C) Chemical-resistant footwear plus socks
- D) Protective eyewear
- E) Chemical-resistant headgear for overhead exposure.

COPY F**Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA Chemical resistance category selection chart.

Barrier Laminate

Butyl Rubber ≥ 14 mils

Nitrile Rubber ≥ 14 mils + 150M

COPY 1

^a
Si usted no entiende la etiqueta,
busque alguien para que se la
explique a usted en detalle (If
you do not understand the label
find someone to explain it to
you in detail).