

 <p>United States Environmental Protection Agency Washington, DC 20460</p>	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number <h1 style="margin:0;">222709</h1>
---	---	--

Application for Pesticide - Section I

1. Company/Product Number 2935-284	2. EPA Product Manager R. Forrest	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DIBROM 8 SPRAY	PM# 14	
5. Name and Address of Applicant (Include ZIP Code) WILBUR-ELLIS COMPANY 191 W. Shaw Avenue, Suite 107 Fresno, CA 93704 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input checked="" type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION APR 18 1996
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	


Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of deleted uses: delete cucumbers, rice, Turnip greens, tobacco and tomatoes. Add restriction against use of product on alfalfa in Utah.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name JOANNE C. STRIEBICH	Title Registration and Labeling Specialist	Telephone No. (include Area Code) (209) 226-1934
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, both under applicable law.		6. Date Application Received (Stamped)
2. Signature 		
3. Title Registration and Labeling Specialist		
4. Typed Name JOANNE C. STRIEBICH		5. Date March 12, 1996



DIBROM 8 SPRAY

NALED INSECTICIDE
ORGANOPHOSPHATE

ACTIVE INGREDIENTS:	By Wt.
Naled	58%
INERT INGREDIENTS:	42%
Xylene Range Aromatic Solvent	20%
1,2-Dibromo-2,2-dichloroethyl dimethyl phosphate	100%
Contains 8 pounds technical NALED. Equivalent to 7.2 pounds NALED per gallon.	

EPA Reg. No. 2935-284-ZA

EPA Est. No. 5481-CA-1

KEEP OUT OF REACH OF CHILDREN DANGER — PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

In Case of Contact — Flush skin or eyes with plenty of water; for eyes, get medical attention.

If Swallowed — Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

NOTE TO PHYSICIAN: Naled is a cholinesterase inhibitor. Atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine. Mucosal damage may occur, and if gastric lavage is indicated it should be performed cautiously.

REFER TO SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS: 1 GALLON

WILBUR-ELLIS COMPANY

P.O. Box 16458 • Fresno, CA 93755

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

ENTRY RESTRICTIONS: Outdoors and Non-Vapor Treatments in Greenhouses: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. **Vapor Treatment in Greenhouses:** Entry by any person — other than a correctly trained and equipped handler who is performing a WPS-defined handling task — is PROHIBITED in the entire enclosed area from the start of application until one of the WPS ventilation criteria has been met. After ventilation is completed, do not enter or allow worker entry into the entire enclosed area during the restricted-entry interval (REI) of 24 hours, except as allowed by the WPS.

PPE for Entry During the Restricted Period: PPE for early entry to treated areas (1) outdoors, (2) in non-vapor-treated greenhouses, or (3) in vapor-treated greenhouses after ventilation is complete — that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: A) Coveralls over long-sleeved shirt and long pants; B) Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, or Viton; C) Chemical-resistant footwear plus socks; D) Protective eyewear; E) Chemical-resistant headgear for overhead exposure.

PPE for entry by handlers into vapor-treated greenhouses before WPS ventilation criteria have been met is listed in the "Hazards to Humans and Domestic Animals" section of this labeling.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter treated areas without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

1. **Prohibitions:** Do not contaminate water, food or feed by storage or disposal.
2. **Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.
3. **Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. **Other:** Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

USE ONLY ACCORDING TO ATTACHED DIRECTIONS

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

DIBROM is a registered trademark of Valent U.S.A. for naled insecticide.
Logo TM of WILBUR-ELLIS Company
IDEAS TO GROW WITH is a registered trademark of WILBUR-ELLIS Company

F-194



DUMONT
PRINTING

(209) 485-6311
FAX: (209) 485-6357

Ink colors _____
Stock _____
Quantity _____ Ship date _____

please
READ YOUR PROOF
CAREFULLY!

1 GALLON
F-194
EPA Est. No. 5481-CA-1

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMAN AND DOMESTIC ANIMALS**

DANGER

Corrosive, causes eye and skin damage. May be fatal if swallowed. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA Chemical resistance category selection chart.

Applicators and other handlers must wear: A) Coveralls over long sleeved shirt and long pants; B) Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber or Viton; C) Chemical-resistant footwear plus socks; D) Protective eyewear; E) Chemical-resistant headgear for overhead exposure; F) Chemical-resistant apron when cleaning equipment, mixing or loading; G) A respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

Exception: When applying to hot pipes in greenhouses or entering vapor-treated greenhouses during application or before ventilation is complete, the protective eyewear must be goggles, unless a full-faced respirator is worn.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not spray on Nectar trees, ornamental Cherries, Liquidamber or Chrysanthemums as injury may occur.

Do not store diluted spray. Make new dilution for each use.

PHYSICAL OR CHEMICAL HAZARDS

Wash metal equipment thoroughly after use to avoid corrosion. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

48032



DIBROM 8 SPRAY

NALED INSECTICIDE
ORGANOPHOSPHATE

ACTIVE INGREDIENTS		By Wt
Naled		58%
INERT INGREDIENTS		42%
Xylene Range Aromatic Solvent	20%	
1,2-dibromo-2,2-dichloroethyl dimethyl phosphate		100%
Contains 8 pounds technical NALED Equivalent to 7.2 pounds NALED per gallon		
EPA Reg No 2935-284 ZA		

KEEP OUT OF REACH OF CHILDREN DANGER — PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

NOT FOR USE OR STORAGE IN OR AROUND HOME

Observe Precautions on Container Label

DIRECTIONS FOR USE

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED

GROUND APPLICATION Vegetable and Field Crops — 30 to 250 gals Cotton — 3 to 20 gals Deciduous Fruit Crops — 60 to 800 gals Citrus — 400 to 3 000 gals (On Young Fruit Plantings — Use a minimum of 15 gals water per acre)

Where Recommended for AIR APPLICATION Field Crops — 3 to 10 gallons

Apply to the foliage of insects before kill of insect and mites, ticks and larvae and to the foliage of plants.

Do not apply within 7 days of harvest. Do not apply with high winds or rain. Do not apply to wet foliage. Do not apply to plants with high humidity or to plants such as lime or border x.

MIXING DIRECTIONS Add to spray tank when to full weight to uniform and maintain agitation until spraying completed.

Do not apply product through any type of irrigation system.

RECOMMENDATIONS FOR USE

NOT FOR
USE IN
UTAH

ALFALFA (Hay and Seed) Grasshoppers — 1 pt per acre when crop is not more than 12" high. Aphids, Loopers, Leafhoppers, Lygus Bugs, Alfalfa Webworms, Armyworms — 1 pt per acre. ALFALFA (Seed Only) Lygus Mites — 2 applications at 7 day intervals. Thrips — 1/2 pt per acre. Corn borer, etc. recommended chemicals for residual control. By Air — Use at least 10 gallon water per acre.

BEANS (Dry or Field) SOYBEANS LIMA BEANS and PEAS for Processing Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner, GREEN BEANS Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers — 1 pt per acre. Water to cover 100 gals per acre. Large Loopers, Stink Bugs — 1 pt per acre in water to cover.

BROCCOLI CABBAGE CAULIFLOWER BRUSSELS SPROUTS KALE COLLARDS Imported Cabbageworm, Diamond Back Caterpillars, Aphids — 1 pt per acre. Looper — 2 pt per acre. May 1st to 1st of July. Use before the caterpillars appear. Use at least 10 gal water per acre.

CELERY Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners. Use at least 10 gal water per acre. Do not apply to plants less than 10 gal water per acre.

COTTON Lygus — 1 pt per acre. Yellow Ground Squirrel (Perforating) Leafhoppers, Armyworms, Bollworms, Loopers and For Spreading of Mite. Do not apply to cotton bolls. Apply 10 days before harvest. **FOR HEAVY POPULATION OF LISTED INSECTS** Use at least 10 gal water per acre. Do not apply to plants less than 10 gal water per acre.

CUCUMBERS SUMMER SQUASH Cucumber Beetles, Aphids, Spider Mites, Looper, Thrips, Armyworms — 1 pt per acre. Water to cover. Pickleworms, Leaf Miners — 1 pt per acre. Water to cover. By Air — Use a minimum of 10 gal water per acre. May be used pre-harvest.

EGGPLANT PEPPERS Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly — 1 pt per acre. May be used up to 10 days before harvest.

GRAPES Drosophila, Leafhoppers — 1 pt per 100 sq ft (100 to 23) of diluted spray. Do not apply to clusters. Use only 1 pt per 10 gal water. Do not apply to clusters. **CAUTION** Do not apply to clusters. Do not apply to clusters. Do not apply to clusters.

HOPS Aphids, Spider Mites, Caterpillars — 1 pt per acre.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION: Vegetable and Field Crops — 30 to 250 gals. Cotton — 3 to 20 gals. Deciduous Fruit Crops — 60 to 800 gals. Citrus — 400 to 3,000 gals. (On Young Fruit Plantings — Use a minimum of 15 gals. water per acre.)

Where Recommended for AIR APPLICATION: Field Crops — 3 to 10 gallons.

A short residual contact insecticide for kill of insect and mite adults and larvae present at the time of application.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when ¼ to ½ full with agitator running and maintain agitation until spraying is completed.

Do not apply this product through any type of irrigation system.

RECOMMENDATIONS FOR USE

NOT
FOR
USE
IN
UTAH

ALFALFA (Hay and Seed): Grasshoppers — ¾ pt. per acre when crop is not more than 12" high. **Aphids, Loopers, Leafhoppers, Lygus Bugs, Alfalfa Webworms, Armyworms** — 1 pt. per acre. **ALFALFA (Seed Only): Lygus, Mites** — 2 applications at 7-day intervals. **Thrips** — ½ pt. per acre. Combine with recommended chemicals for residual control. **By Air** — Use at least 10 gallons water per acre.

BEANS (Dry or Field), SOYBEANS, LIMA BEANS and PEAS for Processing: Alfalfa Loopers, Aphids, Spider Mites, Lygus, Leaf Miner; GREEN BEANS: Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers — 1 pt. per acre in water to cover. 100 gals. per acre is preferable. **Large Loopers, Stink Bugs** — 1½ pts. per acre in water to cover.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE, COLLARDS: Imported Cabbageworm, Diamond-Back Caterpillars, Aphids — 1 pt. per acre. **Looper** — 2 pts. per acre. May be used 2 days before harvest, except do not apply to kale and collards within 4 days of harvest.

CELERY: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals. water per acre on plants less than ¾ grown. On larger plants use 1½ pts. in 100 to 150 gals. water per acre.

COTTON: Lygus — ½ to 1 pt. per acre by air or ground. **Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and For Suppression of Mites** — 1 pt. per acre by air or ground. Apply at 7 to 10 day intervals while pests persist. **FOR HEAVY POPULATION OF LISTED INSECTS** — Use ½ pt. per acre by air or ground in control programs with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms — 1 pt. per acre in water to cover. **Pickleworms, Leaf Miners** — 1½ pts. per acre in water to cover. **By Air** — Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly — 1 pt. per acre. May be used up to 1 day before harvest.

GRAPES: Drosophila, Leafhoppers — ¼ to ¾ pt. per 100 gals. water (100 to 230 gals. diluted spray per acre depending on size of vines). Use only ½ pt. per 100 gals. water on Muscat varieties. **CAUTION:** Postbloom applications can cause fruit russetting. Do not use on Italia varieties as fruit injury may occur. Do not apply within 1 day of harvest.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm — 1 pt. per acre.

MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH: Loopers, Armyworms, Leafhoppers, Spider Mites, Leaf Miners — 1 to 2 pts. per acre (200 gals. diluted spray per acre). May be used 1 day before harvest. Do not apply when temperature is over 90°F.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale Crawlers, Mites, Citrus Cutworms, Leaf Roller (Tortrix) — 1 pt. to 100 gals. water, Thrips (Bloom) — 1/2 pt. per 100 gals. water. Use up to 7 days of harvest. (Calif. and Ariz. — Do not apply within 1 day of harvest.)

PEACHES (Western States Only): Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only) — 3/4 pt. to 100 gals. water. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90°F.

~~RICE: Grasshoppers — 1/2 to 3/4 pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. May be used 2 days before harvest.~~

SPINACH, CHARD, TURNIP-GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals. water per acre on plants less than 3/4 grown. On larger plants, use 1 1/2 pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tler, Spittlebugs (Northwest) — 1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider Mites, Leafhoppers — 1 pt. per acre by air or ground. By Air — Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

~~TOBACCO: Hornworm — 1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre. Flea Beetles — 1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as necessary.~~

~~TOMATOES (Field): Fruit Flies (Drosophila) — 1 pt. in no less than 40 gals. water per acre with ground sprayer, 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. (Field and Greenhouse): Tomato Fruit Worm, Hornworms, Leaf Miners, Flea Beetles, Mites — 1 pt. in 100 gals. water. Spray plants thoroughly. For Field — Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 90°F. May be used 1 day before harvest.~~

WALNUTS: Aphids — 1/2 to 1 pt. to 100 gals. water. Use up to 10 days of harvest.

GRASSHOPPERS: Rangeland, Field Areas (crops listed on label) and Pastures including dairy pastures; Range Caterpillars on Rangeland and Pastures — 1/2 to 3/4 pt. per acre by air or ground. Young Grasshoppers — 1/2 pt. per acre by air or ground. Armyworms (except rice) — 1 pt. per acre by air or ground. By Air — Use in 1 to 5 gals. water per acre. Apply when hoppers hatch is completed. Repeat as necessary. Animals may be present during treatment. See crop recommendations for use limitations near harvest. Forage plants on pastures or field areas should not be cut for hay to be sold or shipped interstate within 4 days of application.

~~GREENHOUSE, VAPOR TREATMENT OF ROSES; OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs — 1 fluid ounce per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes (or in open pans on hot plates). Apply continuously or to intermittent sections of pipe on each side and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160°F. In steam houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment before harvest. Spider Mites (Resistant and Non-resistant) — Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Adult Whiteflies — Make 2 applications at 7-day intervals. Repeat as necessary. Aphids — Apply when aphids appear. Repeat as necessary. WARNING — Avoid overtreatment and direct application to plants as injury may result. White Butterfly and Golden Rapture roses, Pink Champagne Chrysanthemums, Green Wondering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor.~~

FRUIT FLIES (Drosophila sp.) In and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills — Use 2 ounces in 2 1/2 gals. water (5 pints to 100 gals.) as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as necessary. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying.

ADULT MOSQUITO, GNAT, STABLE FLY (Dog Flies) and HOUSE FLY CONTROL: Residential Areas, Crops Listed on Label, Tidal Marshes, Swamps, Woodland, Livestock Pastures, Feed Lots and Pastures Including Dairy Cattle — Consult your State Fish & Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to tidal or marsh waters which are important shrimp producing areas. AIRCRAFT: East — Apply 0.8 to 4 fluid ozs. per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. West — Apply 1.6 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 qts. of diluted spray per acre. MIST OR COLD FOG — Use 3 to 5 qts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

CORRALS, ADJACENT PASTURES, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitos — Airplane Application — Use 1.6 to 4 fl. ozs. per acre. Dilute 1 to 2 1/2 pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

CLOVER MITES, ROACHES, EARWIGS and LEAFHOPPERS: Outside Dwellings and In Lawns — Use 1 fl. oz. in 3 gals. water. Apply to turf and to soil surfaces around flowers, shrubs and trees for general pest cleanup.