

# THIODAN® S SULFUR 25 DUST

This product formulated for application by ground or air.

## RECOMMENDATIONS FOR USE:

**ARTICHOKES:** Use 20 lbs. per acre to control ARTICHOKE APHID and GREEN PEACH APHID. Repeat as needed for control. Do not apply within 7 days of harvest.

**BEANS:** Use 20 lbs. per acre to control MEXICAN BEAN BEETLE, BEAN APHID and WESTERN BEAN CUTWORM. Do not apply more than 3 times per season. Do not feed treated threshings to livestock or allow livestock to graze in treated fields. Do not use on Lima beans that are to be removed from the field for processing. Do not apply within 3 days of harvest.

**BRASSICA, CABBAGE and CAULIFLOWER:** Use 15 lbs. per acre to control CABBAGE LOOPER, IMPORTED CABBAGE WORM, DIAMOND-BACK MOTH LARVAE, CROSS-STRIPED CABBAGE WORM, FLEA BEETLE, HARLEQUIN BUG and STINK BUGS. Repeat as necessary to maintain control. Do not apply on Cauliflower within 14 days of harvest. Do not apply on Broccoli or Cabbage within 7 days of harvest.

**COTTON:** Use 30 lbs. per acre to control COTTON STINK BUGS, LYGUS BUGS, BOLLWORM and CABBAGE LOOPER. Repeat as necessary to maintain control. Do not apply after bolls begin to open. Do not graze meat or dairy animals in treated fields.

**EGGPLANTS:** Use 10 lbs. per acre to control APHIDS, FLEA BEETLES, COLORADO POTATO BEETLE and WHITEFLY. Repeat as necessary to maintain control. Do not apply within 1 day of harvest.

**GRAPES:** Use 25 to 30 lbs. per acre to control GRAPE LEAFHOPPER. Do not apply within 7 days of harvest. Do not use on Concord variety as injury is likely to occur.


**LETUCE (Seed):** Use 20 lbs. per acre to control CABBAGE LOOPER, GREEN PEACH APHID, DIAMOND-BACK MOTH LARVAE & IMPORTED CABBAGE WORM. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove outside leaves at harvest. Do not feed crop waste to livestock.

**PEAS (Seed):** Use 20 to 24 lbs. per acre to control PEA APHID and PEA WEEVIL. Do not feed treated vines or threshings to dairy or meat animals.

**PEPPERS:** Use 20 lbs. per acre to control FLEA BEETLE and GREEN PEACH APHID. Do not apply within 1 day of harvest.

**POTATOES:** Use 20 lbs. per acre to control FLEA BEETLE, COLORADO POTATO BEETLE, LEAFHOPPERS, APHIDS, POTATO TUBERWORM, LEAF-FOOTED PLANT BUG, SOUTHERN ARMYWORM, GREEN STINK BUG, POTATO PSYLLID and EUROPEAN CORN BORER. Apply as needed to harvest.

**SQUASH (Summer and Winter):** Use 15 to 20 lbs. per acre to control APHIDS, CUCUMBER BEETLES, SQUASH VINE BORERS and STRIPED FLEA BEETLES. For SQUASH VINE BORER, apply to flower buds, stems and vines as soon as moths appear. Do not apply more than 20 lbs. per acre. May be applied up to day of harvest.



## ACTIVE INGREDIENTS:

Sulfur	5.00%
Titanium Dioxide	21.00%
Inert Ingredients	73.00%
	100.00%

\* Hexachlorhexahydromethano-2, 4, 3-benzodioxathiepin oxide

EPA Reg. No. 2935-305-AA

## KEEP OUT OF REACH OF CHILDREN!

## CAUTION:

Harmful If Swallowed, Inhaled or Absorbed Through Skin. Avoid inhalation and contact with skin, eyes or clothing. Avoid contamination of feed and foodstuffs. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas within 24 hours of treatment should wear protective clothing.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

## NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or flames.

Do not plant root crops other than carrots, potatoes, sweet potatoes and sugar beets as a follow-up crop.

Thiodan is a registered trademark of Canadian Hoechst Limited.

## RECOMMENDATIONS FOR USE: (Continued)

**STRAWBERRIES:** Use 20 lbs. per acre to control STRAWBERRY APHID and MEADOW SPITTLEBUG. Do not apply within 4 days of harvest. Do not apply at intervals of less than 15 days or more than twice during any 35-day period.

**Tomatoes:** Use 20 lbs. per acre to control COLORADO POTATO BEETLE, FLEA BEETLE, HORNWORM, TOMATO FRUITWORM, CABBAGE LOOPER, GREEN PEACH APHIDS and WHITEFLY. Repeat as necessary to maintain control. Do not apply within 1 day of picking.

**WALNUTS:** Use 40 lbs. per acre to control WALNUT APHIDS. Repeat as required to maintain control. Do not apply after husk split. Do not graze livestock in treated groves.