

THIODAN® 5 DUST

DIRECTIONS FOR USE: This material may be applied by ground or air.

ARTICHOKES: Use 20 lbs. per acre to control ARTICHOKE APHID and GREEN PEACH APHID. Repeat as needed for control. Do not apply within 7 days of harvest.

BEANS: Use 20 lbs. per acre to control MEXICAN BEAN BEETLE, BEAN APHID and WESTERN BEAN CUTWORM. Do not apply more than 3 times per season. Do not feed treated threshings to livestock or allow livestock to graze in treated fields. Do not use on lima beans that are to be removed from the field for processing. Do not apply within 3 days of harvest.

BROCCOLI, CABBAGE and CAULIFLOWER: Use 15 lbs. per acre to control CABBAGE LOOPER, IMPORTED CABBAGE WORM, DIAMOND-BACK MOTH LARVAE, CROSS-STRIPED CABBAGE WORM, FLEA BEETLE, HAR-LEQUIN BUG and STINK BUGS. Repeat as necessary to maintain control. Do not apply on Cauliflower within 14 days of harvest. Do not apply on Broccoli or Cabbage within 7 days of harvest.

COTTON: Use 30 lbs. per acre to control COTTON STINK BUGS, LYGUS BUGS, BOLLWORM and CABBAGE LOOPER. Repeat as necessary to maintain control. Do not apply after bolls begin to open. Do not graze meat or dairy animals in treated fields.

CUCUMBERS, MELONS, PUMPKINS, SUMMER and WINTER SQUASH: Use 20 lbs. per acre to control APHIDS and CUCUMBER BEETLES. Repeat as necessary to maintain control.

EGGPLANTS: Use 12 lbs. per acre to control APHIDS, FLEA BEETLES, COLORADO POTATO BEETLE and WHITEFLY. Repeat as necessary to maintain control. Do not apply within 1 day of harvest.

GRAPES: Use 25 to 30 lbs. per acre to control GRAPE LEAFHOPPER. Do not apply within 7 days of harvest. Do not use on Concord variety as injury is likely to occur.

LETTUCE (Head): Use 20 lbs. per acre to control CABBAGE LOOPER, GREEN PEACH APHID, DIAMOND-BACK MOTH LARVAE & IMPORTED CABBAGE WORM. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove outside leaves at harvest. Do not feed crop waste to livestock.

PEAS (Seed): Use 20 to 24 lbs. per acre to control PEA APHID and PEA WEEVIL. Do not feed treated vines or threshings to dairy or meat animals.

POTATOES: Use 20 lbs. per acre to control FLEA BEETLE, COLORADO POTATO BEETLE, LEAFHOPPERS, APHIDS, POTATO TUBERWORM, LEAF-FOOTED PLANT BUG, SOUTHERN ARMYWORM, GREEN STINK BUG, POTATO PSYLLID and EUROPEAN CORN BORER. Apply as needed to harvest.

PEPPERS: Use 20 lbs. per acre to control FLEA BEETLE and GREEN PEACH APHID. Do not apply within 1 day of harvest.

STRAWBERRIES: Use 20 lbs. per acre to control STRAWBERRY APHID and MEADOW SPITTLEBUG. Do not apply within 4 days of harvest. Do not apply at intervals of less than 15 days or more than twice during any 35-day period.

F-774 EPA Reg. 2935-CA-1

ACCEPTED
2935-306

AUG 27 1974

ACTIVE INGREDIENT:

Endosulfan® 5.00%

INERT INGREDIENTS: 95.00%

100.00%

* Hexachloro-Hexahydro-methano-2, 4, 3-benzodioxathiepin oxide

EPA Reg. No. 2935-306-AA

KEEP OUT OF REACH OF CHILDREN !

CAUTION: Harmful if Swallowed, Inhaled or Absorbed Through Skin.

Avoid contact with skin, eyes, or clothing. Do not breathe dust.

Avoid contamination of feed and foodstuffs. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Workers entering treated areas within 24 hours of treatment should wear protective clothing.

This product is toxic to bees and should not be applied when bees are visiting the area.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DO NOT PLANT ROOT CROPS OTHER THAN CARROTS, POTATOES, SUGAR BEETS AND SWEET POTATOES AS FOLLOW UP CROPS.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes. IN CALIF.: Dispose of in accordance with state and local regulations.

®Thiodan is a registered trademark of Farbwerke Hoechst A. G.

U. S. Patent No. 2,799,685

RECOMMENDATIONS FOR USE: (Continued)

TOMATOES: Use 20 lbs. per acre to control COLORADO POTATO BEETLE, FLEA BEETLE, HORNWORM, TOMATO FRUITWORM, CABBAGE LOOPER, APHIDS and WHITEFLY. Repeat as necessary to maintain control. Do not apply within 1 day of picking.

WALNUTS: Use 40 lbs. per acre to control WALNUT APHIDS. Repeat as required to maintain control. Do not apply after husk split. Do not graze livestock in treated groves.