

THIODAN 5 SULFUR 50 DUST

APPLICATOR: Use 20 lbs. per acre to control **APRICOT TWIG BORER**. Repeat as needed for control. Do not use on apricots.

APPLES: Use 20 lbs. per acre to control **AMERICAN BARK BEETLE**. Do not apply after pods begin to form. Do not use on apples which are destined for dairy or meat animals.

BRASSICAS: Use 20 lbs. per acre to control **CABBAGE LOOPER, GREEN PEACH APHID, DIAMOND-BACK MOTH LARVAE & WEEVIL, CABBAGE WORM, CABBAGE ROOT MOTH, CABBAGE PUPA and WEEVIL**. Repeat as necessary to maintain control. Do not apply on Cabbageworm after white parts start to form. Do not apply on Broccoli or Cabbage within 7 days of harvest.

COTTON: Use 30 lbs. per acre to control **COTTON STINK BUG, LYGUS BUGS, BOLLWORM** and **CABBAGE BEE**. Repeat as necessary to maintain control. Do not apply after bolls begin to open. Do not use on areas used for dairy animals in treated fields.

CUCUMBERS: Use 10 lbs. per acre to control **BEANS, FLEA BEETLE, COLORADO POTATO BEETLE and WHITEFLY**. Repeat as necessary to maintain control. Do not apply within 1 day of harvest.

CUCUMBERS: Use 25 to 30 lbs. per acre to control **LEAF MINERS**. Do not apply within 7 days of harvest. Do not use on Cucumbers variety as injury is likely to occur.

CUCUMBERS: Use 20 lbs. per acre to control **CABBAGE LOOPER, GREEN PEACH APHID, DIAMOND-BACK MOTH LARVAE & WEEVIL, CABBAGE WORM**. Do not use more than 3 applications during season. Do not apply within 7 days of harvest. Remove outside leaves at harvest. Do not feed crop waste to livestock.

PEAS: Use 20 to 25 lbs. per acre to control **BEA ARMS and BEA WEEVIL**. Do not feed treated vines or freshings to dairy or meat animals.

POTATOES: Use 30 to 55 lbs. per acre to control **FLEA BEETLE, COLORADO POTATO BEETLE, LEAFHOPPERS, APHIDS, POTATO TUBER MOTH, LEAF-ROOTED PLANT BUG, SOUTHERN ARMYWORM, GREEN STINK BUG, POTATO FOLIAR and EUROPEAN CORN BORER**. Apply as needed to harvest.

POTATOES: Use 20 lbs. per acre to control **FLEA BEETLE and GREEN PEACH APHID**. Do not apply within 7 days of harvest.

5.00%
50.00%
65.00%
100.00%

1-methyl-2-(2,4,6-trichlorophenyl)hydrazine-2,4,3-benzodioxathiepin oxide

CAUTION: See Reg. No. 2935 - 307. **Harmful if Swallowed.** Avoid inhalation or contact with skin. Avoid contamination of feed and feedstuffs. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. Workers entering treated areas the day of application should wear protective clothing. Toxic to fish and wildlife. Do not contaminate any body of water for application to any area not

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THE LABEL UNLESS INTENDED USE IS RECOMMENDED BY LOCAL AGRICULTURAL OFFICIALS.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM AREA BEING TREATED.
 ®Thiodan is a registered trademark of Farbwerke Hoechst A.G.

U. S. Patent No. 2,799,485

RECOMMENDATIONS FOR USE: (Continued)

STRAWBERRIES: Use 20 lbs. per acre to control **STRAWBERRY APHID & AMERICAN SPITTLERUG**. Do not apply within 4 days of harvest. Do not apply at intervals of less than 15 days or more than twice during any 35-day period.

TOMATOES: Use 20 lbs. per acre to control **COLORADO POTATO BEETLE, FLEA BEETLE, APHIDS and WHITEFLY**. Repeat as necessary to maintain control. Do not apply within 1 day of picking.

WALNUTS: Use 40 lbs. per acre to control **WALNUT APHIDS**. Repeat as required to maintain control. Do not apply after husk split.

F-664-O

ACCEPTED