

Best Available COPY

red-top

ACCEPTED
FEB 02 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

SUPERIOR SPRAY OIL

2935-311

ACTIVE INGREDIENT:	
Mineral Oil	99.0%
INERT INGREDIENT	1.0%
	100.0%
Unvolatilized Residue (ASTM D-483)	93%

Petroleum Oil Classification: Unclassified
EPA Reg. No. 2935-00311AA

CAUTION: KEEP OUT OF REACH OF CHILDREN!

HARMFUL IF SWALLOWED. DO NOT INDUCE VOMITING. CALL A PHYSICIAN IMMEDIATELY.

CAUTION

- READ ENTIRE LABEL. Use strictly in accordance with label cautions, warnings, and directions; and in conformity with Federal and State regulations.
- DO NOT APPLY spray oil without consulting your State Agricultural Extension Service.
- DO NOT COMBINE WITH SULFUR SPRAYS and DO NOT APPLY spray oil within 30 days before or after an application of sulfur.
- DO NOT MIX with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service for specific compatibilities.
- DO NOT APPLY to trees weakened by disease, drought, drying winds, or high nitrogen applications.
- DO NOT USE Red-Top Superior Spray Oil when temperatures are above 90° Fahrenheit.
- AVOID breathing spray mist.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

DIRECTIONS FOR USE:

Dormant, delayed dormant or summer applications: Almonds, apricots, apples, cherries, peaches, pears, prunes. Apply 5-8 gallons oil per acre for control of European Red Mite, Brown Almond Mite, two-spotted Mite, Pacific Mite, Pear Psylla, McDaniel Mite, Strawberry Mite, Plum Nursery Mite, Willamette Mite, Pear Rust Mite, Peach Silver Mite, San Jose Scale, and Lecanium Scale.

Walnuts: Late spring application—apply 5-7 gallons oil per acre for control of San Jose Scale, Walnut Aphid.

Dilution Rate: Apply rates given in 50 2000 gallons water per acre.

Fill spray tank with 1/2 the required amount of water. Start agitator. Add total amount of spray oil. Continue operation of agitator while adding balance of water and while spraying.

Spray oil may be used as a carrier for appropriate Insecticides and Fungicides which are Registered for use on deciduous fruits and nuts and which can be applied to give a uniform spray. Consult your local Agricultural Extension Service.

GENERAL DIRECTIONS: Red-Top Superior Spray Oil is designed specifically for use on deciduous fruit trees.

Apply so as to give total coverage to all parts of tree. Thorough drenching of trunks, limbs, twigs and leaves is essential for maximum insect control.

Use only in spray machines with powerful positive agitation. Keep agitator running at all times during spray operations. Red-Top Superior Spray Oil is formulated as a "fast break" spray oil and will not stay in solution unless agitated constantly!

F-173-B



WILBUR-ELLIS COMPANY



FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

2935-311

2-2-73

10/1