


SEVIN® 10 DUST



RECOMMENDATIONS FOR USE:

ALFALFA and CLOVERS: Use 15 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, JAPANESE BEETLE, ALFALFA CATERPILLAR, ARMYWORMS, CORN EARWORMS and STINK BUGS.

APRICOTS, PEACHES, PLUMS and NECTARINES: Use 40 to 50 lbs. per acre by air or ground to control Twig Borer and Oriental Fruit Moth. Do not use within 3 days of harvest on Apricots and Nectarines or within 1 day on Peaches and Plums.

BEANS: Use 20 lbs. per acre to control MEXICAN BEAN BEETLE, LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, CORN EARWORMS, ARMYWORMS, VELVET BEAN CATERPILLAR, LYGUS and STINK BUGS. Do not feed treated vines to meat or dairy animals.

CORN: Use 20 lbs. per acre to control CORN EARWORM, FALL ARMYWORM, SAP BEETLES and JAPANESE BEETLE. Do not apply within 7 days of harvest for forage. For Corn Earworm apply to silks as soon as they appear. Make 3 to 4 applications of 2-3 day intervals. Reapply immediately after rain.

COTTON: Use 20 to 25 lbs. per acre to control THRIPS, FLEA BEETLE, FLEA HOPPER, LEAFWORM, BOLLWORM, FALL ARMYWORM, LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLAR, STINK BUGS and GRASSHOPPERS. Do not graze treated fields or feed cotton trash to dairy or meat animals.

GRAPES: Use 20 to 25 lbs. per acre to control GRAPE LEAF FOLDER, GRAPE LEAFHOPPER, GRAPE BERRY MOTH and JAPANESE BEETLE. No time limitation.

EGGPLANTS, PEPPERS and TOMATOES: Use 20 lbs. per acre to control TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES, LEAFHOPPER, TOMATO HORNWORM, EUROPEAN CORN BORER, STINK BUGS, COLORADO POTATO BEETLE and LYGUS BUGS. No time limitation.

POTATOES: Use 15 to 20 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO BEETLE. Apply to foliage only.

STRAWBERRIES: Use 20 lbs. per acre to control MEADOW SPITTLEBUG, and LEAFROLLER. Do not apply within 1 day of harvest.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate)	10%
INERT INGREDIENTS	90%
	100%

EPA Reg. No. 2935-320-AA

KEEP OUT OF REACH OF CHILDREN!

CAUTION: Harmful if Inhaled or Swallowed. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing dust. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. To protect fish and wildlife, do not contaminate any body of water with this material.

Note For Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

Sevin is a trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

F-972-320-B