

# SEVIN<sup>®</sup> 10 DUST

## PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS:** Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid breathing dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

## STATEMENT OF PRACTICAL TREATMENT

**If swallowed:** Call a physician immediately. Give a glass or two of water and induce vomiting by touching back of throat with finger. It is preferable to induce vomiting under medical supervision or to use gastric lavage with a cuffed endotracheal tube because of aspiration hazard. Remove victim immediately to emergency treatment facilities.

**If on skin:** Remove by washing with soap and water. Get medical attention.

**If in eyes:** Flush the eyes with clean water for 10 minutes. Get medical attention.

**If inhaled:** Remove victim to fresh air. Transport immediately to emergency treatment facility.

**ANTIDOTE:** Note to Physician. Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

## ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms. To protect fish and wildlife, do not contaminate any body of water with this material.

**DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.**

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service.

## DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## NOT FOR HOME USE

## STORAGE AND DISPOSAL

- 1. Prohibitions** — Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2. Pesticide Disposal** — Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.
- 3. Container Disposal** — Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.
- 4. General:** Consult Federal, State or Local disposal authorities for approved alternative procedures.

**ALFALFA and CLOVERS:** Use 15 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPERS, JAPANESE BEETLES, ALFALFA CATERPILLARS, ARMYWORMS, CORN EARWORMS and STINK BUGS

SV10 4

## ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-Methylcarbamate)

10%

## INERT INGREDIENTS

90%

100%

\* Sevin is a trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

EPA Reg No 2935-320

EPA Est. 2935-CA-1

**KEEP OUT OF REACH OF CHILDREN!**  
**CAUTION:**

SEE PRECAUTIONARY STATEMENTS IN LEFT PANEL

## DIRECTIONS FOR USE (Continued):

**APRICOTS, PEACHES, PLUMS and NECTARINES:** Use 40 to 50 lbs. per acre by air or ground to control TWIG BORER and ORIENTAL FRUIT MOTH. Do not use within 3 days of harvest on Apricots and Nectarines or within 1 day on Peaches and Plums.

**BEANS:** Use 20 lbs. per acre to control MEXICAN BEAN BEETLES, LEAFHOPPERS, BEAN LEAF BEETLES, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, CORN EARWORMS, ARMYWORMS, VELVET BEAN CATERPILLARS, LYGUS and STINK BUGS. Do not feed treated vines to meat or dairy animals.

**CORN:** Use 20 lbs. per acre to control CORN EARWORM, FALL ARMYWORM, SAP BEETLES and JAPANESE BEETLES. Do not apply within 7 days of harvest for forage. For Corn Earworms apply to silks as soon as they appear. Make 3 to 4 applications of 2-3 day intervals. Reapply immediately after rain.

**COTTON:** Use 20 to 25 lbs. per acre to control THRIPS, FLEA BEETLES, LEAFHOPPERS, LEAFWORMS, BOLLWORMS, FALL ARMYWORMS, LEAF FORATORS, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLARS, STINK BUGS and GRASSHOPPERS. Do not graze treated fields or feed cotton trash to dairy or meat animals.


**EGGPLANTS, PEPPERS and TOMATOES:** Use 20 lbs. per acre to control TOMATO FRUITWORMS, FALL ARMYWORMS, FLEA BEETLES, LEAFHOPPERS, TOMATO HORNWORMS, EUROPEAN CORN BORER, STINK BUGS, COLORADO POTATO BEETLES and LYGUS BUGS. No time limitation.

**FILBERTS (Not for use in California):** Use 40 to 50 pounds per acre to control FILBERT APHID, FILBERT LEAFROLLER, and FILBERT MOTH.

**GRAPES:** Use 20 to 25 lbs. per acre to control GRAPE LEAF FOLDERS, GRAPE LEAFHOPPERS, GRAPE BERRY MOTHS and JAPANESE BEETLES. No time limitation.

**POTATOES:** Use 15 to 20 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO POTATO BEETLES. Apply to foliage only.

**STRAWBERRIES:** Use 20 lbs. per acre to control MEADOW SPITTLEBUGS, and LEAFROLLERS. Do not apply within 1 day of harvest.



F 251

BEST DOCUMENT AVAILABLE