

SEVIN® 10 DUST

N-METHYL CARBAMATE

ACTIVE INGREDIENT:

EPA Reg. No. 2935-320-ZA

ACCEPTEL

APR 2 1998

Under the Federal Insocioide, Fundicide, and Redundicide Act, as amended, for the posticide registered under EPA Reg. To 2935-320

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: May be harmful if swallowed. Ayold breathing dust, Ayold contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: A) Long sleeved shirt; B) Long pants; C) Waterproof gloves; D) Shoes plus socks,

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intentidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BRING TREATED.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

USER SAFETY RECOMMENDATIONS

Users should:

- . Wash hands before eating, drinking, chewing gum, using tobacco or using the tollet
- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Oo not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural posticides. It contains requirements for training, decontamination, rioillication, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated gress that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: A) Coveralls; B) Waterproof gloves; C) Shoes plue socks.

DIRECTIONS FOR USE (cont.)

NOT FOR HOME USE

ALFALFA and CLOVERS: Use 15 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, JAPANESE BEETLES, ALFALFA CATERPILLARS, ARMYWORMS, CORN EARWORMS and STINK BUGS. Do not apply within 3 days of harvest on alfalfa.

-APRICOTO, PEACHES, PLUMS and NECTARINES: Use 40 to 50 lbs. per agre by eir or ground to control TWIG-BORER and ORIENTAL FRUIT MOTH, Do not use — within 3 days of harvest on Apricols and Nectarines or within 1 day on Peaches and Prums.

BEAMS: Use 20 lbs. per acre to control MEXICAN BEAM BEETLES, LEAFHOPPERS, BEAM LEAF BEETLES, CUCUMBER BEETLES, WESTERN BEAM CUT-WORMS, FLEA BEETLES, CORN EARWORMS, ARMYWORMS, VELVET BEAM CATERPILLARS, LYGUS and STINK BUGS, Do not feed treated vines to meat or dairy animals.

CORN: Use 20 lbs. per acre to controt CORN EARWORM, FALL ARMYWORM, SAP BEETLES and JAPANASE BEETLES. Do not apply within 7 days of harvest for forage. For Com Earworms apply to sits as soon as they appear, Make 3 to 4 applications at 2-3 day intervals. Reapply immediately effect rain.

COTTON: Use 25 lbs. per acre to control THRIPS, FLEA BEETLE, FLEAHOPPER, LEAFWORM, BOLLWORM, FALL ARMYWORM, LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLAR, STINK BUGS and GRASSHOPPERS. Do not graze treated fields or feed cotton tresh to delay animals or most animals. Do not use within 7 days of harvest.

KEEP OUT OF REACH OF CHILDREN! CAUTION:

Carbaryl (1-naphthyl N-methylcarbamate)

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately, Give a glass or two of water and induce comiting by touching back of throat with finger. It is preferable to induce comiting under medical supervision or to use gastric lavage with a culled endotracheal tube because of aspiration hazard, Remove victim immediately to emergency treatment leatings.

If on skin: Remove by washing with soap and water. Get medical attention,

If in eyes: Flush the eyes with clean water for 10 minutes. Get medical attention.

If Inhaled: Remove victim to Iresh air, Transport immediately to emergency treatment facility.

ANTIDOTE: Note to Physician, Carbaryl is a moderate, reversible, cholinesterase inhibitor. Alrepine is antidotal.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON LEFT PANEL

STORAGE AND DISPOSAL

- 1. Prohibitions: Do not contaminate water, food or feed by storage or disposal.
- 2. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
- 4. Storage: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

DIRECTIONS FOR USE (cont.)

EGGPLANTS, PEPPERS and TOMATOES: Use 20 lbs. per acre to control TOMATO FRUITWORMS, FALL ARMYWORMS, FLEA BEETLES, LEAFHOPPERS, TOMATO HORNWORMS, EUROPEAN CORN BORER, STINK BUGS, COLORADO POTATO BEETLES and LYGUS BUGS. No lime limitation.

FILBERTS (Not for use in Galifornia): Use 40 to 50 ibs. per acre to control FILBERT APHID; FILBERT LEAFROLLER and FILBERT MOTH;

GRAPES: Use 20 to 25 lbs. per acre to control GRAPE LEAF FOLDERS, GRAPE LEAFHOPPERS, GRAPE BERRY MOTHS and JAPANESE BEETLES. No time limitation.

POTATOES: Use 15 to 20 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO POTATO BEETLES. Apply to foliage only.

STRAWBERRIES: Use 20 lbs. per acre to control MEADOW SPITTLEBUGS and LEAFROLLERS. Do not apply within 1 day of harvest.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the sollor shell be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

Sevin* is a tradomark of Rhone-Poulenc for 1-naphthyl N-methylcarbamate.

WILBUR-ELLIS Loco* and IDEAS TO GROW WITH* are registered trademarks of WILBUR-ELLIS Company.

In Case of Emergency, Call CHEMTREC: (800) 424-9300

NET CONTENTS: 50 POUNDS

WILBUR-ELLIS COMPANY
P.O. BOX 16458 • FRESNO, CA 93755

F-995

100%

EPA Est. No. 70-KY-1

56830

2 2

100