

#### DIRECTIONS FOR USE ON CROPS

Measure the correct amount and add to the sprayer tank while it is running with the agitator running. Stir and mix thoroughly in hand sprayers. Thorough full-coverage of spray application is important and nozzles should be adjusted properly for uniform distribution.

#### RECOMMENDATIONS FOR USE:

**CHERRIES:** For control of cherry fruit fly use 1 qt. in 100 gals. of water. Thorough, full coverage applications should be made beginning when adult flies appear, usually just after the blossom buds open. Repeat application every 10 days until harvest. Do not apply within 2 days of harvest. An additional spray after harvest is suggested, especially if high temperatures have kept flies active. Do not use more than 10 qts. per acre.

**COLE CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower and Kohlrabi):** For control of cabbage worm and cabbage looper use 1 qt. per acre in 25 to 100 gals. of water in ground sprayers or in 5 gals. of water in aerial sprayers. Apply when insects are first noted, making thorough applications at 7 to 10 day intervals as required. Do not apply within 3 days of harvest.

**LETTUCE AND SPINACH:** For control of leafhoppers, cabbage looper, (imported Southern) cabbage worm and alfalfa looper (California) use 1 qt. in 25 to 100 gals. of water in ground sprayers or in 5 gals. of water in aerial sprayers. Apply when insects are present making thorough application. Repeat at 7 to 10 day intervals as required. Do not apply to lettuce within 4 days of harvest or to Spinach within 7 days of harvest.

Do not use more than 1 qt. per acre on Spinach and no more than 2 qts. per acre on lettuce.

**CONTAINER DISPOSAL:** Do not reuse empty container or drums. Return container or drum to reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

# red-top

## PERTHANE 4 E. C.

(contains 4 lbs. technical Perthane per gallon)

#### COMPOSITION:

##### ACTIVE INGREDIENTS:

Diethyl diphenyl dichloroethane

40.6%

Related reaction products

5.5%

Xylene

34.5%

##### INERT INGREDIENTS

19.4%

TOTAL

100.0%

EPA Reg. No. 2935 324-AA

**CAUTION: KEEP OUT OF REACH OF CHILDREN!**

#### CAUTION

This solution if brought into contact with the skin may cause toxic symptoms.

Avoid contact with skin. In case of skin contact wash with soap and water.

Avoid breathing spray mist. Harmful if swallowed.

Avoid contamination of feed and feedstuffs.

Do not use in dairy barns, or dairy animals, or in buildings used for the storage of feed or forage to be fed these animals.

This material is compatible with other materials used in pest control programs.

Consult your State Agricultural Experiment Station for further information on the frequency of application.

This product is poisonous to fish. Keep out of lake, ponds, streams, etc.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

WE

WILBUR-ELLIS COMPANY

WE

FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

RECI

PEA

WAI

NOT  
which  
prese  
use 1  
seller  
buyer  
mater

# red-top

## PERTHANE 4 E. C.

(contains 4 lbs. technical Perthane per gallon)

### COMPOSITION:

#### ACTIVE INGREDIENTS:

Diethyl diphenyl dichloroethane 40.6%

Related reaction products 5.5%

Xylene 34.5%

INERT INGREDIENTS 19.4%

TOTAL 100.0%

EPA Reg. No. 2935 324-AA

### CAUTION: KEEP OUT OF REACH OF CHILDREN!

#### CAUTION

This solution if brought into contact with the skin may cause toxic symptoms.

Avoid contact with skin. In case of skin contact wash with soap and water.

Avoid breathing spray mist. Harmful if swallowed.

Avoid contamination of feed and foodstuffs.

Do not use in areas where children and animals are playing. In case of a spill, do not allow children or animals to enter the area.

This material is compatible with other commonly used insecticides and fungicides.

Consult your State Agricultural Experiment Station for further details as to the correct frequency of application.

This product is poisonous to fish. Keep out of lake, ponds, and streams.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

WE

# WILBUR-ELLIS COMPANY

WE

FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

### RECOMMENDATIONS FOR USE: (Continued)

**PEARS:** For control of pear psylla - Pacific Northwest. Use 1 to 2 quarts in 100 gallons of water in standard dilution sprays, or the equivalent in high concentrate sprays. Sufficient spray should be applied to provide 8 to 16 lbs. active ingredient per acre. Make applications prebloom or very early postbloom **before fruit formation** to control over-wintering and early Spring populations of pear psylla. If summer sprays are required, other insecticides should be used in these applications to avoid residues of Perthane on the fruit. **DO NOT USE PERTHANE WHILE FRUIT IS PRESENT.** A post harvest spray of Perthane is recommended to reduce tree injury from psylla toxin. Do not use more than 16 qts. per acre. Do not apply within 7 days of harvest. If more than 8 qts. per acre are applied, allow 30 days between applications.

**WARNING — FLAMMABLE!** Keep away from heat and open flame.

**NOTICE:** The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

4-19-74  
2935 324

29 JAN 1974  
Pesticides Regulation