

1/4 2935-329 P. 112

RECOMMENDATIONS FOR USE:	Days between final application & harvest	Other limitations & application schedules
Nectarines & Peaches Plum Curculio Oriental Fruit Moth San Jose Scale Peach Twig Borer California Peach Weevil Orange Tortrix Red-Banded Leafhopper Malloxia Pear Spindle Webworm Red Spider Mite Aphids Pear B. Scale Pear Psylla Pear Thrips	1/2 pt/100 gal water BY AIR: 1 pt/acre	Do not apply within 14 days of harvest
Dormant Spray California	1/4 pt/100 gal water BY AIR: 3/4 pt/acre	Do not apply within 14 days of harvest
Onions	Do not exceed 2.5 lbs actual per acre	Do not apply more than once after bloom. Do not apply more than 5 lbs actual Parathion per acre between Jan 1 and harvest
Onions Thrips Onion Maggot	1/2 pt/acre BY AIR: 1/2 pt/acre	15 Repeat at weekly intervals as necessary. Apply at time of planting onion seeds
Pears Pear Bristle Nite Pear Psylla	1/4 to 1/2 pt/100 gal water BY AIR: 1 to 1-1/2 pt/acre	14 Apply as pre-blossom spray
Pears Pear Psylla	1/4 pt/100 gal water BY AIR: 3/4 to 1 pt/acre. Do not exceed 3 lbs actual per acre	Apply as a pre-blossom or post-blossom application Use as 1.5 lb coverage spray
Pears Aphids, Pear Weevil	1/4 to 1/2 pt/acre (Cal use 1/4 to 1/2 pt/acre)	10
Peppers Aphid and Serpentine Leafminer	1/4 pt/acre BY AIR: 1/2 pt/acre	15
Potatoes Colorado Potato Beetle, Flea Beetles, Leafhoppers, Serpentine Leafminer, Grasshoppers, White Fly, Aphids Potato Psyllid	1/4 pt/acre BY AIR: 1/2 pt/acre	5 1/2 to 1 pt/acre by ground or air
Plums & Prunes Plum Curculio (Plum only) Bud Moth, Milky Plum Louse, Tortrix, Leaf Rollers, Leafhoppers, Pear B. Scale, Pear Spindle Webworm, Pear Thrips	1/4 to 1/2 pt/acre BY AIR: 1 to 1-1/2 pt/acre. Do not exceed 4.0 lbs actual per acre	14
Plums & Prunes Peach Twig Borer Pear Thrips Pear B. Scale Pear Spindle Webworm	1/2 pt/100 gal water as above spray 1 to 1-1/2 pt/acre in 50 to 100 gal water as concentrate BY AIR: 1 to 1-1/2 pt/acre	14
Dormant Spray San Jose Scale, Pear Thrips, Pear B. Scale, Pear Spindle Webworm, Pear Thrips	1/2 pt/100 gal water BY AIR: 1/2 pt/acre	14
Rape Crops For Seed (Not for use in Calif.) Cabbage Seedpod Weevil (Anthrenus) (Larvae)	1/2 to 1 pt/acre. Use higher rate during cool weather or when infestations are heavy	28 Make first application when two or more may be found per plant on young pods. If treatment is necessary during bloom, treat only from evening to morning to protect bees. Do not apply later than 3 hours after sunrise. Do not graze treated field or feed forage or straw to livestock
Sugar Beets Flea Beetles, Leafhoppers, Curculionids, Armyworms	3/4 pt/acre	15
Squash Melon Worm, Pickworm, Serpentine Leafminer, Aphids	1/4 pt/acre BY AIR: 1/2 pt/acre	15
Squash Aphids	1/4 pt/acre BY AIR: 1/2 pt/acre	14
Tomatoes Flea Beetle, Leafhoppers, Serpentine Leafminer, Grasshoppers, White Fly, Aphids	1/4 pt/acre BY AIR: 1/2 pt/acre	10
Strawberries Leafhopper and Red Spider Mite	1/4 to 1/2 pt/acre BY AIR: 1/2 pt/acre	14 Repeat applications at 7 to 10 days intervals until desired control is achieved
Small Grains (Wheat, Oats and Barley) Greenbugs (Aphid), Armyworms, Grasshoppers	1/4 to 3/4 pt/acre by ground or air (Cal use 1/4 to 1/2 pt/acre)	15
Turnips Diamond-back Moth, Imported Cabbageworm, Armyworms, Aphids, Thrips	1/4 to 1/2 pt/acre BY AIR: 1/2 pt/acre	10 Do not apply within 21 days of harvest if eggs are to be used for food or feed
Witch Greenbugs (Aphid), Armyworms, Grasshoppers	1/4 to 1/2 pt/acre by ground or air	15

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, expressed or implied, concerning the material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

RESTRICTED USE PESTICIDE
 Due to very high acute toxicity to humans and birds

For retail sale to and conformity by certified applicator or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Commercial certified applicators must also insure that all persons involved in these activities are informed of the precautionary statements.

Wilbur-Ellis
PARATHION 8 FLOWABLE

Directions For Use
 Aqueous Emulsion Concentrate Sold for Agricultural Insecticide Purposes

Parathion® 80%
 INERT INGREDIENTS 20%
 *O, O-dimethyl O-p-nitrophenyl phosphorothioate.
 Contains 8 lbs. Parathion per gallon. 100%

EPA Reg. No. 2935-329-AA EPA Est: 2935-CA-1

KEEP OUT OF REACH OF CHILDREN

DANGER  **POISON**
PELIGRO

ACCEPTED
 DEC 1 0 1987
 2935-329

OBSERVE PRECAUTIONS ON SAFETY LABEL
 NOT FOR USE OR STORAGE IN OR AROUND THE HOME
 Do not apply this product through any type of irrigation system.

ENDANGERED SPECIES RESTRICTIONS

The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of federal laws. The use of this product is prohibited to prevent death or harm to endangered or threatened species that occur in the following counties or elsewhere in their range.

Before using this pesticide in the following counties you must obtain the EPA Cropland Endangered Species Bulletin. The use of this pesticide is prohibited in these counties unless specified otherwise in the Bulletin. The EPA Bulletin is available from either your County Agricultural Extension Agent, the Endangered Species Specialist in your State Wildlife Agency Headquarters or the appropriate Regional Office of either the U.S. Fish and Wildlife Service (FWS) or the U.S. Environmental Protection Agency. THIS BULLETIN MUST BE REVIEWED PRIOR TO PESTICIDE USE.

IF USED ON CROPS:	STATE (Regional Office FWS) Species	COUNTY
ALABAMA (Atlanta, GA)	Stackwater darter Alabama cavefish Freshwater mussels	Lauderdale, Limestone, Madison Lauderdale Colbert, Jackson, Marshall, Morgan
ARIZONA (Albuquerque, NIA)	Woundfin Bonytail chub Gila topminnow	Mohave Mohave Graham, Santa Cruz
ARKANSAS (Atlanta, GA)	Freshwater mussels	Clay, Clark, Cross, Lawrence, Randolph, Sharp, St. Francis Benton Sevier
ARIZONA (Portland, OR)	Delta green ground beetle Valley elderberry longhorn beetle Austrian Canada goose Kern pinnose sphinx moth Blunt-nosed leopard lizard Santa Cruz long-toed salamander Unarmored three-spine stickleback	Solano Merced, Sacramento Colusa, Merced, Sutter, Stanislaus Kern Fresno, Kern, Kings, Madera, Merced, Monterey, San Luis Obispo, Santa Barbara, Stanislaus, Tuolumne Monterey, Santa Cruz Los Angeles, Santa Barbara
COLORADO (Denver, CO)	Colorado aqueduct Humpback chub	Blanco, Delta, Garfield, Mesa, Moffat, Rio, Rout Mesa
FLORIDA (Atlanta, GA)	Everglade kite	Broward, Dade, Glades, Palm Beach
KENTUCKY (Atlanta, GA)	Freshwater mussels	Ballard, Edmundson, Jackson, Laurel, Marshall, McCracken, Pulaski, Rockcastle, Warren, Wayne

Wilbur-Ellis

P.O. BOX 16458 • FRESNO, CA 93755

PLPUS

ENDANGERED SPECIES (cont.)

STATE (Regional Office FWS) Species	COUNTY
MARYLAND (Newton Corners, MA) Maryland darter	Harford
MISSISSIPPI (Atlanta, GA) Bayou darter	Copiah
MISSOURI (Twin Ck's, MN) Ozark cavefish	Christian, Greene, Jasper, Lawrence, Newton, Barry, Stone
NEVADA (Portland, OR) Woundfin	Clark
Pahrangat bonytail	Lincoln
Cut-throat	Washoe
Fahumpki fish	Clark, White Pine
NEW MEXICO (Albuquerque, NM) Pecos gambusia	Chaves, Eddy
NORTH CAROLINA (Atlanta, GA) Spotfin chub	Macon, Swain
OHIO (Twin Ck's, MN) Scioto madtom	Champagne, Franklin, Logan, Madison, Pickaway, Union
OKLAHOMA (Albuquerque, NM) Leopard darter Ozark cavefish	Leflore, McCurtain Delaware
TENNESSEE (Atlanta, GA) Stackwater darter Slender chub Spotfin chub Freshwater mussels	Lawrence, Wayne Claborn, Hancock Cumberland, Ferreston, Morgan Blount, Claiborne, Decatur, Franklin, Hancock, Hardin, Lincoln, Loudon, Marshall, Maury, Rhea, Roane, Scott, Sequatchie, Smith, Sullivan
TEXAS (Albuquerque, NM) Arwater's greater prairie chicken Comanche Springs pupfish Pecos gambusia Texas blind salamander San Marcos salamander San Marcos gambusia Houston loach Fountain darter	Arkansas, Austin, Colorado, Fort Bend, Jeff Davis, Goliad, Refugio, Victoria Arkansas, Austin, Colorado, Fort Bend, Jeff Davis, Goliad, Refugio, Victoria Jeff Davis, Pecos, Reeves Hays Hays Hays Bastrop, Burleson Hays
UTAH (Denver, CO) Woundfin Humpback chub Bonytail chub Colorado squawfish	Washington Grand Grand, Uintah Carbon, Duchesne, Emery, Garfield, Grand, Kane, San Juan, Uintah, Wayne
VIRGINIA (Newton Corners, MA) Spotfin chub Freshwater mussels	Scott, Washington Lee, Russell, Scott, Smyth, Tazewell, Washington, Wise

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Use designated amount of product in the following gallons of water per acre:
GROUND APPLICATION: Vegetable and Field Crops — 20 to 250 gal's Cotton — 3 to 20 gal's
 Deciduous Fruit Crops — 40 to 800 gal's Citrus — 100 to 3,000 gal's (On Young Fruit Plantings — Use a minimum of 15 gal's water per acre.)
AIR APPLICATION: Field Crops — 3 to 10 gal's Orchard Crops — 10 to 20 gal's Use fine mist spray; coarse droplet spray w/3 cause spot injury to plants and fruit.

Crop	Pest controlled	Rates	Days between final application & harvest	Other limitations & application schedules
Almonds	Oriental Fruit Moth, Twig Borer, Tent Caterpillar, Leaf Roller	1/3 pt/100 gal's water BY AIR 1-2 pts per acre in 10 to 20 gal's of water. Do not apply more than 2.0 lbs actual per acre.	Do not apply after buds open.	Apply as a full coverage spray.
Doanant Spray		1/2 pt to 1 pt per 100 gal's of water. Do not exceed 8.0 lbs actual per acre.		
Apples	Codling Moth, Plum Curculio, Orange Tortrix, San Jose, Foxes and Spurs, Scab, Red Blended Leaf Roller (Second & Third Brood), Woolly Acale, Aphid and Mite/bug	1/4 to 1/3 pt/100 gal's water BY AIR 1-1/2 pts per acre. Do not exceed 8.0 lb actual per acre.	14	Certain mites such as Two Spined and Western, repeat application at 14 to 30 day intervals during summer months. This product may cause injury to fruit and to type of leafhopper, aphid and related mites. Do not apply to susceptible varieties. Use as a full coverage spray.

Crop	Pest controlled	Rates	Days between final application & harvest	Other limitations & application schedules
Apples (cont.)	Bud Moth, Red Bud, Fruit Tree Leaf Roller, Rosy Apple Aphid, Green Apple Aphid, Red Blended Leaf Roller (First Brood), Grasshoppers, European Red Saw, Two Spined	2.5 pt/100 gal's water BY AIR 1 pt/acre. Do not exceed 8.0 lbs actual per acre.		
Apricots	Codling Moth, Oriental Fruit Moth and Tortrix, Bud Moth, Fruit Tree Leaf Roller, Sucker Mites and Aphids	1/3 pt/100 gal's water. Do not exceed 3.6 lbs actual per acre. 1/4 pt/100 gal's water	14	Do not apply until danger of bee poisoning has passed.
Alfalfa	Greenbug (Aphid), Armyworms, Grasshoppers	1/4 to 1/3 pt per acre	15	
Artichoke	Plum Moth and Aphid	1/4 to 1/3 pt/100 gal's water BY AIR 1/2 pt/acre	7	
Beans	Mexican Bean Beetle, Mexican Aphids, Red Spider Mites, Two Spined Spider Mite, Armyworms, Leaf Miner and Leaf Miner	1/3 pt/acre BY AIR 1/2 pt/acre	15	Repeat applications at 7 to 10 day intervals until control is achieved.
Beets	Webworm	1/3 pt/acre BY AIR 1/2 pt/acre	15	21 days before use for food or feed.
	Aphids, Leafhoppers	1/4 pt/acre BY AIR 1/2 pt/acre		
Broccoli	Diamond Back Moth, Imported Cabbage Worm, Armyworm, Aphid, Thrips	1/4 to 1/2 pt/acre BY AIR 1/2 pt/acre	7	
Brussels Sprouts				
Cauliflower (Not for use in Calif.)	Diamond Back Moth, Imported Cabbage Worm, Armyworm, Aphids, Thrips	1/4 to 1/2 pt/acre BY AIR 1/2 pt/acre	10	
Cabbage	Diamond Back Moth, Imported Cabbage Worm, Armyworm, Aphids, Thrips	1/2 to 1/4 pt/acre BY AIR 1/2 pt/acre	10	
Carrots	Aphids	1/2 pt/acre BY AIR 1/2 pt/acre	15	Do not use treated carrot tops for food or feed.
Celery	Celery Worms, Aphid	1/4 pt/acre BY AIR 1/3 pt/acre	21	
Cherries	Plum Curculio, Cherry Fruitworm, Oriental Fruit Moth, Tortrix and Cankerworm	1/4 pt/100 gal's water BY AIR 1 pt/acre	14	Apply as a full coverage spray.
	Aphids and Red Spider Mites	1/4 pt/100 gal's water BY AIR 1/2 pt to 3/4 pt per acre. Do not exceed 2.1 lbs actual per acre.		
Citrus (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines)	Aphids, Orange Tortrix, Thrips, Citricola, Citrus Canker, Black and Citrus Red Scales	1/4 pt/100 gal's water. Do not apply more than 4 lbs actual per acre.	14	
Clover	Grasshoppers, Greenbugs (Aphids), Armyworms	1/4 to 3/4 pt/acre (Cal use 1/4 to 1/2 pt/acre)	15	
Cotton (Not for use in Calif.)	Aphids and Spider Mites	1/4 to 1/2 pt/acre	7	Apply when aphids appear or for spider mites. Do not apply when mites appear or when leaves begin turning yellow to reddish brown. Repeat once or twice at 7 to 10 day intervals.
Cranberries (Not for use in Calif.)	Cranberry Tipworm, Fireworms, Blackbeetles, Cranberry & Spinyhorn, Leafhoppers	3/4 pt/acre	15	
Cucumbers	Cucumber Beetle, Melon Worm, Pickworm, Spinyhorn Leaf Miner and Aphids	1/4 pt/acre BY AIR 1/3 pt/acre	15	Do not apply before evening.
Grapes	Mite/bug and Leaf Folder	1/4 pt/100 gal's water BY AIR 1/2 to 3/4 pt/acre. Do not exceed 2.5 lbs actual per acre.	14	On Red Spider Mites make applications at 5 to 7 day intervals until desired control is reached. Apply only before bud set or bud set or after harvest.
Grapes (Domant)	Mite/bugs	2-1/2 pts of Parathion 85% plus 2 to 8 pts of Superior Oil in 200 to 300 gal's of water per acre.		Apply before bud swell. Spray should be directed toward trunks and limbs.
Hops (Not for use in Calif.)	Aphids	1/2 to 3/4 pt/acre	10	
Kale	Diamond Back Moth, Imported Cabbage Worm and Armyworm, Thrip	1/4 - 1/3 pt/acre BY AIR 1/2 pt/acre	10	
Leafy	Aphid, Armyworms, Imported Cabbage Worm, Leafhoppers, Leafminers, Looper, Lygids, Thrips, San March Caterpillar	1/2 pt/acre		7 days on field lettuce 21 days on Bibb and Leaf lettuce
Melons	Melon Worm, Pickworm, Spinyhorn Leaf Miner, Aphids	1/4 pt/acre BY AIR 1/3 pt/acre	7	
Mustard Greens	Diamond Back Moth, Imported Cabbage Worm, Armyworm	1/4 to 1/3 pt/acre BY AIR 1/2 pt/acre	10	
	Aphids & Thrips	1/4 pt/acre BY AIR 1/2 to 1/2 pt		

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Poisonous if Swallowed, Inhaled or Absorbed Through Skin. Rapidly absorbed through skin. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to Parathion poisoning.

This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed. Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (and goes through clothes). Breathing vapors, spray mist or dust may be fatal. Workers entering treated fields within 48 hours must wear protective clothing.

STATEMENT OF PRACTICAL TREATMENT

Call a doctor (physician), clinic or hospital immediately. Explain that victim has been exposed to parathion and describe his condition.

If inhaled and breathing has stopped, start artificial respiration immediately and maintain until doctor sees victim.

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF IN EYES: Flush with plenty of water and get medical attention immediately.

IF ON SKIN: Immediately wash with soap and water and get medical attention.

POISON SIGNS (Symptoms)

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die. Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

NOTE TO PHYSICIAN

Antidote — administer atropine sulfate in large doses; TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement; **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

P&FS-9

RESTRICTED USE PESTICIDE

Due to very high acute toxicity to Humans and Birds

For retail sale to and use only by certified applicator or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

Wilbur-Ellis

PARATHION 8 FLOWABLE

ACTIVE INGREDIENTS:	By Weight
Parathion: O, O-diethyl O-p-nitrophenyl Phosphorothioate	80.0%
INERT INGREDIENTS:	20.0%
This product contains 8 lbs. of Parathion per gallon. 100.0%	
EPA Reg. No. 2935-329-AA	Est. No. 2935-CA-1

STOP — READ LABEL BEFORE USING!

**PELIGRO
POISON**



DANGER

PRECAUTION ALUSUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

See Side Panels for Antidote and Precautions

KEEP OUT OF REACH OF CHILDREN

Wilbur-Ellis

P.O. Box 16458, Fresno, CA 93755

ENVIRONMENTAL

This pesticide is highly toxic to fish and will not apply directly to water or wetlands (swamps and drift) from target areas may be hazardous. Do not contaminate water by cleaning.

This product is extremely toxic to bees blooming crops or weeds. Do not apply the weeds if bees are visiting the treatment area.

CHEMICAL AND P

KEEP AWAY FROM HEAT OR OPEN FLAME

ENDANGERED SPECIES

The Agency informs the pesticide applicator of the presence of endangered species. The pesticide and endangered species labeling that follows identifies endangered species. See attached label.

WORK SAFELY

Keep all unprotected persons and children away from drift.

Do not rub eyes or mouth with hands. If you get it on your face, wash it off immediately. Call a doctor (physician), clinic or hospital if you have been exposed to parathion and if a hospital is recommended.

THIS PRODUCT MAY BE FATAL IF SWALLOWED. CONTACT SKIN FAILURE TO PROPERLY WEAR PROTECTIVE CLOTHING AND EQUIPMENT.

USE ONLY WHEN WEARING THE FOLLOWING EQUIPMENT DURING MIXING/LOADING OF APPLICATION EQUIPMENT, DISPOSING OF TREATED AREAS: Waterproof pants, gloves; rubber boots or rubber overshoes; face shield; NIOSH approved respirator; chemical resistant apron when using the concentrate; nonenclosed cockpit, a helmet with a visor; brimmed hat and safety goggles or face shield.

IF MIXING/LOADING IS PERFORMED OUTSIDE OF PROTECTIVE CLOTHING AND EQUIPMENT: Heavy-duty chemical resistant gloves; shirt (or gauntlets and short sleeve shirt); Safety goggles or a face shield must be worn.

All other protective clothing and equipment required for use with open systems must be available nearby.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE: Clean long-sleeved shirt and long-legged pants. All other protective clothing and equipment required for use during application must be available in the cab and must be worn when exiting the cab into treated areas. If used for this purpose, contaminated clothing may not be brought back into the cab unless in an enclosure such as a plastic bag. **REMEMBER — THIS CLOTHING IS NOT INTENDED TO PROTECT YOU DURING REPAIR AND CLEANING OF APPLICATION EQUIPMENT AND EARLY REENTRY! REFER TO THE INSTRUCTIONS ABOVE.**

HUMAN FLAGGERS ARE STRICTLY PROHIBITED DURING AERIAL APPLICATION.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water and consult a physician immediately. Always wash hands, face and arms with soap and water before smoking, eating, drinking or toileting.

AFTER WORK: Wash gloves with soap and water before removing. Take off all work clothes and shoes. Store protective clothing separately from personal clothing. Launder protective clothing after each use. Shower using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Personal clothing worn during mixing/loading, application, repair and cleaning of application equipment, disposal of pesticide, and early reentry into treated fields must be stored and laundered separately from household articles. Clothing and equipment heavily contaminated or drenched with parathion must be destroyed according to state and local regulations.

HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

Respirators should be cleaned and cartridges replaced according to instructor's included with respirators. Replace gloves frequently.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas for 48 hours unless appropriate protective clothing is worn. Because certain states may require more restrict re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Reentry into apple, citrus, peach, nectarine or grape fields in the states of Arizona, California, Nevada, New Mexico, Oklahoma, Texas, and Utah is prohibited for seven days after the end of application, unless appropriate protective clothing and equipment required

or early reentry as described elsewhere on the label is worn. Reentry into all other treated areas is prohibited for 48 hours after end of application, unless all required protective clothing and equipment is worn.

Written or oral warnings must be given to workers who are expected to be in a treated area if in an area shown to be treated with this product. Oral warnings should specify protective clothing (see left), areas or fields that cannot be entered without this clothing, period of time area is to be treated, and statement of practical treatment (see left) in case of exposure. Written warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Danger: Area treated with Parathion on (date of application). Do not enter without appropriate protective clothing for 48 hours. (Insert here actions to take in case of accidental exposure — see left)"

POST TREATED AREA: Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

STORAGE AND DISPOSAL

1. Prohibitions: Do not contaminate water, food or feed by storage or disposal.
2. Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
3. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. Storage: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing, or pouring.

CROP USE DIRECTIONS

Use only according to attached labeling.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other, from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

F587

LARGER & BOLDER

IN CASE OF EMERGENCY,
CALL CHEMTREC: (800) 424-9300



POISON

DEC 1 0 1987

U.S. DEPARTMENT OF AGRICULTURE
FEDERAL AGENCY FOR PROTECTION OF THE ENVIRONMENT
EPA Reg. No. 2935-329-1

BOLDER
LINES

RQ, PARATHION MIXTURE, LIQUID
NA 2783

LARGER & BOLDER

BEST AVAILABLE COPY