MANEB 6 DUST

(Right Panel)

COMPOSITION:

ACTIVE INGREDIENT:

Maneb (Manganese ethylene bisdithiocarbamate. 6%

INERT INGREDIENTS. 94%

100%

USDA Reg. No 2935 -

CAUTION: KEEP OUT OF REACH OF CHILDREN!
(18 point) (12 point)

May cause irritation of nose and throat! Do not breathe dust. Store in cool, dry place. Keep bag closed when not in use.

This product is toxic to fish. Keep out of lakes, streams, and ponds.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM AREA BEING TREATED.

RECOMMENDATIONS FOR USE: (Continued)

ONIONS: Use 30-40 lbs. per acre to control BLAST (Botrytis leaf blight), DOWNY MILDEW and PURPLE BLOTCH. Apply when first spots appear; repeat weekly as needed.

POTATOES: Use 40-50 lbs. per acre to control EARLY and LATE BLIGHT. Apply as soon as plants are 2 to 6 inches high; repeat at 5 to 10 day intervals as long as blights threaten.

TOMATOES: Use 40-50 lbs. per acre to control ANTHRACNOSE, EARLY and LATE BLIGHT, GREY LEAF SPOT (Stemphylium), and SEPTORIA LEAF SPOT. Apply when fruit of first clusters is well formed; repeat at 7 to 10 day intervals. Do not use on young seedlings in green houses as injury may occur.



MANEB 6 DUST

(Left Panel)

RECOMMENDATIONS FOR USE:

BEANS: Use 30-40 .bs. per acre to control ANTHRACNOSE and DOWNY MILDEW. Do not apply within 4 days of harvest. Begin at first sign of disease and repeat at 4 to 6 day intervals as needed.

CABBAGE: Use 30-40 lbs. per acre to control ALTERNARIA LEAF SPOT and DOWNY MILDEW. Apply 7 to 10 days after planting. Apply when disease threatens; repeat weekly as needed. Do not apply within 7 days of harvest.

CARROTS: Use 30-40 lbs. per acre to control LEAF SPOT disease. Apply when disease threatens; repeat every 7 to 10 days.

CELERY: Use 35-40 lbs. per acre to control EARLY and LATE BLIGHT. Remove residues by washing, stripping and trimming. Application should begin in the plant beds as soon as plants emerge. Apply at 3 to 5 day intervals. Continue at 7 to 10 day intervals when plants are established in the field.

CUCUMBERS, MELONS and SQUASH: Use 30-40 lbs. per acre to control ANGULAR LEAF SPOT, ANTHRACNOSE & DOWNY MILDEW. Apply as soon as plants begin to run or when disease threatens; repeat at 7 to 10 day intervals as needed.

LETTUCE and ENDIVE: Use 30 lbs. per acre to control DOWNY MILDEW. Apply when disease appears; repeat at 7 to 10 day intervals under average conditions, 3 to 5 days when disease conditions are severe. Do not apply within 7 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce and endive by washing or other effective means.

Cameraman's Note

Poor Copy