

MANEB 6 DUST

ACCEPTED
 2935-336
 FEB 14 1984
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2935-336

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May cause irritation of eyes, nose and throat. Do not breathe dust. Do not get in eyes, on skin, or on clothing. Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Avoid drift. Do not contaminate feed or food products.

APPLICATOR PRECAUTION

Protective clothing (long pants, long sleeve shirt, impermeable gloves, hat and boots) must be worn during mixing and loading.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where run-off is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- 1. Prohibitions:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2. Pesticide Disposal** — Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.
- 3. Container Disposal**—Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.
- 4. General** — Consult Federal, State or Local Disposal Authorities for additional restrictions and approved alternatives.

Note: Use higher rates under conditions of high pest pressure when a dosage range is indicated.

BEANS: Use 30-40 lbs. per acre to control ANTHRACNOSE and DOWNY MILDEW. Do not apply within 4 days of harvest. Apply when disease threatens; repeat at 4 to 6 day intervals as needed.

CABBAGE: Use 30-40 lbs. per acre to control ALTERNARIA LEAF SPOT and DOWNY MILDEW. Apply 7 to 10 days after planting. Apply when disease threatens; repeat weekly as needed. Do not apply within 7 days of harvest.

CARROTS: Use 30-40 lbs. per acre to control ALTERNARIA and CERCOSPORA LEAF SPOT. Apply when disease threatens; repeat every 7 to 10 days.

CELERY: Use 35-40 lbs. per acre to control EARLY and LATE BLIGHT. Apply when plants are established in the field; repeat as frequently as required (weekly under average conditions; every 3 to 5 days under severe conditions). Remove residues by washing, stripping and trimming. Do not apply within 14 days of harvest.

MB6-3

NOTICE: The statements made on this label, as believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to issue, any warranty, guarantee or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

ACTIVE INGREDIENT:

Maneb (Manganese ethylene bisdithiocarbamate) 6%

INERT INGREDIENTS 94%

100%

State Reg. No. 2935-336-AA

EPA Est. No. 33691-AZ-1

KEEP OUT OF REACH OF CHILDREN!

CAUTION:

ONLY FOR SALE TO, USE, AND STORAGE BY
 AGRICULTURAL AND SERVICE PERSONS.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

— SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS —

DIRECTIONS (Continued)

CUCUMBERS, MELONS and SQUASH: Use 30-40 lbs. per acre to control ANGULAR LEAF SPOT, ANTHRACNOSE and DOWNY MILDEW.

Apply as soon as plants begin to run or when disease threatens; repeat at 7 to 10 day intervals as needed. Do not apply within 5 days of harvest.

GRAPES: Use 25 lbs. per acre to control BUNCH ROT. Make first application when berries are starting to break color and repeat every 14-21 days for three applications. Do not apply within 7 days of harvest.

LETTUCE and ENDIVE: Use 30 lbs. per acre to control DOWNY MILDEW. Apply when disease appears; repeat at 7 to 10 day intervals under average conditions, 3 to 5 days when disease conditions are severe. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce and endive by washing or other effective means.

ONIONS: Use 30-40 lbs. per acre to control BLAST, DOWNY MILDEW and PURPLE BLOTCH. Apply when first spots appear; repeat weekly as needed.

POTATOES: Use 40-50 lbs. per acre to control EARLY and LATE BLIGHT. Apply as soon as plants are 2 to 6 inches high; repeat at 5 to 10 day intervals as long as blights threaten.

TOMATOES: Use 40-50 lbs. per acre to control ANTHRACNOSE, EARLY and LATE BLIGHT, GREY LEAF SPOT (Stemphyllium), and SEPTORIA LEAF SPOT. Apply when fruit of first clusters is well formed; repeat at 7 to 10 day intervals. Do not use on young seedlings in green houses as injury may occur. Do not apply within 5 days of harvest

F-184

[Wilbur-Ellis]

FRESNO • LOS ANGELES • GLENDALE, ARIZ. • PORTLAND • SEATTLE