

RED-TOP BRAND 2, 4, 5-TP 400

A plant hormone-type product effective in controlling the harvest drop of apples of all varieties tested including McIntosh, Delicious, Golden Delicious, Starking, Stayman, Turley, Winesap & Rome Beauty.

RECOMMENDATIONS

FOR DROP CONTROL OF APPLES - Conventional full-volume ground sprayers: Use 2.5 fl. oz. of 2, 4, 5-TP 400 Fruit Spray per 100 gallons of water. This dilution will provide a spray solution of 20 ppm of silvex. Do not exceed this concentration.

Apply 4 days before harvest drop normally begins (7 to 20 days before harvest). 2, 4, 5-TP 400 becomes effective within 7 to 12 days after application and maintains a high degree of drop control for 5 to 7 weeks.

TO IMPROVE SET & YIELD OF ANJOU PEARS - Use 1 fl. oz. of 2, 4, 5-TP 400 per 110 gallons of water to make a 7.5 ppm spray solution. Do not spray a stronger spray. Apply a full coverage spray during the week after harvest to increase fruit set the following year.

RED-TOP

2, 4, 5-TP 400 A HORMONE TYPE SPRAY

For control of preharvest drop of apples.

To Improve Set & Yield of Anjou Pears, for control of preharvest drop of Italian type prunes.

ACTIVE INGREDIENT:

Silvex (2-(2, 4, 5-Trichlorophenoxy) propionic acid) as the triethanolamine salt*. 17.3%

INERT INGREDIENTS

*Equivalent to 11.1% of 2-(2, 4, 5-Trichlorophenoxy) propionic acid or 400 grams per gallon

PROTECT FROM FREEZING

USDA Reg. No. 2935


CAUTION: KEEP OUT OF REACH OF CHILDREN
(18 point) (12 point)

CAUTION: Avoid contact with skin, eyes, and clothing. Wash hands after use.

Harmful if swallowed. Avoid spray drift to nearby susceptible plants as this product may injure cotton, beans, peas, grapes, and ornamentals. Coarse sprays are less likely to drift. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Avoid contamination of irrigation ditches or water used for domestic purposes.

FOR CONTROL OF PREHARVEST DROP OF ITALIAN TYPE PRUNES
Use 2½ fl. oz. of 2, 4, 5-TP 400 per 100 gallons of water (a 20 ppm solution) 20.6 and spray trees until dripping wet. Applications should be made not less than 60 days before harvest and not earlier than 800 to 1100 degree days after full bloom when above 43°F. For correct time of application consult local agricultural authorities since these directions are based on use in

DIRECTIONS FOR USE - Keep agitator running during filling, mixing and spraying operations. Apply as a spray, wetting foliage uniformly and completely. Do not delay harvest as fruit condition and keeping qualities will be impaired if left on trees beyond normal maturity.



in controlling the harvest drop of apples of all varieties tested including McIntosh, Delicious, Golden Delicious, Starking, Stayman, Turley, Winesap & Rome Beauty.

RECOMMENDATIONS

FOR DROP CONTROL OF APPLES - Conventional full-volume ground sprayers: Use 2.5 fl. oz. of 2, 4, 5-TP 400 Fruit Spray per 100 gallons of water. This dilution will provide a spray solution of 20 ppm of silvex. Do not exceed this concentration.

TIME OF APPLICATIONS - Fruit drop control: Apply 4 days before harvest drop normally begins (7 to 20 days before harvest). 2, 4, 5-TP 400 becomes effective within 7 to 12 days after application and maintains a high degree of drop control for 5 to 7 weeks.

TO IMPROVE SET & YIELD OF ANJOU PEARS - Use 1 fl. oz. of 2, 4, 5-TP 400 per 110 gallons of water to make a 7.5 ppm spray solution. Do not spray a stronger spray. Apply a full coverage spray during the week after harvest to increase fruit set the following year.

2, 4, 5-TP 400
A HORMONE TYPE SPRAY

For control of preharvest drop of apples.

To Improve Set & Yield of Anjou Pears, for control of preharvest drop of Italian type prunes.

ACTIVE INGREDIENT:

Silvex (2-(2, 4, 5-Trichlorophenoxy) propionic acid) as the triethanolamine salt*. 17.3%

INERT INGREDIENTS 82.7%

*Equivalent to 11.1% of 2-(2, 4, 5-Trichlorophenoxy) propionic acid or 400 grams per gallon

PROTECT FROM FREEZING

USDA Reg. No. 2935

CAUTION: KEEP OUT OF REACH OF CHILDREN (18 point) (12 point)

CAUTION: Avoid contact with skin, eyes, and clothing. Wash hands after use. Harmful if swallowed. Avoid spray drift to early susceptible plants as this product may injure cotton, beans, peas, grapes, and ornamentals. Coarse sprays are less likely to drift. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Avoid contamination of irrigation ditches or water used for domestic purposes.

Use 2 1/2 fl. oz. of 2, 4, 5-TP 400 per 100 gallons of water (a 20 ppm solution) and spray trees until dripping wet. Applications should be made not less than 60 days before harvest and not earlier than 800 to 1100 degree days after full bloom when above 43°F. For correct time of application consult local agricultural authorities since these directions are based on use in the Pacific Northwest.

DIRECTIONS FOR USE - Keep agitator running during filling, mixing and spraying operations. Apply as a spray, wetting foliage uniformly and completely. Do not delay harvest as fruit condition and keeping qualities will be impaired if left on trees beyond normal maturity.

