


DIAZINON 2 DUST



RECOMMENDATIONS FOR USE:

BEANS (Snap): Use 30 to 50 lbs. per acre to control BLACK BEAN APHIDS, MEXICAN BEAN BEETLES, LEAFHOPPERS, DIPTEROUS LEAF MINERS, and SPIDER MITES. Do not apply within 7 days of harvest.

BEETS, CELERY, LETTUCE and SPINACH: Use 25 lbs. per acre to control APHIDS, FLEA BEETLES and DIPTEROUS LEAF MINER. Do not apply within 10 days of harvest on Lettuce or Spinach; within 14 days on Beets and within 10 days on Celery.

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE, PARSLEY and TURNIP TOPS: Use 30 to 50 lbs. per acre to control APHIDS, IMPORTED CABBAGE WORMS, DIAMOND BACK MOTH, HARLEQUIN CABBAGE BUGS and CORN EARWORMS. Do not apply within 5 days of harvest on Broccoli or Cauliflower; within 7 days on Cabbage, within 10 days on Collards, Kale or Turnip Tops, and within 12 days on Parsley.

FIGS: Use 40 lbs. per acre to control VINEGAR FLIES (*Drosophila* Sp.) and DRIED-FRUIT BEETLES. Do not apply within 5 days of harvest.

GRAPES: Use 50 lbs. per acre to control LEAFHOPPERS, GRAPE BERRY MOTH, GRAPE LEAF FOLDER, and PACIFIC MITES. Do not apply within 10 days of harvest.

ONIONS (including Green Onions): Use 40 to 60 lbs. per acre to control ONION THRIPS and ONION MAGGOTS. Do not apply within 10 days of harvest. For ONION MAGGOT control, direct the dust to the furrow at time of planting.

PEPPERS: Use 30 to 50 lbs. per acre to control APHIDS. Do not apply within 5 days of harvest.

STRAWBERRIES: Use 40 to 60 lbs. per acre to control APHIDS, CYCLAMEN and TWO-SPOTTED MITES. Do not apply within 5 days of harvest.

TOMATOES: Use 30 to 35 lbs. per acre to control APHIDS, VINEGAR FLIES (*Drosophila* Sp.) and DIPTEROUS LEAF MINERS. Do not apply within 1 day of harvest.

F-374

EPA Est. 2935-CA-1

ACTIVE INGREDIENT:

O,O-diethyl o-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate

2%

INERT INGREDIENTS:

98%

100%

EPA Reg. No. 2935-343-AA

KEEP OUT OF REACH OF CHILDREN!

CAUTION: Harmful if Swallowed! May be absorbed through the skin! Do not breathe dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of food or foodstuffs. Keep out of reach of children.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed.

Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

SUGGESTIONS FOR USE: Apply when insects are present on plants and when air is quiet, preferably in early morning. Excessive dust drift is wasteful. Do not mix or dilute this dust with other materials. Even coverage of infested parts of plants is essential for control. Dosage per acre will depend on size of plants, type of duster, degree or severity of infestation. Use only sufficient material to effect desirable control. This product is formulated to be applied by ground or air application.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM AREA BEING TREATED.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by burying with wastes or burning (keep out of smoke).