

MD-102

MIDOLIN MOSQUITO EXCISOL  
(Contains 1% P.M.P. 10%)

For Use Only by Mosquito Abatement Districts, Public Health Officials, and  
Other Trained Persons in Mosquito Control Programs.

ACTIVE INGREDIENT:

0,0-Diethyl (2-(4-methylthio)-p-tolyl) phosphorothioate - 1%

INERT INGREDIENTS - -----

1%

\* U.S. Pat. Off. of Farbwerke Bayer A.G.,  
Chemagro Corporation Licensee.  
U.S. Patent No. 3,042,713

CAUTION (14 point type)

KEEP OUT OF REACH OF CHILDREN (10 point type)

Harmful if swallowed, inhaled, or absorbed through the skin.

Do not get in eyes, on skin, or on clothing.

Do not breathe dust.

Wash thoroughly with soap and warm water after handling.

Wash clothes with soap and hot water after use.

NOTE: This inhibitor of trophic sulfate in large quantities does,  
however, possess a low point of toxicity. Liberal doses of 1 pound  
per acre will give the best results in the control of mosquitoes.

HOW TO USE THIS INSECTICIDE

Apply liberally to all areas of the body.  
It is highly toxic to fish and wildlife. Do not use where fish, snails or  
other animals are abundant. This product is also toxic to bees.  
Do not use in areas where bees are abundant. Do not use in areas where  
wild birds are abundant. Do not use in areas where wild birds are abundant.

ACCEPTED  
MAR 1 1969  
FEDERAL BUREAU OF INVESTIGATION  
U.S. DEPARTMENT OF JUSTICE  
WASHINGTON, D.C. 20535  
435  
COMMENTS

DESCRIPTION

RED-TOP RESIDUAL is recommended for use to control mosquito breeding in  
standing water, residual control and it is effective against DD1-resistant  
and non-resistant strains.

For mosquito larvae control, applications should be made to moist areas,  
standing water, shallow ponds, and around the edges of lawns or ponds, ditches,  
marshes, swamps, and tidal areas. Also apply around garbage dumps,  
manure piles, catch basins, and similar areas where mosquitoes breed.

RECOMMENDATIONS

INSECT	FORMULATION	Directions
Mosquito Larvae	6-TGA 1%	Apply specified dosage per acre. Repeat as necessary. Use the product where ponds or water are present to root in 10 days.

RESTRICTIONS

1. Do not use inside homes, barns, areas, poultry houses, or other areas where food is processed.
2. Do not use on crops used for feed or forage.
3. Do not use treated manure on soil in which food crops are grown.