

RED-TOP

RESIDUAL MOSQUITO LARVICIDE
(Contains 2% DDT)

For Use Only by Mosquito Abatement Districts, Public Health Officials,
and Other Trained Personnel of Public Mosquito Control Programs.

(left panel)

ACTIVE INGREDIENT:
O,O-Dimethyl O-[(p-(methylthio)-m-tolyl)] phosphorothioate - - 2%
INERT INGREDIENTS - - - - - 98%
100%

*Reg. U.S. Pat. Offs. by Farbwerke Bayer A.G.,
Chemagro Corporation Licensee.
U. S. Patent No. 3,042,703

CAUTION (14 point type)

KEEP OUT OF THE REACH OF CHILDREN (10 point type)

Harmful if swallowed, inhaled, or absorbed through the skin.

Do not get in eyes, on skin, or on clothing.

Do not breathe dust.

Wash thoroughly with soap and warm water after handling.

Wash clothing with soap and hot water before re-use.

ANTIDOTE: Administer atropine sulfate in large therapeutic doses.
Repeat as necessary to the point of tolerance. Liberal doses of
emetics appear to enhance the antidotal properties of atropine.

SEE ALSO THE FOLLOWING USE PRECAUTIONS

Apply continuously on food and feed.
Be toxic to fish and wildlife. Do not use where fish, birds or other
wildlife are present. This product is highly toxic to sea, exposed
coastal areas. Do not use on crops. Protective measures
should be taken to protect sensitive agricultural areas.

ACCEPTED

MAY 1966

1966

Control of Mosquito Larvae
(with DDT - Baygon and Dieldrin)

DIRECTIONS FOR USE

DDT (DIBENYL) is recommended for use against mosquito larvae. It provides long residual control and is effective against DDT-resistant strains of non-resistant strains.

For mosquito larvae control, application should be made to moist areas, standing water, shallow ponds, and around the edges of larger ponds, lakes, marshes, swamps, and tidal areas. Also apply around and on garbage dumps, manure piles, catch basins, and similar areas where mosquitos breed.

RECOMMENDATIONS

INSECT	POUNDS PER ACRE	REMARKS
Mosquito Larvae	2½ to 5	Apply specified dosage per acre uniformly. Repeat as necessary. Use the higher rate where pools of water are greater than one foot in depth.

RESTRICTIONS

1. Do not use inside homes, dairy barns, poultry houses, or buildings where food is processed.
2. Do not use on crops used for seed or forage.
3. Do not use treated manure on soil in which food crops are grown.