- READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.
- Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.
- USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:
 - GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Deciduous Fruit Crops—60 to 800 gals. Citrus—400 to 3,000 gals. On young Fruit Plantings—Use a minimum of 15 gals. water per acre)
 - AIR APPLICATION—Vegetable and Field Crops—3 to 10 gals. Orchard Crops—10 to 20 gals.
- APPLES: EARLY DELAYED DORMANT ONLY—Green & Rosy Apple Aphids, European Red Mite, San Jose Scale, Fruit Tree Leafroller—1/4 pt. plus 2 gals. VOLCK 70 Supreme (1 1/2 gals. VOLCK Supreme Spray in Western Strates) per 100 gals. water as full coverage sprays. Do not combine oil sprays with Sulfur, ORTHOCIDE or PHALTAN. PRE-BLOSSOM and POST-BLOSSOM FIRST BROOD COVER SPRAYS—First Brood Codling Moth, Leafhoppers, Green and Rosy Apple Aphids, Leaf Miners—1/4 pt. per 100 gals. water Repeat at 7 to 10-day intervals (14 to 21 day intervals in the West). Repeat Application to Prevent Aphid Buildup—1/6 pt. per 100 gals. water. Do not use Phosphamidon on Apples within 30 days of harvest.
- BROCCOLI, CAULIFLOWER: Aphids, Imported Cabbageworm to 1 pint in water to cover one acre. Apply every 10 to 14 days in a preventative program for aphid control. Do not apply within 3 days of harvest.
- CANTALOUPE, WATERMELON: Aphids, Leaf Miner, Leafhopper 1/4 to 1/2 pt. per acre in water to cover. Do not apply within 1 day of harvest.
- COTTON (California and Arizona Only): Early Season (two through ten leaf stage)

 Aphids, Thrips, Fleahoppers, Spider Mites—3 fl. oz. per acre with ground equipment.

 Spray directly on young plants in sufficient water to cover. Make first application at the second true leaf stage. Mid and Late Season—Lygus, Flea Beetles, Leafhoppers—one half pt. per acre in water to cover, by ground or air. Aphids—one fourth pt. per acre by ground or air. Air Application—Use in one to five gals, water per acre. Do not apply more than three sprays: Do not use within 30 days of harvest. Do not graze treated cotton fields.

- CUCUMBERS: Aphids, Leaf Miner, Leafhopper—1/4 to 1/2 pt. per 100 gals. Water per acre. Do not apply within 3 days of harvest.
- TANGERINES (California and Arizona only), ORANGES, GRAPEFRUIT, LEMONS: Aphids, Thrips—1/4 to 1/2 pt. per 100 gals. water Ground Application—1/4 to 2/2 pts. per acre in water to cover. Air Application—1/2 to 3/4 pt. per acre in water to cover. Repeat as necessary. Apply during periods of aphid or thrips attack on new flush growth. Do not apply within 15 days of harvest.
- PEPPERS: Aphids, Leaf Miners, White Fly—1/2 pt. per acre in water to cover when insects appear European Corn Borer—1 pt. per acre in water to cover when infestations are expected. Do not apply within 6 days of harvest.
- POTATOES: Aphids, Colorado Potato Beetles, Leafhoppers, Leafminers, Flea Beetles, Stinkbugs—One half pt. per acre in water to cover by ground application (one to five gals. by air). Repeat as necessary. Do not apply within 14 days of harvest.
- TOMATOES: Aphids, Colorado Potato Beetles, Leaf Miner, White Fly Flea Beetles, Stinkbugs 1/2 pt. per acre in water to cover by ground application (1 to 5 gals. by air). Repeat as necessary. Do not apply within 10 days of harvest.
- WALNUTS: 'Aphids, Codling Moth—1 pt. per acre '(Ground—400 gals. water. Air—10 to 20 gals. water). Walnut Husk Fly—1 pt. per acre (Ground—200 to 300 gals. water. Air—10 to 20 gals. water), applying after egg stings first occur in the husks. Do not use within 7 days of harvest.

Contains phosphamidon manufactured by CIBA.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize my agent or representative to make, any warranty, guaranty or representation, exp., is or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability from improper handling, storage or use and accepts and uses this material on these conditions.

ANTIDOTE: If Swallowed. __
repeat until vomit fluid is
CIAN IMMEDIATELY! Note to
may be administered in coclething and immediately v

DANGER:

Wear a mask or respirator don protection. Rapidly Abs Wear natural rubber gloves mediately with soap and vexter before eating or smooth before re-use. Keep all unpermay be danger of drift. Note transport near feed or food FLAME. Do not use on Mapavoid drift to these varieties burying in non-crop lands a punching holes in them and possible drift to food, forag of rendered unfit for sale,

This product is toxic to wild where runoff is likely to or disposal of wates.

/This product is highly toxic the tive information may be

WHEN CONTAINER IS E

TANGER

LENIO

ACCEPTED

APR 1.0,1973

FUNGIONE AND ROUNTINDE ACT
FOR TOTAL FOR RESISTRATION SUBJECT

Phospierit H