

**RESTRICTED USE PESTICIDE**

Due to very high acute toxicity to Humans and Birds

For retail sale to and use only by certified applicator or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

**Wilbur-Ellis**

**ETHYL-METHYL PARATHION  
6-3 SPRAY**

**Directions For Use**

**ACTIVE INGREDIENTS:**

Parathion\* ..... 58.0%  
O, O-dimethyl O-p-nitrophenyl Phosphorothioate ..... 29.0%

**INERT INGREDIENTS** ..... 13.0%

\*O, O-diethyl O-p-nitrophenyl Phosphorothioate ..... 100.0%

(1 gallon contains 6 lbs. Parathion and 3 lbs. Methyl Parathion.)

EPA Reg. No. 2935-360

**KEEP OUT OF REACH OF CHILDREN**

**DANGER  
PELIGRO**



**POISON**

**OBSERVE PRECAUTIONS ON SAFETY LABEL**

**NOT FOR USE OR STORAGE IN OR AROUND THE HOME**

**DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM**

**ENDANGERED SPECIES RESTRICTIONS**

The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of federal laws. The use of this product is controlled to prevent death or harm to endangered or threatened species that occur in the following counties or elsewhere in their range.

Before using this pesticide in the following counties you must obtain the EPA Cropland Endangered Species Bulletin. The use of this pesticide is prohibited in these counties unless specified otherwise in the Bulletins. The EPA Bulletin is available from either your County Agricultural Extension Agent, the Endangered Species Specialist in your State Wildlife Agency Headquarters or the appropriate Regional Office of either the U.S. Fish and Wildlife Service (FWS) or the U.S. Environmental Protection Agency. **THIS BULLETIN MUST BE REVIEWED PRIOR TO PESTICIDE USE.**

**IF USED ON CROPS:**

**STATE (Regional Office FWS)  
Species**

**COUNTY**

**ALABAMA (Atlanta, GA)**

Slackwater darter

Alabama cavefish

Freshwater mussels

Lauderdale, Limestone, Madison


Lauderdale

Colbert, Jackson, Marshall, Morgan

**BEST AVAILABLE COPY**

<b>ARIZONA (Albuquerque, NM)</b>	
Woundfin	Mohave
Bonytail chub	Mohave
Gila topminnow	Graham, Santa Cruz
<b>ARKANSAS (Atlanta, GA)</b>	
Freshwater mussels	Clay, Clark, Cross, Lawrence, Randolph, Sharp, St. Francis
Ozark cavefish	Benton
Leopard darter	Sevier
<b>CALIFORNIA (Portland, OR)</b>	
Delta green ground beetle	Solano
Valley elderberry longhorn beetle	Merced, Sacramento
Aleutian Canada goose	Colusa, Merced, Sutter, Stanislaus
Kern primrose sphinx moth	Kern
Blunt-nosed leopard lizard	Fresno, Kern, Kings, Madera, Merced, Monterey, San Luis Obispo, Santa Barbara, Stanislaus, Tulare
Santa Cruz long-toed salamander	Monterey, Santa Cruz
Unarmored three-spine stickleback	Los Angeles, Santa Barbara
<b>COLORADO (Denver, CO)</b>	
Colorado squawfish	Blanco, Delta, Garfield, Mesa, Moffat, Rio, Routt
Humpback chub	Mesa
<b>FLORIDA (Atlanta, GA)</b>	
Everglade kite	Broward, Dade, Glades, Palm Beach
<b>KENTUCKY (Atlanta, GA)</b>	
Freshwater mussels	Ballard, Edmundson, Jackson, Laurel, Marshall, McCracken, Pulaski, Rockcastle, Warren, Wayne
<b>MARYLAND (Newton Corners, MA)</b>	
Maryland darter	Harford
<b>MISSISSIPPI (Atlanta, GA)</b>	
Bayou darter	Copiah
<b>MISSOURI (Twin Cities, MN)</b>	
Ozark cavefish	Christian, Greene, Jasper, Lawrence, Newton, Barry, Stone
<b>NEVADA (Portland, OR)</b>	
Woundfin	Clark
Pahrnagal Bonytail	Lincoln
Cui-ui	Washoe
Pahrump killifish	Clark, White Pine
<b>NEW MEXICO (Albuquerque, NM)</b>	
Pecos gambusia	Chaves, Eddy
<b>NORTH CAROLINA (Atlanta, GA)</b>	
Spotfin chub	Macon, Swain

PR63U-5



**BEST AVAILABLE COPY**

### ENDANGERED SPECIES (cont.)

STATE (Regional Office FWS) Species	COUNTY
OHIO (Twin Cities, MN) Scioto madtom	Champagne, Franklin, Logan, Madison, Pickaway, Union
OKLAHOMA (Albuquerque, NM) Leopard darter Ozark cavefish	Leflore, McCurtain Delaware
TENNESSEE (Atlanta, GA) Slackwater darter Slender chub Spotfin chub Freshwater mussels	Lawrence, Wayne Claiborne, Hancock Cumberland, Fentress, Morgan Blount, Claiborne, Decatur, Franklin, Hancock, Hardin, Lincoln, Loudon, Mar- shall, Maury, Rhea, Roane, Scott, Sequa- chie, Smith, Sullivan
TEXAS (Albuquerque, NM) Attwater's greater prairie chicken  Comanche Springs pupfish Pecos gambusia Texas blind salamander San Marcos salamander San Marcos gambusia Houston toad Fountain darter	Arkansas, Austin, Colorado, Fort Bend, Goliad, Refugio, Victoria Jeff Davis Jeff Davis, Pecos, Reeves Hays Hays Hays Bastrop, Burleson Hays
UTAH (Denver, CO) Woundfin Humpback chub Bonytail chub Colorado squawfish	Washington Grand Grand, Uintah Carbon, Duchesne, Emery, Garfield, Grand, Kane, San Juan, Uintah, Wayne
VIRGINIA (Newton Corners, MA) Spotfin chub Freshwater mussels	Scott, Washington Lee, Russell, Scott, Smyth, Tazewell, Washington, Wise

### DIRECTIONS FOR USE:

*It is a violation of Federal law to use this product in a manner inconsistent with its labeling.*

**GROUND APPLICATION:** Vegetable and Field Crops use 30 to 250 gals. of water per acre. Use 3 to 20 gals. of water per acre on Cotton. Where recommended for air application: Use 3 to 10 gals. of water per acre.

## RECOMMENDATIONS FOR USE:

**ALFALFA, CLOVER, GRASS and VETCH:** For aphids, alfalfa weevils, alfalfa caterpillars, armyworms, cutworms and grasshoppers, use  $\frac{1}{4}$  to  $\frac{1}{2}$  pint per acre. Do not apply within 15 days of harvest.

**SEED ALFALFA:** Use 1 pint per acre by air or ground in sufficient water to provide thorough coverage to control aphid, lygus bug and stink bug. Crop residues cannot be grazed or used as feed for livestock of any type.

**COTTON:** For control of boll weevil and bollworm apply at rate of  $1\frac{1}{3}$  pints per acre. For control of aphids, cotton leaf perforator, armyworms up to third instar, cabbage looper and stink bugs, apply at rate of 1 pint per acre. For control of cotton leafworm, fleahopper, lygus bugs, red spider mite and thrips, apply at rate of  $\frac{2}{3}$  pint per acre. Applications to cotton should be made at 4 or 5-day intervals until control is obtained.

**CAUTION:** When necessary to enter treated cotton fields within 48 hours after application, protective clothing should be worn. Do not apply within 7 days of harvest or of hand picking.

**SMALL GRAINS, (Wheat, Oats, Barley):** For aphids (greenbugs), armyworms, fall armyworms and winter grain mites, use  $\frac{1}{3}$  pint per acre and for grasshoppers, use  $\frac{1}{2}$  pint per acre. Do not apply within 15 days of harvest.

**SOYBEANS:** For stink bugs, three-cornered alfalfa hopper and velvet bean caterpillar use  $\frac{1}{3}$  pint per acre. Do not apply within 21 days of harvest.

**SUGAR BEETS:** For alfalfa loopers, aphids, armyworms, blister beetles, flea beetles, leafhoppers, leafminers, lygus bugs, stink bugs and webworms, use  $\frac{1}{2}$  pint per acre. Do not apply within 21 days of harvest. If tops are to be used for feed, do not apply within 60 days of harvest.

### VEGETABLES:

**BEANS (Dry and Green):** For control of aphids, armyworms, leafhoppers, leaf rollers, and red spider mites, use  $\frac{1}{2}$  to  $\frac{2}{3}$  pints per acre. Do not use within 21 days of harvest.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER and KOHLRABI:** For control of aphids, imported cabbage worms, diamond back moths and vegetable weevils, use  $\frac{1}{3}$  pint per acre. Do not apply within 21 days of harvest.

**CARROTS:** For aphid and vegetable weevils, use  $\frac{1}{2}$  pint per acre. Do not apply within 15 days of harvest. Do not use treated tops for food or feed.

**CELERY:** For control of aphids, celery leaf tiers, celery worms and spider mites, use  $\frac{1}{2}$  pint per acre. Do not apply within 21 days of harvest.

**CORN (Field):** For controlling aphids, budworms, european corn borer, armyworms and rootworms, use  $\frac{1}{2}$  pint per acre. Do not apply within 12 days of harvest or cutting.

**LETTUCE (Leaf, Bibb, Head):** For control of aphids, cabbage loopers, leafhoppers and thrip, use  $\frac{1}{3}$  to  $\frac{1}{2}$  pint per acre. Do not apply within 21 days of harvest.

**PEAS (Garden and Blackeyed):** For control of aphids and pea weevils use  $\frac{1}{2}$  pint per acre. Do not apply within 15 days of harvest.

**POTATOES:** For controlling aphids, blister beetles, leafhoppers, leafminers, plant bugs, potato psyllids, thrips and vegetable weevils, use  $\frac{1}{4}$  to  $\frac{1}{2}$  pint per acre. Do not apply within 5 days of harvest.

**TABLE BEETS:** For aphids, blister beetles and webworms use  $\frac{1}{2}$  pint per acre. Do not apply within 15 days of harvest; 21 days if tops are used for food.

**TURNIPS:** For aphids, imported cabbage worms and vegetable weevils, use  $\frac{1}{3}$  pint per acre. Do not apply within 15 days of harvest; 21 days if tops are used for food.

**NOTICE:** The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

**NOTICE:** The above are the only uses registered for use in California. An expanded label is available but is only for use in other states.