

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION: Vegetable and Field Crops: 50 to 100 gals. Cotton: 3 to 20 gals.

AIR APPLICATION: 3 to 10 gallons.

RECOMMENDATIONS FOR USE:

ALFALFA and VETCH: Use 1/2 pt. per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, and SPIDER MITES. Do not apply within 15 days of harvest or pasturing.

CARROTS, CELERY, PEAS and PEPPER: Use 1 pint per acre to control ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS, and SPIDER MITES. Do not apply within 15 days of harvest. Do not use the material on plants for food or feed.

BEANS, CABBAGE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use 1 1/2 to 2 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES and SPIDER MITES. Do not apply within 21 days of harvest. (May be applied within 15 days of harvest on dry beans.)

BEETS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops are to be used for food).

COTTON: Use 3 to 4 1/2 pints per acre to control APHIDS, BOLL WEEVIL, COTTON LEAFWORM, FLEAHOPPER, THRIPS, COTTON LEAF PERFORATOR, ARMYWORMS (up to 3rd instar) and SPIDER MITES. Do not apply within 15 days of harvest or during bloom. Workers entering cotton fields within 24 hours of application should wear protective clothing.

LETTUCE and SPINACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS, and SPIDER MITES. Do not apply within 15 days of harvest.

POTATOES: Use 1 1/2 pints per acre to control APHIDS, ARMYWORM (up to 3rd instar), FLEABEETLES, LEAFHOPPERS, and SPIDER MITES. Do not apply within 5 days of harvest.

SMALL GRAINS (Barley, Oats, Rye and Wheat): Use 1 pint per acre to control APHIDS (Greenbug), ARMYWORMS (up to 3rd instar), and FLEABEETLES. Do not apply within 15 days of harvest.

SORGHUM (Milo) (California Only): Use 1 1/2 pints per acre to control GREENBUG. Leaf injury may occur on plants that are not fully developed. Spraying plants as more severe injury may occur. Do not apply within 15 days of harvest.

SUGAR BEETS: Use 1 1/2 pints per acre to control ARMYWORMS, WEBWORMS and FLEABEETLES. Do not apply within 15 days of harvest or within 60 days of harvest if tops are to be used for food.

TOMATOES: Use 1 qt. per acre to control APHIDS, ARMYWORMS and CABBAGE MOPIES. Do not apply within 15 days of harvest.

red-top

METHYL PARATHION 5 SPRAY

ACCEPTED
By Weight
4.5%
0.5%
5.0%
100.0%

ACTIVE INGREDIENTS:

O, O-Dimethyl O-p-nitrophenyl thiophosphate
Xylene

INERT INGREDIENTS

EPA Reg. No. 2925-363-AA

KEEP OUT OF REACH OF CHILDREN!

DANGER: Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Methyl Parathion Poisoning. See Antidote Statement and Other Warning on Side Panel.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. Wear clean, natural rubber gloves, clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 15 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

WILBUR-ELLIS
APR 14 1974
FRESNO

WILBUR-ELLIS COMPANY
FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

METHYL PARATHION 5 SPRAY

ACCEPTED

ACTIVE INGREDIENTS

O, O-Dimethyl O-p-nitrophenyl thiophosphate
Xylene

By Weight
4.5%
0.5%
5.0%
100.0%

INERT INGREDIENTS

EPA Reg. No. 2935-363-AA

KEEP OUT OF REACH OF CHILDREN !

Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Methyl Parathion Poisoning. See Antidote Statement and Other Warning on Side Panel.

DO NOT USE, POUR, SPILL OR STOVE NEAR HEAT OR OPEN FLAME.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use.

INTERNAL: If there is any possibility of ingestion, swallow immediately. If it may be absorbed by the mouth, wash out of the mouth by gargling with warm salt water (1 table-spoonful of salt per quart of water). Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet.

EXTERNAL: If there is any possibility of absorption, immediately remove patient from the vicinity of the Methyl Parathion 5 Spray, remove contaminated clothing and wash skin with soap and running water. If it has gotten into the eyes, wash immediately with running water for at least ten minutes.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

DANGER IN USING

Do not use in enclosed spaces. Do not use in the presence of children, pregnant women, or nursing infants. Do not use in the presence of livestock or other animals. Do not use in the presence of bees, butterflies, or other beneficial insects. Do not use in the presence of fish or other aquatic life. Do not use in the presence of birds or other wild animals. Do not use in the presence of domestic animals. Do not use in the presence of pets. Do not use in the presence of children. Do not use in the presence of pregnant women or nursing infants. Do not use in the presence of livestock or other animals. Do not use in the presence of bees, butterflies, or other beneficial insects. Do not use in the presence of fish or other aquatic life. Do not use in the presence of birds or other wild animals. Do not use in the presence of domestic animals. Do not use in the presence of pets.

NOT FOR USE OR STORAGE IN OR NEAR THE HOME

NOTICE: The state of California has determined that this pesticide is highly toxic to humans and animals. It is therefore classified as a "restricted use" pesticide. It may only be used by or under the direct supervision of a person who is licensed by the State of California as a pesticide applicator. For more information on this pesticide, contact the California Department of Pesticide Regulation, 4005 La Jolla Village Drive, San Diego, California 92161. For more information on this pesticide, contact the California Department of Pesticide Regulation, 4005 La Jolla Village Drive, San Diego, California 92161.

APR 1974
STRESS REGULATORY
EPA