

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician for an adequate supply of 1-100 grain atropine tablets for emergency use. Atropine is also an antidote for Methyl Parathion and may be administered in conjunction with Atropine.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 table-spoon of salt to a glass of water). Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

DANGER IN USING: Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Treated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate food or feed products. Keep away from heat and open flame. To prevent reuse, empty containers should be decontaminated and destroyed. Decontaminate by washing thoroughly with a strong solution of a lye and detergent and rinse with water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your agricultural extension service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply within 100 feet of water. Do not apply within 100 feet of water. Do not contaminate water by leaking of equipment or disposal of wastes. Apply this product only as specified on this label.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

EPA Reg. No. 2935 363-AA

BEST DOCUMENT AVAILABLE

RESTRICTED USE PESTICIDE:

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

red-top
METHYL PARATHION 5 SPRAY

ACTIVE INGREDIENTS:		By Weight
O, O-Dimethyl-O-p-nitrophenyl Phosphorothioate		54.5%
Xylene Range Aromatic Solvent		40.5%
INERT INGREDIENTS		5.0%
		100.0%

KEEP OUT OF REACH OF CHILDREN !

DANGER: Poisonous if Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Methyl Parathion Poisoning. See Antidote Statement and Other Warning on Side Panel.



POISON

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. Wear clean, natural rubber gloves, clean, waterproof protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 15 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

WILBUR-ELLIS COMPANY
FRESNO • LOS ANGELES • GLENDALE, ARIZ. • PORTLAND • SEATTLE

DIRECTIONS FOR USE:

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION: Vegetable and Field Crops: 30 to 250 gals Cotton: 3 to 20 gals
AIR APPLICATION: 3 to 10 gallons.

RECOMMENDATIONS FOR USE:

ALFALFA and VETCH: Use 1/2 pt. per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMY WORMS (up to 3rd instar), LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest or pasturing.

CABBOTS, CELERY, PEAS and PEPPERS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed.

BEANS, CABBAGE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use 1 1/2 to 2 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES and SPIDER MITES. Do not apply within 21 days of harvest. (May be applied within 15 days of harvest on dry beans.)

BEETS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops are to be used for food).

COTTON: Use 3 to 4 1/2 pints per acre to control APHIDS, BOLL WEEVIL, COTTON LEAFWORM, FLEAHOPPER, THRIPS, COTTON LEAF PERFORATOR, ARMYWORMS (up to 3rd instar) and SPIDER MITES. Do not apply within 7 days of hand picking or harvest. Workers entering cotton fields within 24 hours after treatment should wear protective clothing.

LETTUCE and SPINACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar) IMPONED CABBAGE WORM, FLEABEETLES, LEAFHOPPERS and SPIDER MITES. For CABBAGE LOOPER control use 1 3/5 pints per acre. Do not use within 21 days of harvest.

POTATOES: Use 1 1/2 pints per acre to control APHIDS, ARMYWORM (up to 3rd instar), FLEA BEETLES, LEAFHOPPERS, and SPIDER MITES. Do not apply within 5 days of harvest.

SMALL GRAINS (Barley, Oats, Rye and Wheat): Use 1 pint per acre to control APHIDS (Greenbug), ARMYWORMS (up to 3rd instar) and LEAFHOPPERS. Do not apply within 15 days of harvest.


SORGHUM (Milo) (California Only): Use 1 1/2 pts. per acre to control GREENBUG. Leaf injury may occur on some sorghum varieties. Do not use wetting agent as more severe injury may occur. Do not apply within 21 days of harvest.

SUGAR BEETS: Use 3/5 pt. per acre to control APHID, ARMYWORM, WERWORMS and FLEA BEETLES. Do not apply within 20 days of harvest or within 20 days of harvest if tops are to be fed to livestock.

TOMATOES: Use 1 qt per acre to control APHID, ARMYWORMS and CABBAGE LOOPER. Do not use within 15 days of harvest.

EPA Est. 2935 CA 1 F 67c

ACCEPTED



METHYL PARATHION MIXTURE, LIQUID