

Reg H 2735-305 P.14
PM 12

173015
17/03
300

JUL 11 1986

Ms. Terry Miller
Wilbur-Ellis Company
Agricultural Services Corporate Office
191 West Shaw Avenue, Suite 107
Fresno, CA 93704-2876

Dear Ms. Miller:

Subject: Revision of Label Ingredient Statement
Red-Top Methyl Parathion 5 Spray
EPA Registration No. 2935-363
Your Application Dated April 11, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Xylene Range Aromatic Solvent does not need to be listed as an inert ingredient in your label ingredient statement.

Sincerely yours,

Lawrence J. Schnaubelt
Lawrence J. Schnaubelt
Acting Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

92963.Schnaubelt.173015.E-11-KENCO:6/5/86:6/11/86:PAR:LF
CONCURRENCE

SYMBOL ▶							
SURNAME ▶							
DATE ▶							

PRECAUTIONARY STATEMENTS

DANGER: Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly absorbed through skin. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to Methyl Parathion poisoning. See Antidote Statement and Other Warning on Side Panel.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. Wear clean, natural rubber gloves; clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 15 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. "2-Pam" is also an antidote for Methyl Parathion and may be administered in conjunction with Atropine.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your agricultural extension service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

MPRS-4

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators under their direct supervision and only for those uses covered by the Applicator's certification.

RED-TOP METHYL PARATHION

ACTIVE INGREDIENTS:

O, O-Dimethyl-O-p-nitrophenyl phosphorothioate

INERT INGREDIENTS:

Xylene Range Aromatic Solvent 40.5%

EPA Reg. No. 2935-363-AA

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**



POISON

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

See Side Panels for Antidote and First Aid Treatment

Wilbur-Ellis

P. O. Box 16458, Fresno, CA 93716

DIRECTIONS FOR USE:

is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. not enter treated areas for 48 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing (see left), areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment (see left) in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Danger. Area treated with Methyl Parathion on (date of application). Do not enter without appropriate protective clothing for 48 hours. (Insert here actions to take in case of accidental exposure - see left)."

STORAGE AND DISPOSAL

Prohibitions — Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Storage: Store in original container only and keep sealed. Store in closed storage areas. Keep away from heat and open flames. Use caution when moving, opening, losing, or pouring. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

SIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE LESS OTHERWISE NOTED:

GROUND APPLICATION: Vegetable and Field Crops: 30 to 250 gals. Cotton: 3 to 20 gals.

APPLICATION: 3 to 10 gallons.

DIRECTIONS FOR USE (Continued):

RECOMMENDATIONS FOR USE:

ALFALFA and VETCH: Use $\frac{1}{2}$ pt. per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMYWORMS (Up to 3rd instar), LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest or pasturing.

CARROTS, CELERY, PEAS and PEPPERS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed.

BEANS, CABBAGE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use $1\frac{1}{2}$ to 2 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES and SPIDER MITES. Do not apply within 21 days of harvest. (May be applied within 15 days of harvest on dry beans.)

BEETS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops are to be used for food).

COTTON: Use 3 to 4 $\frac{1}{2}$ pints per acre to control APHIDS, BOLL WEEVIL, COTTON LEAFWORM, LEAFHOPPER, THRIPS, COTTON LEAF PERFORATOR, ARMYWORMS (up to 3rd instar) and SPIDER MITES. Do not apply within 7 days of hand picking or harvest. Workers entering cotton fields within 24 hours after treatment should wear protective clothing.

LETTUCE and SPINACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), IMPORTED CABBAGE WORM, FLEABEETLES, LEAFHOPPERS and SPIDER MITES. For CABBAGE LOOPER control use 1 $\frac{3}{5}$ pints per acre. Do not use within 21 days of harvest.

POTATOES: Use $1\frac{1}{2}$ pints per acre to control APHIDS, ARMYWORM (up to 3rd instar), FLEABEETLES, LEAFHOPPERS, and SPIDER MITES. Do not apply within 5 days of harvest.

SMALL GRAINS (Barley, Oats, Rye and Wheat): Use 1 pint per acre to control APHIDS (Greenbug), ARMYWORMS (up to 3rd instar) and LEAFHOPPERS. Do not apply within 15 days of harvest.

SORGHUM (Milo) (California Only): Use $1\frac{1}{2}$ pts. per acre to control GREENBUG. Leaf injury may occur on some sorghum varieties. Do not use wetting agent as more severe injury may occur. Do not apply within 21 days of harvest.

SUGAR BEETS: Use $\frac{3}{5}$ pt. per acre to control APHID, ARMYWORMS, WEBWORMS and FLEABEETLES. Do not apply within 20 days of harvest or within 60 days of harvest if tops are to be fed to livestock.

TOMATOES: Use 1 qt. per acre to control APHID, ARMYWORMS and CABBAGE LOOPERS. Do not use within 15 days of harvest.

F-584

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on those conditions.

In Case of Emergency



METHYL PA

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 11 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2935-363

DIRECTIONS FOR USE (Continued):

IONS FOR USE:

ETCH: Use 1/2 pt. per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMY-
Jp to 3rd instar), LEAFHOPPERS and SPIDER MITES. Do not apply within 15
rvest or pasturing.

RY, PEAS and PEPPERS: Use 1 pint per acre to control APHIDS, ARMYWORMS
1 instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within
f harvest. Do not use treated carrot tops for food or feed.

RE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use 1 1/2 to 2 pints per
viral APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES and SPIDER MITES.
ly within 21 days of harvest. (May be applied within 15 days of harvest on

int per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES,
RS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops
used for food).

to 4 1/2 pints per acre to control APHIDS, BOLL WEEVIL, COTTON LEAFWORM,
R, THRIPS, COTTON LEAF PERFORATOR, ARMYWORMS (up to 3rd instar) and
ES. Do not apply within 7 days of hand picking or harvest. Workers entering
s within 24 hours after treatment should wear protective clothing.

NACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar),
CABBAGE WORM, FLEABEETLES, LEAFHOPPERS and SPIDER MITES. For CABBAGE
viral use 1 3/5 pints per acre. Do not use within 21 days of harvest.

1 1/2 pints per acre to control APHIDS, ARMYWORM (up to 3rd instar), FLEA-
AFHOPPERS, and SPIDER MITES. Do not apply within 5 days of harvest.

[Barley, Oats, Rye and Wheat]: Use 1 pint per acre to control APHIDS
ARMYWORMS (up to 3rd instar) and LEAFHOPPERS. Do not apply within
harvest.

o) (California Only): Use 1 1/2 pts. per acre to control GREENBUG. Leaf injury
on some sorghum varieties. Do not use wetting agent as more severe injury
Do not apply within 21 days of harvest.


se 3/5 pt. per acre to control APHID, ARMYWORMS, WEBWORMS and FLEA
s not apply within 20 days of harvest or within 60 days of harvest if tops
ed to livestock.

1 qt. per acre to control APHID, ARMYWORMS and CABBAGE LOOPERS. Do
hin 15 days of harvest.

F-584

ents made on this label, are believed to be true and accurate, but because of conditions of use
r control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or rep-
any warranty, guaranty or representation, express or implied, concerning this material or the
in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the
responsible in any manner for any personal injury or property damage or loss resulting to the
r person from handling, storage or use of this material, not in accordance with directions. The
sk and liability resulting from improper handling, storage or use and accepts and uses this
ditions.

In Case of Emergency, Call CHEMTREC: (800) 424-9300



**METHYL PARATHION MIXTURE, LIQUID
NA 2783**

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 11 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2935-363