

Reg # 2935-355 1972

CAUTIONARY STATEMENTS

DANGER: Poisonous if Swallowed, Inhaled or Absorbed Through Skin. Rapidly absorbed through skin. Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to Methyl Parathion poisoning. See Antidote Statement and Other Warning on Side Panel.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. Wear clean, natural rubber gloves; clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 15 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. "2-Pam" is also an antidote for Methyl Parathion and may be administered in conjunction with Atropine.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your agricultural extension service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**NOT FOR USE OR STORAGE IN OR AROUND THE HOME.
DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME
MPRS-4**

RESTRICTED USE PESTICIDE:

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

RED TOP WILBUR-ELLIS

METHYL PARATHION 5 SPRAY

ACTIVE INGREDIENTS:	By Weight
O, O-Dimethyl-O-p-nitrophenyl Phosphorothioate	54.5%
Xylene-Range-Aromatic-Solvent	40.5%
INERT INGREDIENTS	5.0%
includes Xylene	40.5%
EPA Reg. No. 2935-363-AA	100.0%

Est. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**



POISON APR 0 6 1987

PRECAUTION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

See Side Panels for Antidote and Precautions

Wilbur-Ellis

P. O. Box 16458, Fresno, CA 93755

If is a violation of Federal law

Do not apply this product in or other persons. The area. Do not enter treated areas for certain states may require this product, consult your. Written or oral warnings must be or in an area about to be tective clothing (see left), period of time area is to in case of exposure. When customarily understood by believe that written warning following information: "Do cation). Do not enter with actions to take in case of

- 1. Prohibitions — Do not con-
- 2. Pesticide Disposal: Pesticid pesticide, spray mixture, c not be disposed of by use or Environmental Control / EPA Regional office for gu Container Disposal: Triple ing, or puncture and dispo by state and local authorit
- 4. Storage: Store in original areas. Keep away from h-closing, or pouring. Dece- trated solution of alkali c

USE DESIGNATED AMOUNT OF UNLESS OTHERWISE NOTED GROUND APPLICATION: 1/2- AIR APPLICATION: 3 to 10

BEST AVAILABLE COPY

1
2

DIRECTIONS FOR USE:

Violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Apply this product in such a manner as to directly or through drift expose worker other persons. The area being treated must be vacated by unprotected persons. After treated areas for 48 hours unless appropriate protective clothing is worn. Because certain situations may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Oral warnings must be given to workers who are expected to be in a treated area in an area about to be treated with this product. Oral warnings should specify protective clothing (see left), areas or fields that cannot be entered without this clothing and of time area is to be vacated, and statement of practical treatment (see left case of exposure. When oral warnings are given, warnings shall be given in a language commonly understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Danger. Area treated with Methyl Parathion on (date of application). Do not enter without appropriate protective clothing for 48 hours. (Insert directions to take in case of accidental exposure - see left.)"

STORAGE AND DISPOSAL

Containers — Do not contaminate water, food or feed by storage or disposal. Spill Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest Regional office for guidance. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Storage: Store in original container only and keep sealed. Store in closed storage areas. Keep away from heat and open flames. Use caution when moving, opening, cleaning, or pouring. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

IGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE

LESS OTHERWISE NOTED:
GROUND APPLICATION: Vegetable and Field Crops: 30 to 250 gals. Cotton: 3 to 20 gals
AERIAL APPLICATION: 3 to 10 gallons.

PASTURE (Not for use in Calif.): Use 3/4 pint per acre to control EUROPEAN CRANE FLIES. Do not apply after grass growth is over 6" high. Do not apply within 15 days of pasturing or harvesting.

DIRECTIONS FOR USE (Continued):

RECOMMENDATIONS FOR USE:


ALFALFA and VETCH: Use 1/2 pt. per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMYWORMS (up to 3rd instar), LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest or pasturing. SPOTTED ALFALFA: Use 1 qt. per acre to control Aphids by grass and clover. Use 1/2 qt. per acre for grazing of feed for livestock of any kind.
CABBAGE, CELERY, PEAS and PEPPERS: Use 1/2 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABETTER, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed. Do not apply within 15 days of harvest. (May be applied within 15 days of harvest on dry beans.)
BEANS, CABBAGE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use 1 1/2 to 2 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABETTER and SPIDER MITES. Do not apply within 21 days of harvest. (May be applied within 15 days of harvest on dry beans.)
BEETS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABETTER, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops are to be used for food).
COTTON: Use 3 to 4 1/2 pints per acre to control APHIDS, BOLL WEEVIL, COTTON LEAFWORM, FLEAHOPPER, THRIPS, COTTON LEAF PERFORATOR, ARMYWORMS (up to 3rd instar) and SPIDER MITES. Do not apply within 7 days of hand picking or harvest. Workers entering cotton fields within 24 hours after treatment should wear protective clothing.
LETTUCE and SPINACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), IMPORTED CABBAGE WORM, FLEABETTER, LEAFHOPPERS and SPIDER MITES. For CABBAGE LOOPER control use 1 3/5 pints per acre. Do not use within 21 days of harvest.
POTATOES: Use 1 1/2 pints per acre to control APHIDS, ARMYWORM (up to 3rd instar), FLEABETTER, LEAFHOPPERS, and SPIDER MITES. Do not apply within 5 days of harvest.
SMALL GRAINS (Barley, Oats, Rye and Wheat): Use 1 pint per acre to control APHIDS (Greenbug), ARMYWORMS (up to 3rd instar) and LEAFHOPPERS. Do not apply within 15 days of harvest.
SORGHUM (Milo) (California Only): Use 1 1/2 pts. per acre to control GREENBUG. Leaf injury may occur on some sorghum varieties. Do not use wetting agent as more severe injury may occur. Do not apply within 21 days of harvest.
SUGAR BEETS: Use 3/5 pt. per acre to control APHID, ARMYWORMS, WEBWORMS and FLEABETTER. Do not apply within 20 days of harvest or within 60 days of harvest if tops are to be fed to livestock.
TOMATOES: Use 1 qt. per acre to control APHID, ARMYWORMS and CABBAGE LOOPERS. Do not use within 15 days of harvest.

F-584
NOTICE: The statements made on this label, are believed to be true and accurate, but because of variations of use which are beyond our control, WILBUR-ELI'S COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELI'S COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

SUNFLOWERS (Not for use in Calif.): Use 1-2 pints per acre to control stem weevils, seed weevils, and sunflower beetles. Repeat application 2-3 times at 5 day intervals if needed. Do not apply at onset of flowering or before 10% of plants begin to flower.

GRAPES: Use 1 1/5 pints per acre to control GRAPE LEAF FOLDER, LARGE TORTRIX, MEALYBUG, and RED SPIDER MITES. Use 125 to 150 gallons of water by ground and 5 to 10 gallons of water by air. Apply once per season. Do not apply within 14 days of harvest.

In Case of Emergency, Call CHEMTREC: (800) 424-9300



METHYL PARATHION MIXTURE, LIQUID
NA 2783

RICE: Use 1/3 to 1 pint per acre to control RICE LEAFMINER and TADPOLE SHRIMP. Hold water in check for 3 days after application. Do not apply within 15 days of harvest. Do not apply more than 1 pt. per acre.

BEST AVAILABLE COPY