

ACCEPTED

JUN 22 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under 2935-366-11 EPA Reg. No. 2935-366-11

RED-TOP SEVIN® 5 PELLETS

(CARBARYL BAIT)
FOR THE CONTROL OF INSECT PESTS

BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS

If swallowed: Call a physician immediately. Give a glass or two of water and induce vomiting by touching back of throat with finger. It is preferable to induce vomiting under medical supervision or to use gastric lavage with a cuffed endotracheal tube because of aspiration hazard. Remove victim immediately to emergency treatment facilities.

If on skin: Remove by washing with soap and water. Get medical attention.

If in eyes: Flush the eyes with clean water for 10 minutes. Get medical attention.

If inhaled: Remove victim to fresh air. Transport immediately to emergency treatment facility.

Note To Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Do Not Use: 2-PAM opiates, or cholinesterase inhibiting drugs.

Environmental Hazards:

Avoid direct application to lakes, streams and ponds. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE OR STORAGE IN OR AROUND HOME.

Read Entire Label. Use Strictly According To Label Directions and Cautions. Do not use application methods, dosages, concentrations, or frequencies not listed on labeling. Do not apply against target pests not listed on labeling. Do not mix with fertilizers.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions.

If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed, harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

STORAGE AND DISPOSAL

- 1. PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2. PESTICIDE DISPOSAL:** Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.
- 3. CONTAINER DISPOSAL:** Dispose of bags according to approved Federal, State, or local procedures under the Resource Conservation and Recovery Act.
- 4. GENERAL:** Consult Federal, State or Local Disposal Authorities for additional restrictions and approved alternatives.

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20-40 pounds per acre of Bait will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sawbugs.

In Case of Emergency, Call CHEMTREC: (800) 424-9300

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthylmethyl Carbamate)

INERT INGREDIENTS

SEVIN® is the registered trademark of Union Carbide Corporation for Carbaryl Insecticide.

EPA Reg. No. 2935-366-AA

Est. No. 2935-CA-1

By Wt.

9%

95%

TOTAL

100%

KEEP OUT OF REACH OF CHILDREN!

CAUTION:**Hazards To Humans and Domestic Animals:**

Harmful if Inhaled or Swallowed. Avoid Breathing of Dust. Do Not Take Internally. Avoid Contact with Skin and Eyes.

Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTION STATEMENTS

RECOMMENDATIONS FOR USE

Crop	Dosage Limitations To Meet Residue Tolerances	Days Between Final Application and Harvest
Cucumbers, Squash, Melons, Peanuts	20 pounds per acre	None
Alfalfa, Cotton, Peas	30 pounds per acre	None
Beans, Carrots, Corn, Eggplants, Grapes, Okra, Peppers, Potatoes, Pumpkins, Tomatoes, Sweet Corn	40 pounds per acre	None
Asparagus, Strawberries	40 pounds per acre	1 Day
	Repeat applications to Asparagus should be made at intervals of at least 3 days apart. Apply to Asparagus during cutting season.	
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce, Garden Beets (roots), Horseradish, Parsnips, Radishes, Rutabagas, and Turnips (roots)	40 pounds per acre	3 Days
Raspberries (Not for use in Calif.)	40 pounds per acre	7 Days
Celery (Not for use in Calif.), Collards, Endive (escarole), Garden Beets (tops), Kale, Leaf Lettuce, Parsley, Spinach, Sugar Beets, Swiss Chard, and Turnips (tops)	40 pounds per acre	14 Days
Sorghum	40 pounds per acre	21 Days