 <div> United States Environmental Protection Agency Washington, DC 20460 </div>	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number 222715
--	--	---------------------------------


Application for Pesticide - Section I		
1. Company/Product Number 2935-366	2. EPA Product Manager D. Edwards	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Sevin 5 Bait	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) WILBUR-ELLIS COMPANY 191 W. Shaw Avenue, Suite 107 Fresno, CA 93704 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II	
<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of deleted uses: sweet sorghum, barley, oats and rye.

Section - III			
1. Material This Product Will Be Packaged in:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt No. per container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____			

Section - IV		
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name JOANNE C. STRIEBICH	Title Registration and Labeling Specialist	Telephone No. (Include Area Code) (209) 226-1934
<div> Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. </div>		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration and Labeling Specialist	
4. Typed Name JOANNE C. STRIEBICH	5. Date March 19, 1996	



SEVIN® 5 BAIT

(CARBARYL BAIT)
FOR THE CONTROL OF INSECT PESTS
N-METHYL CARBAMATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Harmful if inhaled or swallowed. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin and clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: A) Long sleeved shirt; B) Long pants; C) Waterproof gloves; D) Shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

Carbaryl is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is A) Coveralls; B) Waterproof gloves; C) Shoes plus socks.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Carbaryl residues when Carbaryl is used in accordance with label directions.

If Carbaryl insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

STORAGE AND DISPOSAL

PROHIBITIONS. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty containers. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to positioning the container in storage, storage temperature, potential for crushing or damage due to stacking and penetration of moisture.

STORAGE. Store in a safe manner. Store in original container only. Store in a cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang can affect package strength. Personnel should use clothing or equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

RECOMMENDATIONS FOR USE

Broadcast applications may be made with either ground equipment or aircraft, under conventional, minimal and no till practices.

Treatments may be repeated as necessary. Where a range of rates is provided, use lower rates for light infestation, early instars or when foliage is not too dense, use higher rates for moderate to heavy infestation, advanced instars or when foliage is thick. For best results on lawn and turf, cut lawn, remove clippings and water thoroughly. Dosages from 20-48 pounds of bait per acre will control cutworms, armyworms, field crickets, house crickets, mole crickets, darkling beetles, grasshoppers, scabbards and ants.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

NOTE: Do not make more than two applications after grain heads of wheat, alfalfa, and clover emerge from boot. Do not allow animals to graze on treated sunflower crops.

Crop	Dosage Limitations To Meet Residue Tolerances	Days Between Final Application and Harvest
Cucumbers, Melons, Peanuts, Pumpkins, Squash	20 lbs. per acre (1% a.i. per 1000 sq. ft.)	0
Burdockroot (Taro) (Orange and Hay), Clover, Wheat (Pasture and Forage)	30 lbs. per acre	0
Lawn and Turf	30 lbs. per acre (1.1 a.i. per 1000 sq. ft.)	0
GROUND APPLICATION ONLY		
Pasture (Rangeland and Wasteland), Forage Grasses, Fence Row and Non-Crop Areas	20-40 lbs. per acre	0
Beans, Carrots, Corn (Field, Sweet, Pop), Eggplant, Okra, Peppers, Potatoes, Sweet Potatoes and Turnips	20-40 lbs. per acre (2% 14.25 a.i. per 1000 sq. ft.)	0
Soybeans	20-40 lbs. per acre	0
Sorghum Forage	20-40 lbs. per acre	0

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)

By Wt.

INERT INGREDIENTS

TOTAL

5%

95%

100%

EPA Reg. No. 2935-366-AA

EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN!

CAUTION:

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. Give a glass or two of water and induce vomiting by touching back of throat with finger. It is preferable to induce vomiting under medical supervision or to use gastric lavage with auffed endotracheal tube because of aspiration hazard. Remove victim immediately to emergency treatment facilities.

If on Skin: Remove by washing with soap and water. Get medical attention.

If in eyes: Flush the eyes with clean water for 10 minutes. Get medical attention.

If inhaled: Remove victim to fresh air. Transport immediately to emergency treatment facility.

Note To Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do Not Use 2-PAM opiates or cholinesterase inhibiting drugs.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

RECOMMENDATIONS FOR USE (CONT.)

Crop	Dosage Limitations To Meet Residue Tolerances	Days Between Final Application and Harvest
Tobacco Beds	20-40 lbs. per acre	3
Peas	30 lbs. per acre (1.1 a.i. per 1000 sq. ft.)	7
Alfalfa	30 lbs. per acre	7
Asparagus, Strawberries	40 lbs. per acre (14.25 a.i. per 1000 sq. ft.)	1
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cawpeas, Garden Beets (Beets), Head Lettuce, Horseradish, Kohlrabi, Parsnips, Radishes, Rutabagas, Salsify (Roots), Turnips (Roots)	20-40 lbs. per acre (7% 14.25 a.i. per 1000 sq. ft.)	3
Tobacco (Field)	20-40 lbs. per acre	7
Cotton	30 lbs. per acre	14
Celery, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden Beet Tops, Hanoover Salad, Kale, Leaf Lettuce, Mustard Greens, Parsley, Salsify (Tops), Spinach, Swiss Chard, Turnip Tops (CELERY NOT FOR USE IN CALIFORNIA)	20-40 lbs. per acre (7% 14.25 a.i. per 1000 sq. ft.)	14
Rice	20-40 lbs. per acre	14
<i>Do not use on rice fields in which crayfish and/or catfish farming is included in the cultural practice</i>		
Sugar Beets	20-40 lbs. per acre	14
Wheat (Grain - Fall or Spring)	30 lbs. per acre	21
Sorghum (Milo, Grain Sorghum, Sweet Sorghum and Hybrids)	20-40 lbs. per acre	21 (Grain)
*Cats, Rabbits and Hogs (all treatments)		
<i>Do not feed treated forage to dairy animals or animals being finished for slaughter or whose class is grain on crops treated with this product</i>		
Raspberries	40 lbs. per acre (14.25 a.i. per 1000 sq. ft.)	7
Sunflowers	20-40 lbs. per acre (7% 14.25 a.i. per 1000 sq. ft.)	50 (oil seed, or feed) 0 (seedling purposes only)
NOT FOR USE IN CALIFORNIA		
The following are for all insects listed above EXCEPT darkling beetles.		
Annual Plants	40-48 lbs. per acre	
Herbaceous Plants and Shrubs, Trees, Woody Plants (Ornamental)	14.25-17% a.i. per 1000 sq. ft.)	
Grapes	40-48 lbs. per acre (14.25-17% a.i. per 1000 sq. ft.)	7
Apples, Cherries, Peaches, Pears and Plums	40-48 lbs. per acre (14.25-17% a.i. per 1000 sq. ft.)	1
Apricots and Nectarines	40-48 lbs. per acre (14.25-17% a.i. per 1000 sq. ft.)	3
Citrus	40-48 lbs. per acre (14.25-17% a.i. per 1000 sq. ft.)	5

NOTICE. The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, yet in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accept, and uses this material on these conditions.

Sevin® is a registered trademark of Rhone-Poulenc for Carbaryl insecticide.
WILBUR-ELLIS logo® and IDEAS TO GROW WITH® a registered trademarks of WILBUR-ELLIS Company

IN CASE OF EMERGENCY, CALL CHEMTREC: (800) 424-9300

NET WEIGHT: 50 POUNDS

P.O. BOX 16458 - FRESNO, CA 93755