



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

April 9, 2025

Anne Downs  
Senior Registration Specialist  
Wilbur-Ellis Company LLC  
2903 S. Cedar Avenue  
Fresno, CA 93725

Subject: Registration Review Label Amendments Incorporating Mitigation from the National Marine Fisheries Services (NMFS) Biological Opinions on the Effects of Carbaryl  
Product Name: Sevin 5 Bait  
EPA Registration Number: 2935-366  
Application Date: 5/6/24  
Decision Number: 595803

Dear Anne Downs:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application of the above referenced product in connection with the NMFS' Biological Opinion (BiOp) on the effects of carbaryl on federally listed endangered and threatened species and designated critical habitat in accordance with section 7 of the Endangered Species Act (ESA) of 1973, as amended (16 U.S.C. 1531 et seq.). The Agency has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or


claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

The Agency received the U.S. Fish and Wildlife Service's (FWS) BiOp on the effects of EPA's national registration of carbaryl and posted it on the [website](#) on April 9, 2025. The Agency plans to incorporate and implement this FWS BiOp when issuing the interim decision for carbaryl. EPA will notify the registrants of any need to submit label amendments within 60 days of the issuance of the interim decision.

If you have any questions about this letter, please contact Carolyn Smith at [smith.carolyn@epa.gov](mailto:smith.carolyn@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Jaclyn Pyne". The signature is written in a cursive, flowing style.

Jaclyn Pyne, Team Leader  
Risk Management and Implementation Branch 3  
Pesticide Re-evaluation Division  
Office of Pesticide Programs

Enclosure

**ACCEPTED**

04/09/2025

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No.

2935-366



**INSECTICIDE**

**SEVIN® 5 BAIT**

For the control of cutworms, armyworms, field crickets, house crickets, mole crickets, darkling beetles, grasshoppers, sowbugs and ants (except fire ants, harvester ants, pharaoh ants and carpenter ants).

**ACTIVE INGREDIENT**

**BY WT.**

Carbaryl (1-naphthyl N-methylcarbamate) ..... 5.00%

**OTHER INGREDIENTS** ..... 95.00%

**TOTAL** ..... 100.00%

20 lb of SEVIN 5 BAIT per acre – 1 lb of A.I. per acre

EPA Reg. No. 2935-366

EPA Est. No. 66876-OR-001

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**FIRST AID**

Have product container or label with you when calling a poison control center or doctor or going for treatment. Poison Control Center: 1-800-222-1222

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes
- Call a poison control center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**Note to Physician:** Carbaryl is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**Caution:** Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are natural rubber, polyethylene or neoprene. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves, and
- Shoes plus socks.

In addition, handlers loading bait formulations into airplanes under the auspices of the Animal and Plant Health Inspection Service (APHIS) Rangeland Grasshopper and Mormon Cricket Suppression Program, and allied state programs must wear a NIOSH-approved dust/mist filtering respirator with NIOSH/MSHA approved number prefix TC-21C or a NIOSH-approved respirator with any N, R, P or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROLS

Pilots applying applications under the auspices of the Animal and Plant Health Inspection Service (APHIS) Rangeland Grasshopper and Mormon Cricket Suppression Program or related state programs must use an enclosed cockpit in a manner that is consistent with the WPS for Agricultural Pesticides [40 CFR 170.240(d)(6)]. In addition, the cockpit air intake must be located above the bait release point and the bait bin must be completely separate from the cockpit.

Flaggers supporting aerial applications permitted above must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] for dermal protection. In addition, flaggers must:

- wear long-sleeve shirt, long pants, shoes, and socks,
- either wear a NIOSH-approved dust/mist filtering respirator with NIOSH/MSHA approval number prefix TC-21C or a NIOSH-approved respirator with an N, R, P, or HE filter **OR** use an enclosed cab that is declared in writing by the manufacturer or by a government agency to provide at least as much respiratory protection as this type of respirator,
- be provided and have immediately available for use in an emergency when they must exit the cab in the treated area: chemical-resistant gloves and chemical-resistant headgear and, if using an enclosed cab that provides respiratory protection, a respirator of the type specified above,
- take off any PPE that was worn in the treated area before re-entering the cab, and
- store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

When applicators use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(5), the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

**For terrestrial uses:** Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

## DIRECTIONS FOR USE

### ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS

Before using this product, you must obtain any applicable Endangered Species Protection Bulletins ('Bulletins') within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at <https://www.epa.gov/pesticides/bulletins>. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area where you are applying the product, including any restrictions on application timing if applicable. It is a violation of Federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov).

### Reporting Ecological Incidents

For guidance on reporting ecological incidents, including death, injury, or harm to plants and animals, including bees and other non-target insects, see EPA's Pesticide Incident Reporting website: <https://www.epa.gov/pesticide-incidents> or call (720) 306-6340.

Do not apply during rain. Do not apply when soil in the area to be treated is saturated (if there is standing water on the field or if water can be squeezed from soil) or if NOAA/National Weather Service predicts a total rainfall of 1 inch or greater over the 48 hours following the day of application, only considering a 48-hour period when, at any point during the 48-hour period, the precipitation potential is 50% or greater. Detailed National Weather Service forecasts for local weather conditions should be obtained on-line at [www.weather.gov](http://www.weather.gov) or by contacting your local National Weather Service Forecasting Office.

## USE RESTRICTIONS

Do not apply this product for agricultural uses in Hawaii.

Do not apply by air for all uses except for USDA APHIS and affiliated State grasshopper and Mormon cricket suppression programs.

Do not product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply to vertical surfaces directly above pervious or impervious surfaces that drain into ditches, storm drains, gutters, or surface waters.

Do not apply or irrigate to the point of runoff.



## MANDATORY SPRAY DRIFT MANAGEMENT

### AERIAL APPLICATIONS:

(For USDA APHIS and affiliated State grasshopper and Mormon cricket suppression programs only)

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- During application, the Sustained Wind Speed, as defined by the National Weather Service (standard averaging period of 2 minutes) must register between 3 and 10 miles per hour.
- Wind speed and direction must be measured on location using a windsock, an anemometer, or an aircraft smoke system.
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- Applicators must use a minimum of ½ swath displacement upwind at the downwind edge of the treatment area. The boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.
- Do not apply by ground within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, or ephemeral streams when water is present, wetlands or natural ponds, estuaries, and commercial fish farm ponds).

### GROUND BOOM APPLICATIONS:

- During application, the Sustained Wind Speed, as defined by the National Weather Service (standard averaging period of 2 minutes), must register between 3 and 15 miles per hour (or 3 and 10 mph for the USDA APHIS and affiliated State grasshopper and Mormon cricket suppression programs).
- Wind speed and direction must be measured on location using a windsock or anemometer (including systems to measure wind speed or velocity using application equipment).
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- Do not release spray at a height greater than 2 feet above the ground or crop canopy except for USDA APHIS and affiliated State grasshopper and Mormon cricket suppression programs.
- For USDA APHIS and affiliated State grasshopper and Mormon cricket suppression programs, apply at the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply during temperature inversions.
- For soil or foliar applications, do not apply by ground within 25 feet of aquatic habitats (such as, but not limited to lakes, reservoirs, rivers, permanent streams, or ephemeral streams when water is present, wetlands or natural ponds, estuaries, and commercial fish farm ponds).

### USDA APHIS REDUCED AREA AND AGENT TREATMENTS (RAATS):

The RAATs approach takes advantage of grasshopper movement to allow SEVIN 5 BAIT to be applied at reduced rates on a reduced treated area while maintaining acceptable protection against grasshopper. Under this program, Sevin 5 Bait may be applied to as little as 50% of the infested area (treating a 100 ft. swath, skipping a 100 ft. swath) up to 100% of infested area. The amount of area treated will depend on grasshopper age, density and plant canopy. By heaving untreated swaths, the RAATs program provides reserves for natural biological control agents. This strategy fully utilizes an Integrated Pest Management approach for grasshopper. Apply Sevin 5 Bait by air or ground or both depending on current labeling on 50 – 100% of infested area for grasshopper nymphs between the 2<sup>nd</sup> and 5<sup>th</sup> instar. The rate to use per acre will depend on grasshopper age, population density, and plant canopy. Use the higher rates on more mature grasshoppers, severe infestations and dense vegetation. Use the lower rate on younger grasshoppers, light to moderate infestations and sparse vegetation. Consult with the local grasshopper control experts in your area for the proper rate and swath width to use under your conditions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions, and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the "Restricted-entry interval (REI). The REI for carbaryl is 12 hours unless otherwise specified in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, are:

- Coveralls over short-sleeve shirt and short-pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant headgear if overhead exposure

When the Restricted-Entry Interval for a crop is 7 days or longer, you must notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

### **IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION**

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Carbaryl residues when Carbaryl is used in accordance with label directions.

If Carbaryl insecticide is used in accordance with label directions, carrot tops, sugar beet tops and corn forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

### **APPLICATION METHOD**

This product may be applied only by ground using a mechanical spreader or underground shank applicator. Do not apply by hand, spoon or belly grinder. **EXCEPTION:** Aerial applications only to rangeland are allowed only under the auspices of the Animal and Plant Health Inspection Service (APHIS) Rangeland Grasshopper and Mormon Cricket Suppression Program, and/or allied state programs.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, SEVIN 5 BAIT contains a Group 1A insecticide. Any insect population may contain individuals naturally resistant to SEVIN 5 BAIT and other Group 1A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of SEVIN 5 BAIT or other Group 1A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC): 1. Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species. 2. Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management. 3. When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s). 4. Mixtures become less

effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits. 5. The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance, contact Wilbur-Ellis Company LLC representatives at (720) 306-6340.

Commodity	Use Rate
Asparagus	<p><u>Application to Spears:</u> Apply 20 lbs. per acre per application with a maximum yearly usage of 60 lbs. per acre.</p> <p><u>Applications to Ferns (after harvest of spears):</u> Apply 40 lbs. per acre per application with a maximum yearly usage of 80 lbs. per acre.</p> <p>•<b>IMPORTANT:</b> Total combined maximum yearly usage for spears and ferns is 100 lbs. per acre.</p> <p>•Retreatment interval = 7 days</p> <p>•Do not apply within 1 day of harvest.</p> <p>•Do not make more than 4 applications to spears. Do not make more than 2 applications to ferns.</p>
Garden Beets (roots), Carrots, Horseradish, Radish, Parsnip, Rutabaga, Salsify, Turnips (roots)	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 120 lbs. per acre.</p> <p>•Retreatment interval = 7 days.</p> <p>•Do not apply within 7 days of harvest.</p> <p>•Do not make more than 3 applications per year.</p>
Garden Beets (tops), Turnips (tops)	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 120 lbs. per acre.</p> <p>•Retreatment interval = 7 days.</p> <p>•Do not apply within 14 days of harvest.</p> <p>•Do not make more than 3 applications per year.</p>
Chinese Cabbage, Collards, Kale, Mustard Greens	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 120 lbs. per acre.</p> <p>•Make applications only within 30 days of crop emergence or transplanting.</p> <p>•Retreatment interval = 7 days.</p> <p>•Do not apply within 14 days of harvest.</p> <p>•Do not make more than 3 applications per year.</p>
Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kohlrabi	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 120 lbs. per acre.</p> <p>•Make applications only within 30 days of crop emergence or transplanting.</p> <p>•Retreatment interval = 7 days.</p> <p>•Do not apply within 3 days of harvest</p> <p>•Do not make more than 3 applications per year.</p>



Commodity	Use Rate
Corn (field & pop)	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 160 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 14 days.</li> <li>•Do not apply within 14 days of harvest (forage, silage).</li> <li>•Do not apply within 48 days of harvest (grain, fodder).</li> <li>•Do not make more than 4 applications per year.</li> <li>•Restricted-entry interval (REI) = 24 hours.</li> <li>•<b>EXCEPTION:</b> the REI is 21 days for workers detasseling corn. Hand harvesting is prohibited.</li> </ul>
Corn (sweet)	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 160 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment = 3 days.</li> <li>•Do not apply within 2 days of harvest (ears).</li> <li>•Do not apply within 14 days of harvest (forage).</li> <li>•Do not apply within 48 days of harvest (fodder).</li> <li>•Do not make more than 4 applications per year.</li> <li>•Restricted-entry interval (REI) = 24 hours.</li> <li>•<b>EXCEPTION:</b> the REI is 21 days for workers detasseling corn. Hand harvesting is prohibited.</li> </ul>
Cucumber, Melon, Squash	<p>Apply 20 lbs. per acre per application with a maximum yearly usage of 80 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 7 days.</li> <li>•Do not apply within 3 days of harvest.</li> <li>•Do not make more than 4 applications per year.</li> </ul>
Pistachios (Non-bearing trees only)	<p>Only for use on trees that will not bear fruit for one year from the date of application.</p> <p>Apply 40 lbs. per acre per application with a maximum yearly usage of 200 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 7 days.</li> <li>•If weed or cover crop bloom is present, mow orchard or between rows prior to applying this product.</li> </ul>
Potato	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 120 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 7 days.</li> <li>•Do not apply within 7 days of harvest.</li> <li>•Do not make more than 3 applications per year.</li> </ul>
Sugar Beets	<p>Apply 30 lbs. per acre per application with a maximum yearly usage of 60 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 14 days.</li> <li>•Do not apply within 28 days of harvest (roots, forage).</li> <li>•Do not make more than 2 applications per year.</li> </ul>
Tomato, Pepper, Eggplant	<p>Apply 40 lbs. per acre per application with a maximum yearly usage of 160 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 7 days.</li> <li>•Do not apply within 3 days of harvest.</li> <li>•Do not make more than 4 applications per year.</li> </ul>
Pasture & grasses (grown for hay or seed)	<p>Apply 30 lbs. per acre per application with a maximum yearly usage of 60 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 14 days.</li> <li>•Do not apply within 7 days of harvest or grazing.</li> <li>•Do not make more than 2 applications per year.</li> </ul>

Commodity	Use Rate
Rangeland	
Ground application only	<p>Apply 20 lbs. per acre.</p> <ul style="list-style-type: none"> <li>• Do not make more than one application per year.</li> </ul>
USDA APHIS Program for Mormon Cricket and Grasshopper Suppression	<p>Apply 4 lbs. per acre per application with a maximum yearly usage of 8 lbs. per acre.</p> <ul style="list-style-type: none"> <li>•Retreatment interval = 14 days</li> <li>•Do not make more than 2 applications per year.</li> <li>•Do not apply within 14 days of grazing</li> <li>•Use of a flaky cereal grain bait is prohibited.</li> </ul> <p>Aerial application is allowed only through USDA APHIS and affiliated state grasshopper and Mormon Cricket suppression programs.</p>

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container only and keep sealed. Store in a cool, dry and closed storage area. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Use caution when moving, opening, closing or pouring.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:**

**NONREFILLABLE CONTAINER:** Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

**ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.**

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors

Seller warrants that this product conforms to the chemical description on the label.

**EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.**

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

INSECTICIDE® logo and WILBUR-ELLIS® logo are registered trademarks of Wilbur-Ellis Company LLC.

All other trademarks cited here are the property of their respective owners.

**NET WEIGHT:** \_\_\_\_\_

**Batch Code**\_\_\_\_\_

2025-0321



**ATTENTION:** Cancer and Reproductive Harm • [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)

**WILBUR-ELLIS COMPANY LLC**  
2903 S. Cedar Avenue  
Fresno, CA 93725  
559-442-1220



**WILBUR-ELLIS<sub>x</sub>**