TOXAPHENE 40 - WP

A WETTABLE POWDER FOR USE IN WATER AS AN INSECTICIDE

DIRECTIONS FOR USE: Use designated amount of product in the following gallons of water per acre unless otherwise noted.

GROUND APPLICATION: Vegetables and Field Crops: 20 to 250 gals. **Cotton:** 5 to 20 gals.

AIR APPLICATION: 5 to 10 gals.

Dosage rates are given in terms of Toxaphene 40 WP per acre.

Measure the specified amount of this product, then wash through filler screen in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of [illegible]

[illegible] combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants. Emulsifiable liquids should always be thoroughly emulsified with at least half of total water before adding this product.

ACTIVE INGREDIENT:

Toxaphene (Technical Chlorinated Camphene) (Chlorine content 67% to 69%)	**40%**
INERT INGREDIENTS:	**60%**
	100%

EPA Reg. No. 2935 381 AA

WARNING: KEEP OUT OF REACH OF CHILDREN!

[illegible] **May be absorbed through skin. Do not get in eyes, on skin or on clothing. Do not breathe dust or spray mist. Wash thoroughly after using. Do not contaminate food and feed.**

BURN OR BURY EMPTY CONTAINER - KEEP OUT OF SMOKE

OBSERVE THE FOLLOWING USE PRECAUTIONS

RECOMMENDATIONS FOR USE:

BEANS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER

CELERY

EGGPLANT

GRASS RANGE: [illegible]

SPRUCE

PEPPERS

TOMATOES

NOTICE: The statements made in this label are believed to be [illegible] WILBUR-ELLIS COMPANY does not make [illegible] authorize any agent or representative to make any warranty [illegible] WILBUR-ELLIS COMPANY [illegible] held responsible in any manner for any personal injury [illegible] resulting from improper handling, storage or use and [illegible] this material on these conditions.

TOXAPHENE 40 - WP

A WETTABLE POWDER FOR USE IN WATER AS AN INSECTICIDE

DIRECTIONS FOR USE: Use designated amount of product in the following gallons of water per acre unless otherwise noted.

GROUND APPLICATION: Vegetables and Field Crops: 20 to 250 gals. **Cotton:** 5 to 20 gals.

AIR APPLICATION: 5 to 10 gals.

Dosage rates are given in terms of Toxaphene 40 WP per acre.

Measure the specified amount of this product, then wash through filter screen in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to [illegible]

combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants. Emulsifiable liquids should always be thoroughly emulsified with at least half of total water before adding this product.

RECOMMENDATIONS FOR USE:

[illegible]

BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER

[illegible]

ACTIVE INGREDIENT:	
Toxaphene (Technical Chlorinated Camphene)	40%
(Chlorine content 67% to 69%)	
INERT INGREDIENTS:	60%
	100%

EPA Reg. No. 2935 381 AA

WARNING: KEEP OUT OF REACH OF CHILDREN!

[illegible] **or on clothing. Do not breathe dust or spray mist. Wash thoroughly after using. Do not contaminate [illegible] and feed.**

BURN OR BURY EMPTY CONTAINER - KEEP OUT OF SMOKE.

OBSERVE THE FOLLOWING USE PRECAUTION:

[illegible]