

DIRECTIONS FOR USE: Do not mix this material with other materials not compatible with oil. Observe precaution when applying oil-based insecticides to crops during high temperatures since some crops are then more susceptible to injury. Do not apply within 14 days of sulfur or sulfur containing sprays. Consult your county agent for proper dosages when applying by aircraft.

Use designated amount of product in the following gallons of water per acre unless otherwise noted: **Ground Application** - Field Crops - 30 to 250 gals. **Air Application** - Field Crops - 3 to 10 gals.

FORAGE CROPS: On alfalfa, grass, soybeans, apply at the rate of 2 - 3 quarts (1 to 1½ lbs. of active) per acre for the control of alfalfa weevil (larvae), alfalfa webworm, fall armyworm, pea weevil, soybean caterpillar and velvet bean caterpillar. (May be used in conjunction with malathion for alfalfa weevil control.)

Mix with water (both high and low volume spraying) to give correct coverage per acre. Start applications at first sign of infestation and repeat at 7 - 10 day intervals as needed for control. Do not apply later than 7 days before harvest or grazing.

LIVESTOCK (Beef Cattle & Pigs): For control of hornflies, mix 1 quart in 12½ gallons of water and apply two (2) quarts of spray to each mature animal. For control of lice, apply as drenching spray. For control of hog lice, mix 1 quart in 12½ gallons of water and apply as a drenching spray.

DO NOT USE ON DAIRY ANIMALS OR IN DAIRY BARNS.

DO NOT USE IN DIPPING VATS.

SHADE TREES, ORNAMENTALS:

For control of Japanese beetles, leafhoppers, tent caterpillars, cankerworms and rose chafer, apply at the rate of 2 - 3 quarts (1 - 1½ lbs. active) in 100 gallons of water (3 - 5 tablespoons in 1 gallon of water). Spray foliage thoroughly. Repeat applications every 7 - 14 days.

ACTIVE INGREDIENT:

Methoxychlor, technical*

25.0%

INERT INGREDIENTS:

75.0%

100.0%

*Equivalent to 22% 2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 3% other isomers and related compounds

EPA Reg. No. 2935-00385-AA

CAUTION: KEEP OUT OF REACH OF CHILDREN!

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of spray from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONTAINER DISPOSAL: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

WARNING - FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME. WASH AFTER HANDLING. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. AVOID BREATHING OF SPRAY MIST. AVOID CONTAMINATION OF FEED OR FOODSTUFFS. HARMFUL IF SWALLOWED.

DIRECTIONS FOR USE (Continued):

For elm bark beetle (assists in control of Dutch elm disease), apply at the rate of 4 gallons per 100 gallons of water. Spray trunk and limbs thoroughly. Repeat every 7 - 14 days thru growing season.

PEAS: For control of pea weevil and armyworm, use 3 - 4 quarts (1½ to 2 lbs. active) per acre. Do not apply within 7 days of harvest or grazing.

NOTICE: The statements made on this label, are believed to be true and accurate, but, because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use of this material.