

red-top

DIAZINON[®] 4 SPRAY

For control of insects on fruits, nuts, vegetables, field crops, and ornamentals

Do Not Re-use Empty Container. Destroy it by perforating or crushing or returning drum to manufacturer. Bury or discard perforated or crushed metal and container in a safe area away from water supplies.

WARNING: KEEP OUT OF REACH OF CHILDREN!

May be fatal if swallowed. May be fatal if inhaled. May be fatal if absorbed through skin. May irritate eyes, nose, throat, and skin. Do not get in eyes, on face, or on clothes. Do not breathe spray. Do not get on skin or clothes. Do not get in mouth. Do not get on hands. Do not get on face. Do not get on hair. Do not get on neck. Do not get on chest. Do not get on back. Do not get on arms. Do not get on legs. Do not get on feet. Do not get on shoes. Do not get on socks. Do not get on underwear. Do not get on pajamas. Do not get on nightgown. Do not get on bathrobe. Do not get on towel. Do not get on blanket. Do not get on bedspread. Do not get on pillow. Do not get on mattress. Do not get on rug. Do not get on carpet. Do not get on floor. Do not get on wall. Do not get on ceiling. Do not get on furniture. Do not get on drapery. Do not get on curtain. Do not get on shade. Do not get on blind. Do not get on screen. Do not get on door. Do not get on window. Do not get on handle. Do not get on knob. Do not get on latch. Do not get on lock. Do not get on key. Do not get on coin. Do not get on button. Do not get on zipper. Do not get on fastener. Do not get on hook. Do not get on eyelet. Do not get on ring. Do not get on stud. Do not get on screw. Do not get on bolt. Do not get on nut. Do not get on washer. Do not get on gasket. Do not get on seal. Do not get on O-ring. Do not get on grommet. Do not get on plug. Do not get on cap. Do not get on lid. Do not get on cover. Do not get on case. Do not get on box. Do not get on container. Do not get on package. Do not get on label. Do not get on sticker. Do not get on tape. Do not get on string. Do not get on cord. Do not get on wire. Do not get on cable. Do not get on hose. Do not get on pipe. Do not get on tube. Do not get on hose. Do not get on pipe. Do not get on tube. Do not get on hose. Do not get on pipe. Do not get on tube.

ACTIVE INGREDIENTS:

O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothiate 48.72%

Xylene 39.00%

INERT INGREDIENTS 12.28%

Net Weight: 4.0 lbs. (1.81 kg) per gallon

~~Company Name: Wilbur-Ellis Company~~
E-14

USE ONLY ACCORDING TO DIRECTIONS ON ACCOMPANYING SHEET

Not for use on food crops. Do not use on ornamentals. Do not use on lawns. Do not use on golf courses. Do not use on tennis courts. Do not use on swimming pools. Do not use on hot tubs. Do not use on spas. Do not use on saunas. Do not use on showers. Do not use on bathtubs. Do not use on toilets. Do not use on sinks. Do not use on drains. Do not use on pipes. Do not use on faucets. Do not use on valves. Do not use on handles. Do not use on knobs. Do not use on latches. Do not use on locks. Do not use on keys. Do not use on coins. Do not use on buttons. Do not use on zippers. Do not use on fasteners. Do not use on hooks. Do not use on eyelets. Do not use on rings. Do not use on studs. Do not use on screws. Do not use on bolts. Do not use on nuts. Do not use on washers. Do not use on gaskets. Do not use on seals. Do not use on O-rings. Do not use on grommets. Do not use on plugs. Do not use on caps. Do not use on lids. Do not use on covers. Do not use on cases. Do not use on boxes. Do not use on containers. Do not use on packages. Do not use on labels. Do not use on stickers. Do not use on tape. Do not use on string. Do not use on cord. Do not use on wire. Do not use on cable. Do not use on hose. Do not use on pipe. Do not use on tube. Do not use on hose. Do not use on pipe. Do not use on tube.



WILBUR-ELLIS COMPANY



FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

Transplant Water (Drench) Application: Mix 1/2 gal. of transplant water when 1/2 gal. of water are used per plant. May be applied at the rate of 1/2 gal. of spray per 1000 plants. In case of the plant, the spray should be applied to the soil at least at time of planting.

Radishes: Mix 1 qt. of water. Apply 80-100 gals. per acre (1/2 gal. per 1,000 linear feet of row) at planting time.

WIREWORMS: ~~Brassica, Brussels Sprouts, Cabbage, Cantaloupe, Carrots, Cauliflower, Celery, Collards, Cucumbers, Eggplant, Kale, Lettuce, Lima Beans, Musk Melons, Onions, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Corn, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelon, Winter Squash.~~

Broadcast 3-4 qts. per acre prior to planting, and work into soil 4-6 inches immediately.

Potatoes: Wireworms: Broadcast 3 qts. per acre prior to planting. Immediately disk or harrow into soil 4-6 inches.

FIELD AND FORAGE INSECTS

Red-Top Diazinon 4 Spray should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil, depending on the type of application.

ALFALFA and CLOVER: Spotted Alfalfa Aphid, Yellow Clover Aphid, Pea Aphid, Leafhoppers, Grasshopper, Armyworm, Cutworm, Caterpillar, etc.

Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to alfalfa or clover in bloom. Do not apply to alfalfa or clover in bloom. Do not apply to alfalfa or clover in bloom.

CORN: European Cornborer, Corn Earworm, etc.

Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to corn in bloom. Do not apply to corn in bloom.

PROTEIN LARVAE CONTROL: European Cornborer, Corn Earworm, etc.

Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to protein larvae in bloom. Do not apply to protein larvae in bloom.

FRY BEETLES: Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to fry beetles in bloom. Do not apply to fry beetles in bloom.

GRASSHOPPERS: Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to grasshoppers in bloom. Do not apply to grasshoppers in bloom.

FRY BEETLES: Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to fry beetles in bloom. Do not apply to fry beetles in bloom.

GRASSHOPPERS: Apply 2-4 qts. per acre, depending on the type of application. Repeat applications may be necessary. Do not apply to grasshoppers in bloom. Do not apply to grasshoppers in bloom.

ACCEPTED

Red-Top

MAR 16 1973

RECEIVED

2935-383

DIAZINON[®] 4 SPRAY

For control of insects on fruits, nuts, vegetables, field crops, and ornamentals.

ACTIVE INGREDIENTS:	
O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	48.72%
Xylene	39.00%
INERT INGREDIENTS	
	12.28%
TOTAL	100.00%

Red-Top Diazinon 4 Spray contains 4 lbs. D in each gallon.

Calif. Reg. No. 2935-50103-AA

WARNING: KEEP OUT OF REACH OF CHILDREN!

May be fatal if swallowed. May be absorbed through the skin. Do not breathe dust or mist. Do not get in eyes, on face, or on clothing. Wash thoroughly after handling. Avoid contact with food and animals. Do not use in or near homes, schools, or day care centers. Do not use in or near children's play areas. Do not use in or near water. Do not use in or near swimming pools.

Do not use in or near homes, schools, or day care centers.

DIRECTIONS FOR USE: Use designated amount of product in the following gallons of water per acre unless otherwise noted:

soil using a disk tiller or cultivator immediately after spraying. Application should be made at the first sign of corn rootworm larvae, but not later than June 24 in the middle Corn Belt or July 1 in the northern Corn Belt.

SUGGESTED RATES ACCORDING TO ROW SPACING FOR CORN ROOTWORM LARVAE CONTROL

Row Spacing	Amount of Red Top Diazinon P-4 Spray per Acre	Planting Time application	Post emergence basal treatment
30 inches	23.55 oz.	14.28 oz.	15.43 oz.
36 inches	24.74 oz.	14.88 oz.	16.03 oz.
42 inches	25.93 oz.	15.48 oz.	16.63 oz.
48 inches	27.12 oz.	16.08 oz.	17.23 oz.
54 inches	28.31 oz.	16.68 oz.	17.83 oz.
60 inches	29.50 oz.	17.28 oz.	18.43 oz.
66 inches	30.69 oz.	17.88 oz.	19.03 oz.
72 inches	31.88 oz.	18.48 oz.	19.63 oz.
78 inches	33.07 oz.	19.08 oz.	20.23 oz.
84 inches	34.26 oz.	19.68 oz.	20.83 oz.
90 inches	35.45 oz.	20.28 oz.	21.43 oz.
96 inches	36.64 oz.	20.88 oz.	22.03 oz.
102 inches	37.83 oz.	21.48 oz.	22.63 oz.
108 inches	39.02 oz.	22.08 oz.	23.23 oz.
114 inches	40.21 oz.	22.68 oz.	23.83 oz.
120 inches	41.40 oz.	23.28 oz.	24.43 oz.

NOTE: For row spacings not listed, calculate the appropriate dosage rates based on the 10-inch row spacing.

See Corn Maggots - 2.4 pts. per acre just prior to planting. Work into soil 2 inches in depth.

Cutworms - Surface and Subterranean - Broadcast 2.4 pts. per acre for surface cutworms, apply just prior to planting or when larvae first appear and work into soil 1-2 inches immediately. Do not apply later than 1 day before subterranean cutworms, apply just prior to planting and work into soil 3-4 inches immediately.

Wireworms - Apply 3-4 pts. per acre prior to planting, and work into the soil 4-6 inches in depth.

Corn Rootworm - Adults - Apply 1 pt. per acre.

Corn Leaf Aphid - Apply 2 pts. per acre in sufficient water to provide complete coverage of the upper portions of plants. Repeat application as necessary.

Mites - Apply 1 pt. per acre in sufficient water to achieve thorough coverage when infestation first appears and repeat applications as necessary.

Flea Beetles - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Grasshoppers - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Sap Beetles - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Stink Bugs - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Pinworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

Tomato Rootworm - Apply 1 pt. per acre in sufficient water to assure thorough coverage when insects first appear and repeat application as necessary.

mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not permit children or pets to go on sprayed grass until spray has completely dried.

See directions sheet for use recommendations.

FISH AND WILDLIFE PRECAUTIONS - This product is toxic to fish and wildlife. Do not apply to water bodies or near water. Do not apply to areas where waterfowl are present.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT RE-USE EMPTY CONTAINER. Destroy it by perforating or crushing or returning drum to reconditioner. Bury or discard perforated or crushed drums and containers in a safe place away from water supplies.

Do not use of larvae or roots of open fields.

WARNING - FLAMMABLE - KEEP AWAY FROM FIRE - KEEP AWAY FROM CHILDREN.

Diazinon is a registered trademark of CIBA-GEIGY Corporation.

DIRECTIONS FOR USE: Use designated amount of product in the following gallons of water per acre unless otherwise noted:

GROUND APPLICATION - Vegetable and Field Crops - 30 to 50 gal. Cotton - 3 to 20 gallons. Deciduous Fruit Crops - 20 to 800 gal. Citrus - 400 to 1,000 gal. (On Young Fruit Plantings - Use a minimum of 15 gal. water per acre). Where Recommended for AIR APPLICATION - FIELD CROPS - 3 to 10 gal. Orchard Crops - 10 to 20 gal.

GENERAL INFORMATION - Red Top Diazinon P-4 spray is a systemic soil and contact insecticide which, when applied to the soil, is taken up by the plant for the control of insect pests of agricultural crops. Do not use on lawns or dwellings. Do not use on livestock.

In the absence of a specific recommendation, Red Top Diazinon P-4 Spray should be mixed with water in the following proportions: 1 pt. per acre for most outdoor use, and 1/2 pt. per acre for indoor use. Apply in a uniform manner.

W E WILBUR-ELLIS COMPANY W E
 FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

RECOMMENDATIONS FOR USE:

FRUIT AND NUT CROPS

ALMONDS: Mites, Twig Borers: Apply 1 1/2 pts. per 100 gals. of water.
Pantano Scales, San Jose Scales: Mix 1 1/2 pts. of zinc plus 2-3 gals. dormant oil per 100 gals. of water and apply as a dormant spray. Do not apply more than 12 pts. per acre.
Almond hulls may be used for feed for livestock.

APRICOTS: Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale, Crawlers, San Jose Scale Crawlers: Mix 1 pt. per 100 gals. of water. Apply when aphids, mites, and scale crawlers first appear.

Olive Scale Crawlers: Mix 1/2 pt. of Red-Top Diazinon 4 Spray plus 1 1/2 gals. light medium kerosene oil per 100 gals. of water. Apply when scale crawlers are present.

Apricot Mealybugs: Mix 1 pt. per 100 gals. of water. Apply as a cover spray from petal fall to June.

Do not apply within 10 days of harvest. Do not apply more than 8 pts. per acre.

CANE BERRIES (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries): Leafhoppers, Thrips, Aphids, Two-spotted Mites, Raspberry Sawflies: Mix 1 pt. per 100 gals. of water and apply up to 200 gals. of spray per acre, or apply 2 pts. per acre in sufficient water for thorough coverage.

Raspberry Fruitworms: Apply at the close rates when blossom buds separate and repeat application immediately before blossoms open.

Dryberry Mites: Mix 2 pts. per 100 gals. water. Apply as a dormant spray in early spring when canes are still on the ground and buds are closed. Make a second application when the buds are well developed, but still closed.

Do not apply within 10 days of harvest. Do not apply more than 2 pts. per acre.

Raspberry Green Scale: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray from petal fall to June. Do not apply more than 12 pts. per acre.

CHERRIES: Eye gnats, Red weevils, Fruit Tree Leaf Rollers: Mix 1 pt. per 100 gals. of water. Apply as a cover spray from petal fall to June.

Leafhoppers: Apply 1/2 pt. per 100 gals. of water in 3-5 applications beginning with petal fall to June.

Cherry Fruit Flies: Mix 1/2 pt. per 100 gals. of water. Make 3 or 4 applications beginning with petal fall to June, at 7-10 day intervals.

Cherry Aphids: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when aphids first appear.

Cherry Sawflies: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when sawflies first appear.

Cherry Mites: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when mites first appear.

Cherry Scale: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when scale first appears.

CITRUS: Citrus Leaf Miners, Citrus Fruit Tree Leaf Rollers, Cottony Cushion Scale Crawlers, Citrus Aphids: Mix 1/2 pt. per 100 gals. of water. Apply as a cover spray when insects first appear.

Citrus Mites: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when mites first appear.

Citrus Scale: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when scale first appears.

Citrus Thrips: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when thrips first appear.

Citrus Leafhoppers: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when leafhoppers first appear.

Citrus Fruit Flies: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when fruit flies first appear.

Citrus Mealybugs: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when mealybugs first appear.

Citrus Aphids: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when aphids first appear.

Citrus Mites: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when mites first appear.

Citrus Scale: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when scale first appears.

Citrus Thrips: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when thrips first appear.

Citrus Leafhoppers: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when leafhoppers first appear.

Citrus Fruit Flies: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when fruit flies first appear.

Citrus Mealybugs: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when mealybugs first appear.

Citrus Aphids: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when aphids first appear.

Citrus Mites: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when mites first appear.

Citrus Scale: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when scale first appears.

Citrus Thrips: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when thrips first appear.

CARROTS: Aphids: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when aphids first appear.

CELERY: Aphids: Apply 1/2 pt. per 100 gals. of water. Apply as a cover spray when aphids first appear.

COLLARDS, KALE, SWISS CHARD, and TURNIP TOPS: Aphids, Cabbageworms, Diamond-back Moths, Harlequin Cabbage Bugs, Corn Earworms: Apply 1/2 pt. per acre. Do not apply to collards, kale, or turnip tops within 10 days of harvest.

CUCUMBERS: Aphids, Serpentine Leaf Miners, Thrips: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

LETTUCE and ENDIVE: Aphids, Dipterous Leaf Miners: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

MELONS and SQUASH: Cantaloupes, Casabas, Greenshaws, Honeydew Melons, Muskmelons, Persian Melons and hybrids of these, Watermelons and their hybrids, Summer and Winter Squash: Aphids, Leafhoppers, Mites: Apply 1/2 pt. per acre. Serpentine Leaf Miners, Thrips: Apply 1/2 pt. per acre. Do not apply to melons and winter squash within 3 days, or to summer squash within 7 days, of harvest.

MUSHROOM HOUSES: Mushroom Flies, Phorids and Sclerids: Mix 1/2 gal. in 50 gals. of water (0.5% to 1.0% actual Diazinon) and spray on outside walls and tent stais of mushroom houses, after compost in beds has been pasteurized by heating. Do not spray compost in beds or after spawning.

To treat wooden sides of beds, posts, doors, and door frames after pasteurizing, mix 1/2 pt. in 6 gals. of water and apply with a brush.

Do not treat compost.

ONIONS: Onion Thrips: Apply 1 pt. per acre. Do not apply within 10 days of harvest.

PEAS: Aphids, Leaf Miners: Apply 1/2 pt. per acre. Beans may be picked immediately. Red vines and pea vine hay may be fed to dairy and beef cattle and sheep. Wait 4 days before using treatment if forage is to be cut for hay.

PEPPERS: Aphids, Serpentine Leaf Miners: Apply 1/2 pt. per acre. Do not apply within 5 days of harvest.

POTATOES: Aphids, Mealybugs, Colorado Potato Beetles, Dipterous Leaf Miners: Apply at the rate of 1/2 pt. per acre.

Banded Cucumber Beetles, Leafhoppers, Root-knot Aphids: Apply 1/2 pt. per acre. Do not apply within 25 days of harvest.

RADISHES, PARSNIPS, and TURNIPS: Aphids, Eye Beetles, Dipterous Leaf Miners: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

SPINACH and BEETS: Aphids, Dipterous Leaf Miners: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

SQUASH: Summer and winter squash: Aphids, Leafhoppers, Mites: Apply 1/2 pt. per acre. Serpentine Leaf Miners, Thrips: Apply 1/2 pt. per acre. Do not apply to summer squash within 7 days, or to winter squash within 3 days, of harvest.

SWEET CORN: Corn Earworms: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

TOMATOES: Aphids, Dipterous Leaf Miners, Colorado Potato Beetles, Banded Cucumber Beetles, Flea Beetles, Root-knot Aphids, Vine Weevils, Flea Beetles: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

RANGE, PASTURE, AND GRASSLAND INSECTS

BERMUDAGRASS and FORAGE GRASSES: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

ANGELICA PASTURE, HIGH BANKS, BROOMS, WHEAT, RYE, BARLEY, and OATS: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

ORNAMENTAL INSECT CONTROL: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

APPLICATIONS: Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

DO NOT USE: Do not use on certain ferns, Parsnips, Hibiscus, Papayas, Pilea, and Gardenia, since injury to the plants may occur.

MIMOSA WEBWORMS: Mix 1/2 fl. oz. in 3 gals. of water or 1 pt. in 100 gals. of water and apply as a cover spray when webworms first appear.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100