


NUSAN[®]

30 E.C.



ACTIVE INGREDIENT:

2-(thiocyanomethylthio) benzothiazole..... 30%

INERT INGREDIENTS..... 70%

Total 100%

Contains 2.635 pounds TCMTB per gallon.

* Trademark of Wilbur-Ellis Company

EPA Reg. No. 2935-389-ZA

EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN!

WARNING — AVISO

IF ON SKIN: Wash promptly and thoroughly with an abrasive soap and cool water and finally with glycerin. In case of persistent irritation of the skin, obtain medical attention. Remove and wash contaminated clothing before reuse.

IF IN EYES: Flush eyes immediately and thoroughly with cool water for 15 to 30 minutes. See a physician if any irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON SIDE PANEL

BEST AVAILABLE COPY

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

ENVIRONMENTAL HAZARDS

Change
to
Copy
"A"

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water or wetlands.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Change
to
Copy
"E"

Label treated seed. Seeds treated with NUSAN 30 E. C. cannot be used for food, feed, oil production or any other purpose except planting.

NUSAN is recommended for the control of soil and seedborne fungi that impair the germination and seedling development of cotton, wheat, barley, oats, rice, sorghum or milo, corn, sugar beets, and safflower. Nusan, when used as directed, will usually increase stands and yields of these crops by preventing seed decay and seedling diseases. Nusan is effective against Rhizoctonia, Pythium, Fusarium, Helminthosporium, Trileptia, Ustilago, Verticillium, and certain other fungi.

STORAGE AND DISPOSAL

1. Prohibitions: Do not contaminate water, food or feed by storage or disposal.
2. Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
3. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. Storage: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing, or pouring.

37504

BEST AVAILABLE COPY

DIRECTIONS (cont.)

NUSAN can be applied directly to the seed from the drum or as an agitated dispersion in water. The recommended rates on the crops presented below may be dispersed in as much as 16 fl. oz. of water and applied to 100 lbs. of seed.

Crop	Application Rates per 100 lbs. seed	Fungi and Diseases Controlled
Cotton *	Acid-delinted seed	Seed and seedling rots, damping-off, sore shin (Pythium, Rhizoctonia, Fusarium)
	1.25-4 fl. oz.	
	Machine-delinted seed	
	1.5-5 fl. oz.	
Wheat	1-1.25 fl. oz.	Scab, head blight (Fusarium, Helminthosporium); Bunt or stinking smut (Tilletia)

**WHEAT: FOR USE ONLY IN THE STATES OF KANSAS,
TEXAS, OKLAHOMA, CALIFORNIA, AND
ARIZONA.**

Barley	1-1.25 fl. oz.	Bunt or stinking smut (Tilletia); covered smut (Ustilago lolae); Barley stripe (Helminthosporium graminum)
Oats	1-1.25 fl. oz.	Loose smut (Ustilago avenae); covered smut (Ustilago lolae)
Rice	1-1.25 fl. oz.	Pythium; Helminthosporium oryzae; Water mold (Achlya)
Sorghum	1.25 fl. oz.	Covered head smut (Sporobolotheca)
Corn	1-1.5 fl. oz.	Helminthosporium, Pythium
Sugar Beet	1-2 fl. oz.	Seed, seedling rots (Phoma betae) and damping-off (Pythium and Rhizoctonia)
Safflower	1-2 fl. oz.	Safflower rust (Puccinia carthami)

* UNDER HEAVY PYTHIUM PRESSURE AND RATES
BELOW 5 FL. OZ. PER 100 POUNDS OF
SEED, WILBUR-ELLIS COMPANY RECOMMENDS
TREATMENT WITH WILBUR-ELLIS APRON TL
OR WILBUR-ELLIS APRON FLOWABLE FOR
ADDITIONAL SYSTEMIC PYTHIUM CONTROL.

SEED SHOULD BE PLANTED WITHIN 5 MONTHS
OF TREATMENT. IF STORED LONGER THAN
5 MONTHS, RE-TREATMENT WITH NUSAN 30 EC
MAY BE NECESSARY.

IMPORTANT

Do not use crops grown from Nusan-treated seed for grazing of livestock or to
harvest as silage or forage for seven weeks after planting. However, grain
harvested from plants grown from Nusan-treated seed can be used for food
or feed.

NOTICE: The statements made on this label are believed to be true and accurate, but
because of conditions of use which are beyond our control, WILBUR-ELLIS COM-
PANY does not make, nor does it authorize any agent or representative to make, any
warranty, guarantee or representation, expressed or implied, concerning this mate-
rial or the use thereof, except in conformity with the statements on the label. Neither
WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for
any personal injury or property damage or loss resulting to the buyer or to the other
person from handling, storage or use of this material, not in accordance with
directions. The buyer assumes all risk and liability resulting from improper han-
dling, storage or use and accepts and uses this material on these conditions.

In Case of Emergency:
Call CHEMTREC: (800) 424-9300

F-291

BEST AVAILABLE COPY

Copy A

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Do not contaminate water when disposing of equipment washwaters.

Copy B

Label treated seed: Seeds treated with NUSAN 30 E.C. cannot be used for food, feed, oil production or any other purpose except planting. Do not use crops grown from NUSAN 30 E.C. treated seed for grazing of livestock or to harvest as silage or forage for seven weeks after planting. However, grain harvested from plants grown from NUSAN 30 E.C. treated seed can be used for food or feed.

15-1-10