

SEP 16 1993

Ms. Joanne Pratt
Wilbur-Ellis Company
191 W. Shaw Ave., Suite 107
Fresno, CA 93704

Dear Ms. Pratt:

Subject: Seed Treatment Discoloration Statement
NUSAN 30 E.C.
EPA Reg. No. 2935-389
Your submission dated Aug. 5, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:9/16/93:305680₀

BEST AVAILABLE COPY

ACCEPTED
with COMMENTS
EPA Letter Dated:

SEP 16 1993

Under the Federal Insecticide
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

253,5-389

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye damage. May be harmful if absorbed through the skin. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

IMPORTANT!

Do not use crops grown from Nusan 30 E. C. treated seed for grazing of livestock or to harvest as silage or forage for seven weeks after planting. However, grain harvested from plants grown from Nusan 30 E. C. treated seed can be used for feed or feed.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

IMPORTANT. Label treated seed: Seeds treated with NUSAN 30 E. C. cannot be used for food, feed, oil production or any other purpose except planting. Do not use crops grown from NUSAN 30 E. C. treated seed for grazing of livestock or to harvest as silage or forage for seven weeks after planting. However, grain harvested from plants grown from NUSAN 30 E. C. treated seed can be used for food or feed.

NOTE: The purchaser of this product is responsible for insuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye added must be cleared for use under 40 CFR, Part 180.1001.

STORAGE AND DISPOSAL

1. Prohibitions: Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

3. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

4. Storage: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing, or pouring.

NUSAN is recommended for the control of soil and seedborne fungi that impair the germination and seedling development of cotton, wheat, barley, oats, rice, sugar beets, and safflower. Nusan, when used as directed, will usually increase stands and yields of these crops by preventing seed decay and seedling diseases. Nusan is effective against Rhizoctonia, Pythium, Fusarium, Helminthosporium, Tilletia, Ustilago, Verticillium, and certain other fungi.

NUSAN can be applied directly to the seed from the drum or as an agitated dispersion in water. The recommended rates on the crops presented below may be dispersed in as much as 1 1/2 fl. oz. of water and applied to 100 lbs. of seed.

In Case of Emergency,
Call CHEMTREC: (800) 424-9300

ACTIVE INGREDIENT:

2-(thiocyanomethylthio) benzothiazole 30%
INERT INGREDIENTS 70%

Total 100%

Contains xylene range aromatic solvent.
Contains 2.635 pounds TCMTB per gallon.

EPA Reg. No. 2935-389-ZA

EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN!

WARNING — AVISO

IF ON SKIN: Wash promptly and thoroughly with an abrasive soap and cool water and finally with glycerin. In case of persistent irritation of the skin, obtain medical attention. Remove and wash contaminated clothing before reuse.

IF IN EYES: Flood eyes immediately and thoroughly with cool water for 15 to 30 minutes. See a physician if any irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON SIDE PANEL

DIRECTIONS (cont.)

Crop	Application Rates per 100 lbs. seed	Fungi and Diseases Controlled
Cotton*	Acid-dolint seed 1.25-4 fl. oz.; Machine-dolint seed 1.5-5 fl. oz.	Seed and seedling rots, damping-off, scorchin (Pythium, Rhizoctonia, Fusarium)

For use only in the states of Kansas, Texas, Oklahoma, California and Arizona.

Wheat	1-1.25 fl. oz.	Scab, head blight (Fusarium, Helminthosporium); Bunt or stinking smut (Tilletia)
Barley	1-1.25 fl. oz.	Bunt or stinking smut (Tilletia); covered smut (Ustilago kollerii); Barley stripe (Helminthosporium graminum)
Oats	1-1.25 fl. oz.	Loose smut (Ustilago avenae); covered smut (Ustilago kollerii)
Rice	1-1.25 fl. oz.	Pythium; Helminthosporium oryzae; Water mold (achlya)
Sugar Beet	1-2 fl. oz.	Seed, seedling rots (Phoma betae) and damping-off (Pythium and Rhizoctonia)
Safflower	1-2 fl. oz.	Safflower rust (Puccinia carthami)

*Under heavy pythium pressure and rates below 4 fl. oz. per 100 pounds of seed, Wilbur-Ellis Company recommends treatment with Wilbur-Ellis Apron TL or Wilbur-Ellis Apron Flowable for additional systemic pythium control.

Seed should be planted within 5 months of treatment. If stored longer than 5 months, re-treatment with Nusan 30 E. C. may be necessary.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

Logo TM of WILBUR-ELLIS Company.

IDEAS TO GROW WITH and Nusan are registered trademarks of WILBUR-ELLIS Company.