# NUSAN 30 E.C.

2-(thiocyanomethylthio) benzothiazole 100%

BEST AVAILABLE COPY

Contains sylene range aromatic solvent. Contains 2.635 pounds TCMTB per gallon.

EPA Rea, No. 2935-389-ZA

ACTIVE INGREDIENT:

HIEVES indling

estock

, grain sed for

water

sams.

wash.

its la

PRE-TATE

DX Of

as ed 2005

age

ited

EPA Est. No. 2935-CA-1

### **KEEP OUT OF REACH OF CHILDREN!**

# **WARNING — AVISO**

IF ON SKIN: Wash promptly and thoroughly with an abrasive soap and cool water and finally with glycerin, in case of persistent initiation of the skin, obtain medical aftention. Remove and wash contaminated clothing before reuse.

IF IN EYES; Flood eyes immediately and thoroughly with cool water for 15 to 30 minutes. See a physician if any initiation persists.

IF SWALLOWED: Call a physician or Poison Control Center, Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of special. Do not induce vomiting or give anything by mouth to an unconscious person,

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON SIDE PANEL

**NET CONTENTS: 21/2 GALLONS** 



P.O. BOX 16458 · Fresno, CA 93755

Contract of the second

WILBU d١

gueren

poor q. posts. vigor. commi conditi

NI ·D × Crc Co. Sar (Cat "Un Wat Wil

546

mor

NL 9+

ca. ...

The st ofcon not me gyerei use (i WIL BL atty pe Per \$01 directi hendili Treatm or to hervest as allage or forage for seven weeks after planting. However, grain hervested from plants grown from Nusen 30 E. C. treated seed can be used for food or feed.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Do not contaminate water when disposing of equipment washwaters.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALWAYS READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRE-CAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Not for use on agricultural establishments in hopper-box, planter box, sturry box or other seed treatment applications at or immediately before planting.

Do not apply this product through any type of irrigation system.

IMPORTANT. Label treated seed: Seeds treated with NUSAN30 E.C. cannot be used for food, feed, oil production or any other purpose except planting. Do not use crops grown from NUSAN30 E.C. treated seed for grazing of livestock or to harvest as slage or forage for seven weeks after planting. However, grain harvested from plants grown from NUSAN30 E.C. treated seed can be used for food or feed.

NOTE: The purchaser of this product is responsible for insuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals, Refer to 21 CFR, Part 2.25. Any dye added must be cleared for use under 40 CFR, Part 180.1001.

## STORAGE AND DISPOSAL

- Prohibitions: Do not contaminate water, food or feed by storage or disposal.
- Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess
  pesticide, spray mixture or rinsate is a violation of Federal law. If these
  wastes cannot be disposed of by use according to label instructions, contact
  your State Pesticide or Environmental Control Agency, or the Hazardous
  Waste representative at the nearest EPA Regional Office for guidance.
- Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by oth x procedures approved by state and local authorities.
- Storeget Store in original container only and keep sealed, Store in closecstorage areas. Use caution when moving, opening, closing, or pouring.

51017

BEST AVAILABLE COPY

.....

#### DIRECTIONS (cont.)

NUSAN is recommended for the control of soil and seedborne fungi that impair the garmination and seeding development of cotton and safflower. Nusan, when used as directed, will usually increase stands and yields of these crops by preventing safed decay and seedling diseases, Nusan is effective against Phizoctonia, Pythium. Furtain turn and certain other fungi.

NUSAN can be applied directly to the seed from the drum or as an agricated dispersion in water. The recommended rates on the crops presented below may be dispersed in as much as 16 ft loz, of water and applied to 100 lbs, of seed.

Cotion\* Application flates per 100 lh.s. seed

Cotion\* Acid-delimed seed Seed and seedling rots, damping-off, sore-shiri (Pythium, Rhizoctonia, Fusarrum)

1.5-5 fl. oz

Satflower 1-2 fl. oz. Satflower rust (Puccinia cartham)

\*Under heavy pythium pressure and rates below 4 fl. oz. per 100 pounds of seed, Wilbur-Ellis Company recommends treatment with Wilbur-Ellis Apron TL or Wilbur-Ellis Apron Flowable for additional systemic pythium control.

(California only)

Seed should be planted within 5 months of treatment. If stored longer than 5 months, re-treatment with Nusan 30 E.C. may be necessary.

## NOTICE — READ CAREFULLY CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the lebel. Neither WILBUR-ELLIS COMPANY nor the selfer shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of WILBUR-ELLIS Company, no claims are made to guarantee germination of carry-over seed.

WILBUR-ELLIS Logo\*, IDEAS TO GROW WITH\* and Nusain\* are registered trademarks of WILBUR-ELLIS Company.

F-1094

In Case of Emergency, Call CHEMTREC: (800) 424-9300

BEST AVAILABLE COPY

....)......30% 70% Total 100%

IL No. 2935-CA-1

ion, Prmove and

ninistering syrup