

This product is labeled for use on the commodities grown on commercial properties by application in conventional hydraulic sprayers, or ground application by airplane, sprayer. When applying by ground equipment, use recommended amount of material with thorough coverage by ground equipment in 2 to 10 gallon of water. Do not apply when weather conditions favor drift or spray from areas treated. Repeat application may be necessary unless otherwise specified. Consult your state agricultural extension service for more information on application.

CROP	PESTS CONTROLLED	RATE	INTERVAL (DAYS) BETWEEN LAST APPLICATION AND HARVEST
FRUIT:			
Apples	Aphids, leafhoppers, mites (except rust mites)	1/2 to 1 1/2 pts./100 gals. water	28
Pears	Aphids	Ground Equipment: 1/2 to 1 1/2 pts./100 gals. water. Apply as an outside coverage spray. Aircraft Equipment: 3 to 6 pts./acre in 15 to 20 gals. water.	Do not apply during the bloom period
	Mites (except rust mites)	Ground Equipment: 1/2 to 1 1/2 pts./100 gals. water. Apply as a thorough distribution coverage spray.	15
Lemons	Scales (except black or snow)	Ground Equipment: 1/2 to 2 1/2 pts./100 gals. water. Apply as a thorough coverage spray.	45
Oranges	Thrips	Ground Equipment: 3/4 to 1 1/2 pts./100 gals. water. Apply as a mist spray. Aircraft Equipment: 3 to 6 pts./acre in 15 to 20 gals. water.	15
	Whiteflies	Ground Equipment: 1 1/2 pts./100 gals. water. Apply as a thorough distribution coverage spray.	15
Do not apply during bloom period. Do not use on citrus seedlings. Make no more than 2 applications to mature fruit. Do not enter treated groves within 4 days of last application.			
Citrus (non-bearing) (Calif. and Ariz.)	Thrips, Aphids	FOILAR SPRAY: 100 gals. Red-Top Dimethoate 2.67 Spray per 100 gallons of water. Repeat applications as necessary. Do not enter treated groves within 4 days of last application. SOIL DRENCH: (Trees 1-3 years old) 3 quarts Red-Top Dimethoate 2.67 Spray per acre in the furrow or basin around the base of the tree. Apply when insect injury to new growth appears. Red-Top Dimethoate 2.67 Spray should be used only on nursery stock and nonbearing trees. Do not apply in the year trees begin to bear a crop. Consult your state agricultural extension service or state agricultural extension service for proper timing of application.	

FIELD CROPS:			
Alfalfa	Aphids, mites, leafhoppers	Ground Equipment: 1/2 to 1 1/2 pts./acre	21
Do not apply to alfalfa in the bloom period. Do not apply within 28 days of harvest or pasturing. Make only one application per season. Effective only on cutting to which applied.			

VEGETABLE CROPS

Beans (green, snap, dry)	Aphids, leafhoppers, leaf miners, mites, lygus bugs	1 to 1 1/2 pts./acre	Beans may be harvested on day of application. Do not feed treated vines.
Broccoli	Aphids	1/2 to 1 1/2 pts./acre	7
Cauliflower	Aphids	1/2 to 1 1/2 pts./acre	3
Cabbage	Aphids, leafhoppers, leaf miners	3/4 pt./acre	7
Head lettuce	Aphids, leafhoppers, leaf miners	3/4 pt./acre	14
Leaf lettuce	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Spinach	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Collards	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Kale	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Turnip (greens and roots)	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Mustard greens	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Swiss-chard	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Endive (escarole)	Aphids, leafhoppers, leaf miners	3/4 pt./acre	
Melons (except watermelons)	Aphids, leafhoppers, leaf miners	1 1/2 pts./acre	3
Watermelons	Aphids, leafhoppers, leaf miners	3/4 to 1 1/2 pts./acre	3
Peas	Aphids	1/2 pt./acre	Peas may be harvested on day of application. Do not feed or graze hay within 21 days after last application when a stationary viner is used. Do not feed or graze when a mobile viner is used.
Peppers	Aphids, leaf miners, maggots	3/4 to 1 pt./acre	Peppers may be harvested on day of application.
Potatoes	Aphids, leaf miners, leafhoppers	3/4 to 1 1/2 pts./acre	Potatoes may be harvested on day of application.
Tomatoes	Aphids, leaf miners, leafhoppers	3/4 to 1 1/2 pts./acre	7

FIELD CROPS

Alfalfa (grown for hay)	Aphids, leafhoppers, lygus bugs, grasshoppers, reduction of alfalfa weevil larvae	1/2 to 1 1/2 pts./acre	Do not apply to alfalfa in the bloom period. Do not apply within 28 days of harvest or pasturing. Make only one application per season. Effective only on cutting to which applied.
-------------------------	---	------------------------	---

Cotton (grown in California and Arizona) Lygus b. leafhopper

Cotton (early season) Aphids, thrips
Cotton (Oklahoma, Texas) Fleahopper
Safflower (grown in California) Aphids, lygus b.

Tobacco Aphids

SEED CROPS:
Alfalfa Aphids, lygus b. hoppers of alfalfa larvae

FOREST INSECTS
TREE
Pinyon Pine Pinyon scale

Douglas Fir Fir cone

Pines Loblolly sawfly

NOTICE: The statements of use which are any agent or represent concerning this material WILBUR-ELLIS COMPANY or property damage or of this material, not in improper handling, etc.

ACCEPTED
JUN 13 1973
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION UNDER NO. 2935-390
TO ATTACHED COMMENTS.

