

2965-371

DIPEL® 150 DUST*

(Bacillus thuringiensis)

APPLICATION RATE:

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE and MUSTARD, and **TURNIP GREENS**: For the control of **IMPORTED CABBAGE WORM** apply 25 to 50 lbs. per acre.

GRAPES: For the control of **GRAPE LEAF FOLLIER** apply 25 to 50 lbs. per acre.

GRAPES (in California only): For the control of **CANNIVOROUS LEAF ROLLER** with single applications, as needed, apply 25 to 50 lbs. per acre. For the control of **CANNIVOROUS LEAF ROLLER** and **GRAPE LEAF ROLLER** with a scheduled program of 4 to 6 applications beginning with bud break, apply 12.5 to 25 lbs. per acre for each application.

TOBACCO and TOMATOES: For the control of **HORNWORMS** apply 12 to 25 lbs. per acre.

ACTIVE INGREDIENTS:

Bacillus Thuringiensis, Berliner, Potency of 320 International Units per mg. (at least 0.5 billion viable spores per g.)* 0.004%

INERT INGREDIENTS

99.936%

*Equivalent to potency of 0.15 billion International Units per pound of this product.

E. P. A. Reg. No. 2935-395-AA

CAUTION: KEEP OUT REACH OF CHILDREN!

DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

DIRECTIONS FOR USE

DIPEL 150 Dust is a biological insecticide intended for use against larvae of certain lepidopteran species.

DIPEL 150 Dust is indicated for use by a broadcast method and by spot treatments of plants.

DIPEL is the registered trademark of Abbott Laboratories, Chicago, Illinois 60064.