

2935-398

ACCEPTED

APR 23 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, or amended, for the pesticide registered under EPA Reg. No. 2935-398

DIRECTIONS FOR USE:

Apply recommended amounts when air is calm, using ground equipment, aircraft or hand duster. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Good underleaf coverage is essential since the insect must ingest BT 320 Dust. After eating BT 320 Dust, some insects stop feeding but may otherwise appear to be unaffected for several days. Use the higher recommendations against very heavy infestations, or where it is difficult to obtain good coverage due to rank foliage. BT 320 Dust is exempt from tolerance requirements, and may be applied in recommended amounts up to the day of harvest.

BT 320 DUST

(BACILLUS THURINGIENSIS)

ACTIVE INGREDIENTS:

Bacillus thuringiensis Berliner, potency of 320 International Units (at least 600 thousand viable spores) per milligram*.... 0.064%

INERT INGREDIENTS: 99.936%
100.000%

* Equivalent to potency of 145 Million International Units per pound.

EPA Reg. No. 2935-398-AA

CAUTION: KEEP OUT OF REACH OF CHILDREN!

RECOMMENDATIONS

Thorough coverage essential when using BT 320 Dust

Crop	Pest	Lbs. per Acre	Crop	Pest	Lbs. per Acre
BROCCOLI	Cabbage looper	20-40	KALE	Cabbage looper	20-40
	Imported cabbageworm	20-35		Imported cabbageworm	20-35
CABBAGE	Cabbage looper	20-40	LETTUCE	Cabbage looper	20-50
	Imported cabbageworm	20-35		MELONS	Cabbage looper
CAULIFLOWER	Cabbage looper	20-50	MUSTARD GREENS	Cabbage looper	20-40
	Imported cabbageworm	20-40		Imported cabbageworm	20-35
CELERY	Cabbage looper	20-50	POTATOES	Cabbage looper	20-40
COLLARDS	Cabbage looper	20-40		SPINACH	Cabbage looper
	Imported cabbageworm	20-35	TOMATOES		Cabbage looper
GRAPES	Grape Leaf Folder	20-50		Tomato Hornworm	20-35
GRAPES (California)	Omnivorous Leaf Roller	20-50	TURNIP GREENS	Imported cabbageworm	20-35

Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply by ground equipment to obtain thorough coverage of leaf surfaces. For the control of OMNIVOROUS LEAF ROLLER and GRAPE LEAF FOLDER with a scheduled program of 4 to 6 applications apply 12.5 to 25 lbs. per acre for each application.

STORE IN COOL PLACE

CONTAINER DISPOSAL: DO NOT REUSE EMPTY CONTAINERS. DISPOSE OF IN ACCORDANCE WITH STATE AND LOCAL REGULATIONS.

2935-398

04/23/1975