

# DIBROM<sup>®</sup> 4 DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

## RECOMMENDATIONS FOR USE:

Use 40 to 50 pounds per acre unless otherwise specified. This product is formulated for application by ground or air equipment. For Airplane Application, use heavier dosages. Begin applications at first sign of insects. Repeat as necessary. Apply for contact and thorough coverage to obtain control of insects.

**BEANS (Dry or Field):** APHIDS, MITES, LYGUS BUGS, THRIPS, LEAFHOPPERS, LEAF MINERS, LOOPERS. Do not apply within 4 days of harvest.

**BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, BRUSSELS SPROUTS, KALE:** CABBAGE LOOPER, IMPORTED CABBAGEWORM, DIAMOND-BACK CATERPILLAR, APHIDS. Do not use within 1 day of harvest on Broccoli, Cabbage, Cauliflower, and Brussels Sprouts. Do not apply to Collards and Kale within 4 days of harvest.

**CITRUS (Oranges, Lemons, Grapefruit, Tangerines):** APHIDS, SPIDER MITES, CITRUS CUTWORMS, THRIPS. Do not apply within 1 day of harvest.

**COTTON:** LYGUS, COTTON LEAF PERFORATOR, ARMYWORMS, LOOPERS, FLEAHOPPERS — 35 lbs. per acre. Apply at 7 to 10-day intervals while pests persist. Do not apply within 4 days of hand-picking.

**CUCUMBERS, SUMMER SQUASH:** CUCUMBER BEETLES, APHIDS, LEAF MINERS, PICKLEWORMS, ARMYWORMS, LOOPERS, LEAFHOPPERS. May be used up to harvest.

**EGGPLANT:** APHIDS, FLEA BEETLES, LEAF MINERS — 25 lbs. per acre. Do not use within 1 day of harvest.

**GRAPES:** LEAFHOPPERS — 40 to 50 lbs. per acre. THRIPS — 20 to 25 lbs. per acre. DROSOPHILA — 40 lbs. per acre. Apply pre-harvest at first sign of Drosophila or bunch rot. Repeat at weekly intervals as necessary to maintain control. Do not apply within 4 days of harvest. Do not apply on Italia varieties of grapes as fruit injury may occur.

**HOPS:** HOP LOOPER — 25 lbs. per acre. Do not apply within 4 days of harvest.

**LETTUCE, SPINACH, CELERY, CHARD, TURNIP GREENS:** LOOPERS, APHIDS — 40 to 50 lbs. per acre. May be used 1 day before harvest on Lettuce, Spinach, Turnip Greens and Chard. Do not apply to Celery within 4 days of harvest.

**MELONS (Cantaloupes, Honeydews, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH:** CABBAGE LOOPERS, ARMYWORMS, LEAFHOPPERS, APHIDS, LYGUS BUGS — Do not use within 1 day of harvest.

**ORNAMENTALS:** APHIDS, MITES, ADULT WHITE FLIES — 25 lbs. per acre.

**PEACHES:** APHIDS, FRUIT FLIES (Drosophila), RED SPIDER MITES, PEACH TWIG BORER, LYGUS BUGS, ORIENTAL FRUIT MOTH, THRIPS, STINK BUGS — 50 lbs. per acre. Do not apply within 4 days of harvest. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

F-1174 EPA Est. 2935-CA-1

© DIBROM is Chevron Chemical Company's Reg. TM

## ACTIVE INGREDIENT:

\* Naled

## INERT INGREDIENTS

\* 1,2-dibromo-2,2-dichloroethyl dimethyl phosphate

U. S. Pat. No. 2,971,882

Use within \_\_\_\_\_ days of manufacture only, as this product decomposes.

Date of Manufacture \_\_\_\_\_

EPA Reg. No. 2935-400-AA

## CAUTION: KEEP OUT OF REACH OF CHILDREN!

**CAUTION:** Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Do not store or transport near feed or food.

**Note to Physician:** Atropine is antidotal.

Do not use on cherries as injury may result. May corrode metal dust equipment, clean equipment after use.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

## RECOMMENDATIONS FOR USE: (Continued)

**PEPPERS:** APHIDS, FLEA BEETLES, LEAF MINERS — 25 lbs. per acre. Do not use within 1 day of harvest.

**ROSES (Field Grown):** APHIDS, MITES, TENT CATERPILLARS, THRIPS, LEAF ROLLERS — 25 lbs. per acre.

**STRAWBERRIES:** SPIDER MITES, APHIDS — Do not apply within 4 days of harvest.

**TOMATOES (Field):** FRUIT FLIES (Drosophila sp.), TWO-SPOTTED MITES, APHIDS, LEAFMINERS, TOMATO FRUITWORMS, HORNWORMS — 25 lbs. per acre. Make first application at 5 to 7 days before first picking and continue at 5 to 7-day intervals until harvest is complete. Do not use within 1 day of harvest.

**WALNUTS:** SPIDER MITES, APHIDS — 50 to 80 pounds per acre. Repeat as necessary. Do not apply within 10 days of harvest.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL DRY PLACE, OUT OF DIRECT SUNLIGHT.

Empty bags should be buried or burned. Stay away from smoke. If the bags are burned, the ashes should be buried in a safe place. IN CALIF.: Dispose of in accordance with state and local regulations.

2935-400