

# red-top

## BAITOX 5 BAIT

CONTAINS 5% DYLOX®

ACCEPTED

OCT 8 1975

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, this pesticide  
registered under  
EPA Reg. No. 2935-401

5.0%

95.0%

100.0%

### RECOMMENDATIONS FOR USE:

Baitox Bait can be applied by ground or air equipment.

Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected to be infested.

### DIRECTIONS FOR USE:

**ALFALFA:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 14 days of harvest. Apply only once between cuttings.

**BARLEY, OATS, FLAX, WHEAT:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 21 days of harvest. Do not graze treated fields. Do not feed treated straw to dairy or meat animals.

**BEANS (Dry, Snap):** Apply 30 lbs. per acre to control ARMYWORMS, CUTWORMS (Surface Feeding), CRICKETS, DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

**BEETS:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 28 days of harvest. Do not eat tops.

**BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 21 days of harvest.

**COLLARDS:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), SALTMARSH CATERPILLARS. Do not apply within 28 days of harvest.

**CORN:** Apply 20 lbs. per acre to control ARMYWORMS, CUTWORMS (Surface Feeding), DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Apply once per season when plants are 3 to 12 inches high. Do not use corn for food, or ensilage within 40 days of treatment.

**COTTON:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application.

**LETTUCE:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), SALTMARSH CATERPILLARS and DARKLING GROUND BEETLES. Do not apply within 28 days of harvest. Do not apply after heads start to form.

### ACTIVE INGREDIENT:

0,0-Dimethyl (2,2,2-Trichloro-1, hydroxyethyl) phosphonate\*

### INERT INGREDIENTS

DYLOX is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

EPA Reg. No. 2935-401-AA

## CAUTION: KEEP OUT OF THE REACH OF CHILDREN!

May be harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. If use results in poisoning, obtain prompt medical aid.

### PHYSICIAN'S NOTE: ATROPINE SULFATE IS ANTIDOTAL.

This product is toxic to birds and other wildlife. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Rinse equipment and container and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

### DIRECTIONS FOR USE (Continued):

**SUGAR BEETS:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Do not apply within 14 days of harvest. Do not feed tops which have been treated within 28 days of harvest to livestock.

**TOMATOES:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS,

and SALTMARSH CATERPILLARS. Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

**BEETS:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 28 days of harvest. Do not eat tops.

**BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 21 days of harvest.

**COLLARDS:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), SALTMARSH CATERPILLARS. Do not apply within 28 days of harvest.

**CORN:** Apply 20 lbs. per acre to control ARMYWORMS, CUTWORMS (Surface Feeding), DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Apply once per season when plants are 3 to 12 inches high. Do not use corn for food, or ensilage within 40 days of treatment.

**COTTON:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application.

**LETTUCE:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), SALTMARSH CATERPILLARS and DARKLING GROUND BEETLES. Do not apply within 28 days of harvest. Do not apply after heads start to form.

**PEPPERS:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Do not apply within 21 days of harvest.

**SAFFLOWER:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and SALTMARSH CATERPILLARS. Do not apply after plant begins to bloom. Do not apply more than twice per year.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

soap and hot water before re-use. Do not contaminate food or feed. If use results in poisoning, obtain prompt medical aid.

**PHYSICIAN'S NOTE: ATROPINE SULFATE IS ANTIDOTAL.**

This product is toxic to birds and other wildlife. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Rinse equipment and container and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

**DIRECTIONS FOR USE (Continued):**

**SUGAR BEETS:** Apply 30 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), and DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Do not apply within 14 days of harvest. Do not feed tops which have been treated within 28 days of harvest to livestock.

**TOMATOES:** Apply 20 lbs. per acre to control ARMYWORMS, CRICKETS, CUTWORMS (Surface Feeding), DARKLING GROUND BEETLES, and SALTMARSH CATERPILLARS. Do not apply within 28 days of harvest. Do not graze treated fields.

**FLOWERS:** (Including - Asters, Gladiolus, Iris, Mums, Stock and Daisies) Apply 20 to 30 lbs. per acre to control ARMYWORMS, CRICKETS and SALTMARSH CATERPILLARS. Caution - May cause injury on certain varieties of carnations and zinnias.

EPA Est. 2935-CA-1 F-675



# WILBUR-ELLIS COMPANY



FRESNO • LOS ANGELES • GLENDALE, ARIZ. • PORTLAND • SEATTLE