


BOTRAN® 6 DUST



RECOMMENDATIONS FOR USE:

APRICOT: Use 50 lbs. per acre to control BLOSSOM BLIGHT **Monilinia** and FRUIT DECAY (Brown Rot **Monilinia** and **Rhizopus** Rot). For BLOSSOM BLIGHT, apply at pink bud and full bloom. For FRUIT DECAY, apply at 18, 10 and 1 day before harvest. Do not apply within 1 day of harvest.

GRAPES: Use 30 lbs. per acre to control **Botrytis** ROT (Bunch Rot and Storage Rot). For best results, apply Botran 75W spray soon after Shatter (See Botran 75W label for directions). Follow after cluster tightening with Botran 6% Dust application and repeat at two-week intervals. Allow 1 day before application and harvest. If dust only is used, begin application while clusters are still loose and repeat application at two-week intervals. Allow 1 day between application and harvest.

TTUCE, HEAD: Use 30 lbs. per acre to control WILT (**Botrytis** Rot and **Sclerotinia**). Apply 1st application at thinning. Repeat applications 7 days later. Do not apply within 14 days of harvest.

ONION and GARLIC: Use 250 to 400 lbs. per acre to control WHITE ROT (**Sclerotium**). For direct seeding, mix (broadcast and incorporate) Botran in top 1½ inches of soil 1 to 2 weeks before onions are seeded. **Planting sets or buds** - Apply as a broadcast application and mix into the top 1½ inches of soil; or use 250 to 400 lbs. per 43,560 feet of row as an in-furrow application to soil around sets or buds. When lettuce is to be planted as a follow up crop, plow soil to 8 inches and cross disc the area prior to planting.

SWEET CHERRY: Use 50 lbs. per acre to control BLOSSOM BLIGHT **Monilinia** and FRUIT DECAY (Brown Rot **Monilinia** and **Rhizopus** Rot). For BLOSSOM BLIGHT, apply at popcorn, bloom, full bloom and petal fall. For FRUIT DECAY, apply 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest.

ACTIVE INGREDIENT:

2,6-Dichloro-4-nitroaniline 6%

INERT INGREDIENTS 94%

100%

EPA Reg. No. 2935-402-AA

WARNING: KEEP OUT OF REACH OF CHILDREN!

Harmful if swallowed. Avoid inhalation of dust. In case of contact, remove contaminated clothing and wash skin with soap and water.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

PRECAUTION: Do not use on germinated seeds or annual seedlings except as suggested. Do not use on dried fruits. Do not plant Spinach as a followup crop in treated soil.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL. DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

Do not reuse container. Destroy when empty.

RECOMMENDATIONS FOR USE: (Continued)

PEACHES and NECTARINE: Use 60 lbs. per acre to control BLOSSOM BLIGHT **Monilinia** and FRUIT DECAY (Brown Rot **Monilinia** and **Rhizopus** Rot). For BLOSSOM BLIGHT, apply at pink bud and full bloom. For FRUIT DECAY, apply 1 day prior to harvest. Do not apply within 1 day of harvest.