

BOTRAN® 6 SULFUR 25 DUST

RECOMMENDATIONS FOR USE:

This product is formulated for application by ground or air.

GRAPES: Use 30 lbs. per acre to control BOTRYTIS ROT (Bunch Rot and Storage Rot). Follow cluster tightening with an application and repeat at two-week intervals. If dust only is used, begin application while clusters are still loose and repeat at two-week intervals. Do not apply within 1 day of harvest.

PEACHES and NECTARINES: Use 60 lbs. per acre to control BLOSSOM BLIGHT *Monilia* and FRUIT DECAY (Brown Rot *Monilia* and *Rhizopus* Rot). For BLOSSOM BLIGHT, apply at pink bud and full bloom. For FRUIT DECAY, apply ~~at 18 days, 10 days and 1 day prior to harvest.~~ at 18 days, 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest.

SWIFT CHERRY: Use 50 lbs. per acre to control BLOSSOM BLIGHT *Monilia* and FRUIT DECAY (Brown Rot *Monilia* and *Rhizopus* Rot). For BLOSSOM BLIGHT, apply at popcorn, bloom, full bloom and petal fall. For FRUIT DECAY, apply 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest.

DO NOT REUSE EMPTY CONTAINERS. DESTROY WHEN EMPTY.

DISPOSE OF IN ACCORDANCE WITH STATE AND LOCAL REGULATIONS.

© Reg. T.M. Tuco Products Co., Division of the Upjohn Co.

EPA Est. 2935-CA-1 F-775

ACTIVE INGREDIENTS:

2,6-Dichloro-4-nitroaniline 6%

Sulfur 25%

INERT INGREDIENTS 69%

TOTAL 100%

EPA Reg. No. 2935-403-AA

WARNING: KEEP OUT OF REACH OF CHILDREN!

Harmful if swallowed. Avoid inhalation of dust. In case of contact, remove contaminated clothing and wash skin with soap and water. Avoid contact with skin and eyes. In case of contact with eyes, flush with water and call a physician.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

PRECAUTION: Do not use on germinated seeds or annual seedlings except as suggested. Do not use on dried fruits. Do not plant Spinach as a followup crop in treated soil.

Do not apply during or before periods of high temperatures. (90° F. or higher)

Do not allow workers to enter vineyards or peach and nectarine orchards where foliage contact will occur until one day after application.

ACCEPTED

9 JUN 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED