


PM 19 2935-405

File 134

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

(A) 	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration	OPP Identifier Number 201611
		<input checked="" type="checkbox"/> Amendment	
Application for Pesticide:		<input type="checkbox"/> Other	

Section I

1. Company/Product Number 2935-405	2. EPA Product Manager D. Edwards	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Supreme Oil	Pl.# 19	
5. Name and Address of Applicant (Include ZIP Code) WILBUR-ELLIS Company 191 W. Shaw Avenue Suite 107 Fresno, CA 93704 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Change to labeling statement prohibiting application to water. The exact wording from PR Notice 93-3 has been used.


NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 93-3

DATE: 8/3/93

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted.				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container	If "Yes," Unit Package wgt.	No. per container	If "Yes," Package wgt.	Other (Specify) _____	
4. Size(s) of Retail Container 5, 55, 110 Gallon	5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product		6. Manner in Which Label is Affixed To Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other (_____)		

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Joanne Pratt	Title Product Registration and Labeling Specialist	Telephone No. (Include Area Code) (209) 226-1934
2. Signature 		6. Date Application Received (Stamped)
3. Title Product Registration and Labeling Specialist		
4. Typed Name Joanne Pratt		
5. Date July 8, 1993		

NOT REGISTERED IN CALIFORNIA
SPECIMEN LABEL

NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 88-8
DATE 8/3/93

INSECTICIDES

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: Harmful if Swallowed. Do Not Induce Vomiting. Call A Physician Immediately! Avoid breathing of spray mist.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Copy A

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS: To avoid injury to trees, fruit or foliage the following precautions should be taken when using emulsifiable spray oils: Do not combine SUPREME SPRAY OIL with sulfur sprays on foliage or apply following a sulfur spray on foliage. Avoid applications of sulfur following sprays of SUPREME SPRAY OIL until sufficient time has elapsed for oil dissipation. For foliage applications, the spray tank should be free of sulfur residues. After bloom do not combine or follow an oil application with Captan, Sevin, Kelthane or Karathane as foliage or fruit injury may result. When through spraying with oil, clean tanks and system of any residues in that oil residues may cause crop injury when combined with some spray materials. On oil sprays both dormant and summer followed or preceded by sprays of other materials that may not be compatible with oils, sufficient interval needs to be maintained as there is the potential for serious leaf and fruit bud injury and a reduction in fruit set.

Do not use if emulsion breaks. If the emulsion should break in the spray tank, free oil will float to the surface causing excessive oil deposits at the end of each tank. Severe injury to trees may result if used under these conditions.

When cool slow drying conditions exist, labeled rates may cause greater potential for delayed bud development and or leaf and fruit bud damage which may result in reduced fruit set. Use lower labeled rates or delay application until weather improves. Do not apply when temperatures are likely to fall below freezing before the spray has had a chance to dry. Do not apply SUPREME SPRAY OIL when temperature is below 40°F or above 85°F. Avoid use in sprayers having hydraulic agitation or those having insufficient agitation to form an emulsion with SUPREME SPRAY OIL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing, areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment (see above) in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Caution. Area treated with Spray Oil on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here action to take in case of accidental exposure — see above.)"

Do not apply this product through any type of irrigation system.

If spray solution containing SUPREME SPRAY OIL has set without agitation for more than five minutes, be sure to flush all spray hoses back into spray tank and agitate for five minutes before resuming spray application.

MIXING DIRECTIONS: With agitator running, start filling spray tank with water. When combining with wettable powder or fungicide products, add these first when tank is 1/2 filled. Add SUPREME SPRAY OIL when tank is about 1/2 full and while there is sufficient agitation of the spray to complete emulsification. Be sure whether spraying by air or ground, dilute or concentrate, that agitation is adequate to keep the spray mixed uniformly.

AERIAL APPLICATION: When applying SUPREME SPRAY OIL with aerial equipment, apply 2 to 5 gallons per acre. On young trees or oil susceptible varieties, use the lower rates. Use sufficient water and total gallonage per acre to give thorough coverage. Consult your local Ag Chem authority for approved combination of SUPREME SPRAY OIL with other materials.

APPLES: Dormant to delayed timing for the control of SAN JOSE SCALE, APHIDS, EUROPEAN RED and BROWN MITE EGGS and GRAPE MEALYBUG.

39440

Wilbur-Ellis
SUPREME OIL
MITICIDE — INSECTICIDE

ACTIVE INGREDIENT:	By Wt.
Mineral Oil	99.0%
INERT INGREDIENT:	1.0%
	TOTAL 100.0%
Minimum Unsulfonated Residue (ASTM D-483)	93.0%
EPA Reg. No. 2935-405	EPA Est. No. 2935-WA-1

KEEP OUT OF REACH OF CHILDREN!

CAUTION:

— SEE PRECAUTIONARY STATEMENTS IN LEFT PANEL —

DIRECTIONS FOR USE (cont.)

Dilute application (200 to 400 gallons per acre): Use 1 to 2 gallons of SUPREME SPRAY OIL per 100 gallons of water. Be sure to maintain thorough coverage for adequate pest control. Do not exceed 8 gallons of SUPREME SPRAY OIL per acre. Consult your local agricultural consultant for specific area rates.

Concentrate application (25 to 125 gallons per acre): Use 2 to 5 gallons of SUPREME SPRAY OIL per acre. On young trees and on oil-susceptible varieties use the lower rate. Consult your local agricultural consultant for specific area rates.

Summer application timing for the control of ORCHARD MITES and SAN JOSE SCALE.

Dilute application (200 to 400 gallons per acre): Use 1 to 4 quarts of SUPREME SPRAY OIL per 100 gallons of water as a dilute spray.

Concentrate application (25 to 125 gallons per acre): Use 1 to 3 gallons of SUPREME SPRAY OIL per acre. On young trees and susceptible varieties use lower rates. Special precautions should be used in summer oil applications. Summer oil applications have greater potential for plant and fruit damage than pre-bloom applications.

PEARS: Dormant to delayed dormant timing for the control of SAN JOSE SCALE, APHIDS, EUROPEAN RED and BROWN MITE EGGS, GRAPE MEALYBUG, a synergist for PEAR PSYLLA insecticides and delay of pear psylla egg deposition. If two applications are used pre-bloom at the dormant and delayed dormant timings, then reduce oil rates to the lower labeled amounts.

Dilute application (200 to 400 gallons per acre): Use 1 to 2 gallons of SUPREME SPRAY OIL per 100 gallons of water. Be sure to maintain thorough coverage for adequate pest control.

NET CONTENTS:

Wilbur-Ellis

P.O. BOX 16458 • FRESNO, CA 93755

(continued on next page)

SUPREME OIL (continued)

DIRECTIONS FOR USE (cont.)

STORAGE AND DISPOSAL

1. **Prohibitions:** Do not contaminate water, food or feed by storage or disposal.
2. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
3. **Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. **Storage:** Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

Concentrate application (25 to 125 gallons per acre): Use 2 to 5 gallons of SUPREME SPRAY OIL per acre. On young trees and oil-susceptible varieties use lower rates.

Summer application for the control of ORCHARD MITES, SAN JOSE SCALE and as a synergist for PEAR PSYLLA insecticides.

Dilute application (200 to 400 gallons per acre): Use 1 to 4 quarts SUPREME SPRAY OIL per 100 gallons of water as a dilute spray.

Concentrate application (25 to 125 gallons per acre): Use 1 to 3 gallons of SUPREME SPRAY OIL per acre. On young trees and susceptible varieties use lower rates. Special precautions should be used in summer oil applications. Summer oil applications have greater potential for plant and fruit damage than pre-bloom applications.

At higher elevations and cooler pear growing districts use lower end of labeled rates.

APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES: Dormant to delayed dormant timing for the control of SAN JOSE SCALE, APHIDS, EUROPEAN RED and BROWN MITE EGGS.

Dilute application (200 to 400 gallons per acre): Use 1 to 2 gallons of SUPREME SPRAY OIL per 100 gallons of water as a dilute spray.

Concentrate application (25 to 125 gallons per acre): Use 3 to 5 gallons of SUPREME SPRAY OIL per acre. On young trees and oil-susceptible varieties use the lower rate.

GRAPES: For the control of MEALY BUGS, apply at the rate of 1 gallon per 100 gallons of water. Use 300 gallons of this mixture per acre with thorough coverage of the vines at delayed dormant stage up to bud break. For concentrate applications, use 3 gallons per acre in 40 to 100 gallons of water.

ORNAMENTALS and SHADE TREES (Conifers — See warning, Holly, Pyracantha, Sycamores, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch): Soft Brown Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, San Jose Scale, Red Spider Mites and Eggs — 1 gallon to 100 gallons water. Dormant spray on deciduous varieties — 2 gallons per 100 gallons water. Dormant spray on other varieties — 1 gallon per 100 gallons of water. WARNING: May cause discoloration of blue spruce. Do not use on conifers or ferns not tolerant to oil sprays. Rates less than 1 gallon per 100 gallons of water can result in poor pest control.

WALNUTS (Late Spring to Mid-Summer only): Summer stages of Frosted Scale, European Fruit Scale, San Jose and other Soft-Shell Scale, Red Spider Mites — 2 to 4 quarts. The addition of an approved phosphate insecticide aids in scale control. See label for recommended dosages. Do not apply after husk split.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

KARATHANE — Reg. TM of Rohm & Haas Co
KELTHANE — Reg. TM of Rohm & Haas Co
SEVIN — Reg. TM of Rhone-Poulenc

F 991

Copy A

"For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark."

11 2
11 2
11 2
11 2
11 2