

PM-13

Reg # 2935-408

1/10/94

File

197



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 10 1994

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Joanne Pratt
WILBUR ELLIS CO.
191 W. Shaw Avenue
Fresno, CA 93704

Subject: Label Amendment Submission of 08/30/93 in Response to PR Notice 93-7
EPA Reg. No. 2935-408
DIAZINON 14G

IMPORTANT NOTICE: This Letter Requires You to Submit a Notification to the Office of Compliance Monitoring within 20 Days.

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

SERIOUS ERROR(S) ON SUBMITTED LABELING:

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons, or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING. Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

Each notification must include the following information:

1. Registrant's name and address.
2. EPA Registration number of the product for which the labeling was submitted.
3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

1. Quantity of the product that has been sold or distributed.
2. EPA Producing Establishment number.
3. Location of the product, if known.
4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director
Compliance Division, (7203)
Worker Protection Notification
Office of Compliance Monitoring
U.S. Environmental Protection Agency
401 M Street SW
Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling as amended

AND

- WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
 Office of Pesticide Programs (7505C)
 U.S. Environmental Protection Agency
 401 M Street, SW
 Washington, D.C. 20460-0001

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 10 1994

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2935-408



DIAZINON 14G

ORGANO PHOSPHATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before smoking and eating. May be absorbed through skin.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unexposed persons.

Copy B - Add

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Copy D - Add

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY TREATMENT

Do not enter treated areas without protective clothing until dusts have settled. Because certain states may require more restrictive re-entry intervals for various uses, consult with this product on your State Department of Agriculture for further information.

Written oral warnings must be given to workers who are expected to be in treated areas or in areas about to be treated with this product. Oral warnings should specify protective clothing, areas of fields that cannot be treated without this clothing, period of time area is to be vacated, and statement of practical treatment in case of exposure. When oral warnings are given, warnings shall be given in a language commonly understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Caution: Area treated with Diazinon (date of application). Do not enter without appropriate protective clothing until dusts have settled. (Insert here action to take in case of accidental exposure)."

STORAGE AND DISPOSAL

- Prohibitions: Do not contaminate water, food or feed by storage or disposal.
- Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.
- Storage: Store in original container and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

ONLY FOR SALE TO, USE AND STORAGE BY AGRICULTURAL AND SERVICE PERSONS

Use within _____ days of manufacture only, as this product decomposes.

Date of Manufacture _____

Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, harrow, or other suitable means for greatest effectiveness.

RECOMMENDATIONS FOR USE

SMALL FRUIT INSECTS

STRAWBERRIES; MOLE CRICKETS — Apply 7 lbs. per acre when insects are present. Make applications to plant beds or fields 1-2 days prior to transplanting and work into soil 1-2 inches.

FIELD AND FORAGE INSECTS

BEANS (Pinto); LESSER CORNSTALK BORERS — Apply 7-14 lbs. per acre in a 6-inch band centered over the row when beans are 1/2- to 1-inch high.

CORN (Not for use in California)

Corn Rootworm Larvae; (Postemergence Basal Treatment) — Apply 4 1/2-8 1/2 oz. per 1,000 linear feet of row in a band over the row at first sign of rootworms, but no later than June 24 in the Middle Corn Belt or July 1 in the Northern Corn Belt. Cover treated band with 2-3 inches of soil.

Apply amount per acre recommended according to row spacing chart for basal treatment. **Lesser Cornstalk Borers:** Apply 3 1/2-17 oz. per 1,000 linear feet of row in a 10-inch band centered over the row. Make application at planting time or when crop begins to emerge. Cover the treated band with 1-2 inches of soil.

Apply amount per acre recommended according to row spacing chart for basal treatment.

46063

ACTIVE INGREDIENTS:

O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%

INERT INGREDIENTS: 85.7%

TOTAL: 100.0%

EPA Reg. No. 2935-408-AA

EPA Est. No. 33691-AZ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION:

— SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONS —

RECOMMENDATIONS FOR USE (cont.)

Suggested Rates According to Corn Row Spacing

Row Spacing	Linear feet of corn row per acre	Amount of Wilbur-Ellis Diazinon 14G per acre Postemergence basal treatment
48"	10,800	3 - 6 lbs.
42"	12,455	3 1/2 - 6 1/2 lbs.
40"	13,658	3 1/2 - 7 lbs.
38"	13,939	3 1/2 - 7 1/2 lbs.
36"	14,520	4 - 8 lbs.
30"	17,424	4 1/2 - 9 1/2 lbs.
24"	21,780	5 1/2 - 11 1/2 lbs.
20"	26,136	7 - 14 lbs.
15"	34,848	9 1/2 - 18 1/2 lbs.

Note: For row spacings not listed, calculate the appropriate dosage rates based on the 40-in. row figures.

European Corn Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls, when 75% of plants show first-generation borers feeding. Make a second application 7 days later if needed. To control second-generation corn borers, apply 7-14 lbs. per acre, as above, when field count shows 100 egg masses per 100 plants.

WILBUR-ELLIS Diazinon 14G may be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above.

Fall Armyworms: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when armyworms first appear. Repeat application if necessary.

Southwestern Corn Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when second-generation borers first appear. Repeat application one week later.

Note: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be picked immediately. Temporary spotting of foliage sometimes occurs following application.

Cutworms: Broadcast 14-28 lbs. per acre. For surface cutworms apply just prior to planting of when larvae first appear and work into soil 1-2 inches. Do not apply after tassel. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches.

Seed Corn Maggots: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2 inches.

Garden Symphylans (Centipedes): Broadcast 70 lbs. per acre just prior to planting and work into soil 4-8 inches. This preplant treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

Wireworms: Broadcast 21-28 lbs. per acre before planting. Work into soil 4-6 inches.

FORAGE CROPS (Alfalfa, Clover, Cowpeas and Leanedaze): CUTWORMS — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

SORGHUM: CUTWORMS — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORN BORERS — Apply 7-14 lbs. per acre in a 10-inch band over the row at planting time or as crop emerges. Cover the band with 1-2 inches of soil.

Note: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOYBEANS; CUTWORMS — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORNSTALK BORERS — Apply 7-14 lbs. per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

SUGAR BEETS; SUGAR BEET ROOT MAGGOT — Apply 7-14 lbs. per acre in a 5-7 inch band (based on row width of 22 ins.) or 4-7-8 oz. per 1,000 linear feet of row in a 5-7 inch band either incorporating 2-3 inches at planting or applying above the seed after the seed is covered. Apply ahead of the press wheel. If heavy maggot infestation is anticipated or a wet spring occurs, a second application may be made, early postemergence, when the adult maggots (flies) first start to appear. If a second application is necessary, apply 7-10 1/2 lbs. per acre in a 5-7 inch band (based on row width of 22 ins.) or 4-7-7, 1 oz. per 1,000 linear feet of row in a 5-7 inch band directly over the row.

Note: Diazinon should not come in direct contact with
WIREWORMS — Broadcast 21-28 lbs. per acre.

VEGETABLES

BROCCOLI, BRUSSELS SPROUTS, CUCUMBERS: Broadcast 14-21 lbs. per acre before planting.

RADISHES; ROOT MAGGOTS: Apply 10 lbs. per acre in the seed furrow at planting time.

ONIONS; ONION MAGGOTS — Apply 1 lb. per acre prior to planting. Thoroughly mix the soil before planting.

POTATOES; VINEGAR FLIES (Drosophila spp.): Repeat in 7-10 day intervals or as needed.

OTHER VEGETABLES

CUTWORMS (Surface and Subterranean Cutworms, Cabbage Cutworm, Celery, Collards, Cucumber, Endive, Kale, Lettuce, Mustard, Peppers, Squash, Swiss Chard, Tomatoes, Watercress) are just prior to planting. Work into soil 2-4 inches to control subterranean cutworms.

POTATOES — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 4 inches.

MOLE CRICKETS; BROCCOLI, BRUSSELS SPROUTS, CUCUMBERS, ENDIVE, KALE, LETTUCE, MUSTARD, PEPPERS are when insects are present. Make application of seed or transplanting and work into soil 2-4 inches.

LESSER CORNSTALK BORERS; BEAN BORERS: 6-inch band centered over the row. Apply 7-14 lbs. per acre.

WIREWORMS: Broccoli, Brussels Sprouts, Celery, Collards, Cucumbers, Endive (or Onions, Parsley, Peas, Peppers, Pole Beans, Potatoes, Swiss Chard, Tomatoes, Watercress) are just prior to planting. Work into soil 4 inches.

POTATOES — Broadcast at 21-42 lbs. per acre just prior to planting, placing 1 lb. per 100 sq. ft. of row.

GARDEN SYMPHYLANS (CENTIPEDE) Radishes, Red Beets, Snap Beans, Tomatoes, Watercress, (see also subterranean), mole crickets, etc.

LAWN INSECTS (Not for use in California)

Keep children and pets off treated areas until dry.

Lawn Chinch Bugs: Apply 1/2-1 lb. per 1,000 sq. ft. to work granules down to the soil surface. Application in 7-10 days if necessary. For lawn on St. Augustine grass, use 1-1 1/2 lbs. per 1,000 sq. ft.

Billbugs, White Grubs such as Japanese Beetles: Apply 1 lb. per 1,000 sq. ft. of lawn control make application any time between application if first observed or where they are first seen.

In Case of
Call CHEMTR

Logo TM of WILBUR-ELLIS Company. IDEAS TO GROW WITH is a registered trademark.

BEST AVAILABLE COPY

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS



DIAZINON 14G

NTS
TC ANIMALS
food and feed. Do not store in contact with skin, eyes and clothing and eating. May be absorbed through drift expose workers or protected persons.

XS
drinking on treated areas may be hazardous where surface water is present. Do not apply this product only as directed.

CS
drinking on treated areas may be hazardous where surface water is present. Do not apply this product only as directed.

consistent with its labeling.

are have settled. Because serious crops treated with this information.

ected to be in a treated area or should specify protective clothing. period of time area is to be vacated. When oral warnings are issued by workers. Oral warnings cannot be understood by whom. Area treated with Diazinon protective clothing until dusts exposure.)

CSAL
y storage or disposal.
is product may be disposed
ation equipment. Then dispose, if allowed by State or local air storage areas.

STORAGE
PERSONS
decomposes.

immediately after application and for greatest effectiveness.

ISE
se when insects are present, planting and work into soil 1-2

7-14 lbs. per acre in a 6-inch

Apply 4 1/2-8 1/2 oz. per 1,000 sq. ft. but no later than June 24 in treated band with 2-3 inches of soil.

ng chart for basal treatment, and row in a 10-inch band centered on crop begins to emerge. Cover

ng chart for basal treatment.

ACTIVE INGREDIENTS:	By Wt.
O,O-Diethyl O-(2-isopropyl-5-methyl-4-pyrimidinyl) phosphorothioate	14.3%
INERT INGREDIENTS:	85.7%
TOTAL:	100.0%

EPA Reg. No. 2935-408-AA

EPA Est. No. 33891-AZ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION:

— SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONS —

RECOMMENDATIONS FOR USE (cont.)

Suggested Rates According to Corn Row Spacing

Row Spacing	Linear feet of corn row per acre	Amount of Wilbur-Ellis Diazinon 14G per acre Postemergence basal treatment
48"	10,800	3 - 6 lbs.
42"	12,455	3 1/2 - 6 1/2 lbs.
40"	13,058	3 1/2 - 7 lbs.
38"	13,939	3 3/4 - 7 1/2 lbs.
36"	14,520	4 - 8 lbs.
30"	17,424	4 1/2 - 9 1/2 lbs.
24"	21,780	5 1/2 - 11 1/2 lbs.
20"	26,136	7 - 14 lbs.
15"	34,848	9 1/2 - 18 1/2 lbs.

Note: For row spacings not listed, calculate the appropriate dosage rates based on the 40-in. row figures.

European Corn Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls, when 75% of plants show first-generation borers feeding. Make a second application 7 days later if needed. To control second-generation corn borers, apply 7-14 lbs. per acre, as above, when field count shows 100 egg masses per 100 plants.

WILBUR-ELLIS Diazinon 14G may be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above.

Fall Armyworms: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when armyworms first appear. Repeat application if necessary.

Southwestern Corn Borers: Apply 7-14 lbs. per acre over the corn so granules fall into whorls when second-generation borers first appear. Repeat application one week later.

Note: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be picked immediately. Temporary spotting of foliage sometimes occurs following application.

Cutworms: Broadcast 14-28 lbs. per acre. For surface cutworms apply just prior to planting or when larvae first appear and work into soil 1-2 inches. Do not apply after layoff. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches.

Seed Corn Maggots: Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2 inches.

Garden Symphylans (Centipedes): Broadcast 70 lbs. per acre just prior to planting and work into soil 4-8 inches. This preplant treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

Wireworms: Broadcast 21-28 lbs. per acre before planting. Work into soil 4-6 inches.

FORAGE CROPS (Alfalfa, Clover, Cowpeas and Leaned-seed): CUTWORMS — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

SORGHUM: CUTWORMS — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORN BORERS: Apply 7-14 lbs. per acre in a 10-inch band over the row at planting time or as crop emerges. Cover the band with 1-2 inches of soil.

Note: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOYBEANS: CUTWORMS — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORNSTALK BORERS: Apply 7-14 lbs. per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

SUGAR BEETS: SUGAR BEET ROOT MAGGOT — Apply 7-14 lbs. per acre in a 5-7 inch band (based on row width of 22 ins.) or 4.7-9.4 oz. per 1,000 linear feet of row in a 5-7 inch band with incorporating 2-3 inches at planting or applying above the seed after the seed is covered. Apply ahead of the press wheel. If a heavy maggot infestation is anticipated or a wet spring occurs, a second application may be made, early postemergence, when the adult maggots (flies) first start to appear. If a second application is necessary, apply 7-10 1/2 lbs. per acre in a 5-7 inch band (based on row width of 22 ins.) or 4.7-7.1 oz. per 1,000 linear foot of row in a 5-7 inch band directly over the row.

Note: Diazinon should not come in direct contact with the seed or injury will occur.

WIREWORMS: Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER; ROOT MAGGOTS: Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 3-4 inches.

RADISHES: ROOT MAGGOTS: Apply 10-14 lbs. per acre (2 1/2-2 3/4 oz. per 1,000 linear feet or row) in the seed furrow at planting time.

ONIONS: ONION MAGGOTS: Apply 14-28 lbs. per acre as broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable means.

TOMATOES: VINEGAR FLIES (Drosophila spp.): Broadcast 7 lbs. per acre when flies first appear. Repeat in 7-10 day intervals or as required.

OTHER VEGETABLE INSECTS

OUTWORMS (Surface and Subterranean): Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collards, Cucumbers, Endive (escarole), Kale, Lettuce, Lima Beans, Muskmelons, Parsley, Peppers, Peas, Pole Beans, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash — Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms, 3-6 inches to control subterranean cutworms.

POTATOES: Broadcast 14-28 lbs. per acre before planting. Work into soil 4-6 inches.

MOLE CRICKETS: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Tomatoes, Turnips — Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

LESSER CORNSTALK BORERS: BEANS (Lima, Pole, Snap): Apply 7-14 lbs. per acre in a 6-inch band centered over the row. Apply when plants are 1/2-1 inch tall.

WIREWORMS: Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrots, Cauliflower, Celery, Collards, Cucumbers, Endive (escarole), Kale, Lettuce, Lima Beans, Muskmelons, Onions, Parsley, Peas, Peppers, Pole Beans, Snap Beans, Spinach, Summer Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelons, Winter Squash — Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

POTATOES: Broadcast 21-42 lbs. per acre before planting. Work into soil 4-8 inches or apply 7-21 lbs. per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

GARDEN SYMPHYLANS (CENTIPEDES): Cabbage, Carrots, Lettuce, Peas, Pole Beans, Radishes, Red Beets, Snap Beans, Tomatoes, Turnips — Broadcast 70 lbs. per acre just prior to planting. Work into soil 4-8 inches. This preplant treatment will also control outworms (surface and subterranean), mole crickets, root maggots and wireworms.

LAWN INSECTS (Not for use in California)

Keep children and pets off treated areas until the material is washed into the soil and granules dry.

Lawn Chinch Bugs: Apply 1 1/2 lb. per 1,000 sq. ft. (2 1/2-5 lbs. per 5,000 sq. ft.) and brush lawn to work granules down to the soil surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control and on dense growing turf such as St. Augustine grass, use 1-1 1/2 lbs. per 1,000 sq. ft. (5-7 1/2 lbs. per 5,000 sq. ft.).

Billbugs, White Grubs such as Japanese Beetle Larva, European Chaffer and Southern Chaffer: Apply 1 lb. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application anytime between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

In Case of Emergency,
Call CHEMTREC: (800) 424-9300

Logo TM of WILBUR-ELLIS Company.
IDEAS TO GROW WITH is a registered trademark of WILBUR-ELLIS Company.

BEST AVAILABLE COPY

COPY B

Personal Protective Equipment
Applicators and other handlers must wear:

- A) Long sleeved shirt
- B) Long pants
- C) Waterproof gloves
- D) Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

COPY D

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with it labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protective Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontaminations, notification, and emergency assistance. It also contains specific instructions, and exceptions pertaining/protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

missing language

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- A) Coveralls
- B) Waterproof gloves
- C) Shoes plus socks



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 10 1994

OFFICE OF
PREVENTION, PESTICIDE
TOXIC SUBSTANCES

Joanne Pratt
WILBUR ELLIS CO.
191 W. Shaw Avenue
Fresno, CA 93704

Subject: Label Amendment Submission of 08/30/93 in Response to PR Notice 93-7
EPA Reg. No. 2935-408
DIAZINON 14G

IMPORTANT NOTICE: This Letter Requires You to Submit a Notification to the Office of Compliance Monitoring within 20 Days.

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

SERIOUS ERROR(S) ON SUBMITTED LABELING:

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING. Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.



Recycled/Recyclable
Printed with SoyCanada Ink
contains at least 50% recycled

Each notification must include the following information:

1. Registrant's name and address.
2. EPA Registration number of the product for which the labeling was submitted.
3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

1. Quantity of the product that has been sold or distributed.
2. EPA Producing Establishment number.
3. Location of the product, if known.
4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director
Compliance Division, (7203)
Worker Protection Notification
Office of Compliance Monitoring
U.S. Environmental Protection Agency
401 M Street SW
Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- **BEFORE** selling or distributing any product bearing the final printed labeling as amended

AND

- **WITHIN** one year from date of this acceptance.


Submit the final printed labeling via the U.S. Postal Service to:

**Document Processing Desk (FIN-LABEL)
 Office of Pesticide Programs (7505C)
 U.S. Environmental Protection Agency
 401 M Street, SW
 Washington, D.C. 20460-0001**

Hand or courier deliveries of final labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

cc: Office of Compliance Monitoring

ACCEPTED
with COMMENTS
in EPA Letter Dated
JAN 10 1994

507x

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
5935-608



DIAZINON 14G

ORGANO PHOSPHATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or food products. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. If swallowed, seek medical attention. May be absorbed through skin.

Do not apply this product to water courses, to directly or through drift to water courses or other persons. The area being treated must be vacated by exposed persons.

Copy B - Add

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Copy D - Add

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

RE-ENTRY TREATMENT

Re-entry restrictions must be given to workers who are exposed to this product. For more information, see the Safety Data Sheet (SDS) for this product. For more information, see the Safety Data Sheet (SDS) for this product.

STORAGE AND DISPOSAL

1. Precautions: Do not contaminate water, food or feed by storage or disposal.
2. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
3. Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.
4. Storage: Store in original container and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

ACTIVE INGREDIENTS:

O,O-Dimethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%

INERT INGREDIENTS: 85.7%

TOTAL: 100.0%

By Wt.

FPA Reg. No. 2935-408-AA EPA Est. No. 30991-AZ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION:

— SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONS —

RECOMMENDATIONS FOR USE (cont.)

Suggested Rates According to Corn Row Spacing

Row Spacing	Linear foot of corn row per acre	Amount of Wilbur-Ellis Diazinon 14G per acre Postemergence basal treatment
48"	10,800	3 - 6 lb.
42"	12,455	3 1/4 - 6 1/4 lb.
40"	13,080	3 1/2 - 7 lb.
36"	13,638	3 3/4 - 7 1/4 lb.
30"	14,520	4 - 8 lb.
24"	17,424	4 3/4 - 9 1/4 lb.
20"	21,780	5 3/4 - 11 1/4 lb.
18"	24,144	6 3/4 - 13 1/4 lb.

Note: For row spacings not listed, estimate the appropriate dosage rates based on the 40-in. row figures.

European Corn Borers: Apply 7-14 lb. per acre over the corn as granules fall into whorls when 75% of plants show first-generation borers feeding. Make a second application 7 days later if needed. To control second-generation corn borers, apply 7-14 lb. per acre, as above, when field count shows 100 egg masses per 100 plants.

WILBUR-ELLIS Diazinon 14G may be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above.

Fall Armyworm: Apply 7-14 lb. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary.

Southwestern Corn Borers: Apply 7-14 lb. per acre over the corn as granules fall into whorls when second-generation borers first appear. Repeat application one week later.

Note: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be picked immediately. Temporary spotting of foliage sometimes occurs following application.

Cutworms: Broadcast 14-28 lb. per acre. For surface cutworms apply just prior to planting of when larvae first appear and work into soil 1-2 inches. Do not apply after layby. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches.

Seed Corn Maggots: Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2 inches.

Garden Symphylans (Centipede): Broadcast 70 lb. per acre just prior to planting and work into soil 4-8 inches. This product treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

Wireworms: Broadcast 21-28 lb. per acre before planting. Work into soil 4-8 inches.

FORAGE CROPS (Alfalfa, Clover, Comps and Leppodas): CUTWORMS — Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Sorghum: CUTWORMS — Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORN BORERS — Apply 7-14 lb. per acre in a 10-inch band over the row at planting time or as a top dressing. Cover the band with 1-2 inches of soil.

Note: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOYBEANS: CUTWORMS — Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORNSTALK BORERS — Apply 7-14 lb. per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

SUGAR BEETS: SUGAR BEET ROOT MAGGOT — Apply 7-14 lb. per acre in a 5-7 inch band (based on row width of 22 in.) or 4-7 lb. per acre, per 1,000 linear foot of row in a 5-7 inch band either incorporating 2-3 inches at planting or applying above the seed after the seed is covered. Apply ahead of the grass wheel. If a heavy maggot infestation is anticipated or a root rotting occurs, a second application may be made, early post-emergence, when the adult maggot (flies) first start to appear. If a second application is necessary, apply 7-10 lb. per acre in a 5-7 inch band (based on row width of 22 in.) or 4-7 lb. per acre, per 1,000 linear foot of row in a 5-7 inch band directly over the row.

Note: Diazinon should not come in direct contact with WIREWORMS — Broadcast 21-28 lb. per acre before planting.

VEGETABLES
BROCCOLI, BRUSSELS SPROUTS, CUCUMBERS — Broadcast 14-21 lb. per acre before planting.

RADISHES: ROOT MAGGOTS: Apply 14 lb. per acre in the seed furrow at planting time.

ONIONS: ONION MAGGOTS — Apply 14 lb. per acre just prior to planting. Thoroughly mix the soil in the seed furrow or other suitable means.

TOMATOES: VINEGAR FLIES (Drosophila) — Apply 14 lb. per acre just prior to planting. Repeat in 7-10 day intervals or as needed.

OTHER VEGETABLES
CUTWORMS (Surface and Subterranean) — Broadcast 14-28 lb. per acre before planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

POTATOES — Broadcast 14-28 lb. per acre before planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

MOLE CRICKETS: Broadcast 21-42 lb. per acre before planting. Work into soil 4-8 inches. This product treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

LESSER CORNSTALK BORERS: BEAN BORERS — Apply 7-14 lb. per acre in a 6-inch band over the row when beans are 1-2 inches high. Cover the band with 1-2 inches of soil.

WIREWORMS: Broadcast 21-28 lb. per acre before planting. Work into soil 4-8 inches.

POTATOES — Broadcast 21-42 lb. per acre before planting. Work into soil 4-8 inches. This product treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

GARDEN SYMPHYLANS (CENTIPEDE): Broadcast 70 lb. per acre just prior to planting and work into soil 4-8 inches. This product treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

LAWN INSECTS (Not for use in California): Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

Keep children and pets off treated areas until dry.

Lawn Chinch Bugs: Apply 1/2 lb. per 1,000 sq. ft. to work granules down to the soil surface. If necessary, for lawns at St. Augustine grass, use 1-1 1/2 lb. per 1,000 sq. ft.

White Grubs: Apply 1 lb. per 1,000 sq. ft. of lawn just prior to planting or when white grubs are 1-2 inches high. Cover the band with 1-2 inches of soil.

In Case of
Call CHEMTR

Logo TM of WILBUR-ELLIS Company
IDEAS TO GROW WITH is a registered trademark

BEST AVAILABLE COPY

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS



DIAZINON 14G

INSTRUCTIONS

STRICTLY ANIMALS
of food and feed. Do not store in a container with skin, eyes and clothing and eating. May be absorbed through skin exposure workers or unprotected persons

SODS

working on treated areas may be in areas where surface water is not. Do not apply where runoff is likely from areas treated. Do not water. Apply this product only as

ACTIVE INGREDIENTS:	By WL
O,O-Dimethyl O-(2-isopropyl-4-methyl-4-pyridinyl) phosphorothioate	94.3%
INERT INGREDIENTS:	5.7%
TOTAL:	100.0%

EPA Reg. No. 2935-408-AA EPA Est. No. 32681-AZ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION:

— SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONS —

RECOMMENDATIONS FOR USE (cont.)

Suggested Rates According to Corn Row Spacing

Row Spacing	Linear feet of corn row per acre	Amount of Wilbur-Ellis Diazinon 14G per acre Post-emergence band treatment
48"	10,890	3 - 6 lb.
42"	12,455	3 1/4 - 6 1/2 lb.
40"	13,058	3 1/2 - 7 lb.
36"	13,898	3 3/4 - 7 1/2 lb.
30"	14,520	4 - 8 lb.
30"	17,424	4 1/4 - 9 1/4 lb.
24"	21,780	5 1/4 - 11 1/4 lb.
20"	28,136	7 - 14 lb.
15"	34,848	9 1/4 - 18 1/4 lb.

Note: For row spacings not listed, calculate the appropriate dosage rates based on the 48-in. row figures.

European Corn Borers: Apply 7-14 lb. per acre over the corn as granules fall into whorls, when 75% of plants show first-generation borers feeding. Make a second application 7 days later if needed. To control second-generation corn borers, apply 7-14 lb. per acre, as above, when field count shows 100 egg masses per 100 plants.

WILBUR-ELLIS Diazinon 14G may be applied by aircraft for the above uses. This material should be uniformly applied to the area to be treated at the recommended rates given above.

Fall Armyworms: Apply 7-14 lb. per acre over the corn as granules fall into whorls when armyworms first appear. Repeat application if necessary.

Southern Corn Borers: Apply 7-14 lb. per acre over the corn as granules fall into whorls when second-generation borers first appear. Repeat application one week later.

Note: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be plucked immediately. Temporary spotting of foliage sometimes occurs following application.

Cutworms: Broadcast 14-28 lb. per acre. For surface cutworms apply just prior to planting of when larvae first appear and work into soil 1-2 inches. Do not apply after tassel. For subterranean cutworms apply just prior to planting and work into soil 3-6 inches.

Seed Corn Maggots: Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2 inches.

Garden Symphyla (Centipedes): Broadcast 70 lb. per acre just prior to planting and work into soil 4-8 inches. This preplant treatment will also control other soil insects such as cutworms (surface and subterranean) and wireworms.

Wireworms: Broadcast 21-28 lb. per acre before planting. Work into soil 4-8 inches.

FORAGE CROPS (Alfalfa, Clover, Compsa and Legumes): CUTWORMS—Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

BORGHUMS: CUTWORMS—Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORN BORERS—Apply 7-14 lb. per acre in a 10-inch band over the row at planting time or as crop emerges. Cover the band with 1-2 inches of soil.

Note: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOY BEANS: CUTWORMS—Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms; 3-6 inches to control subterranean cutworms.

LESSER CORNSTALK BORERS—Apply 7-14 lb. per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

SUGAR BEETS: SUGAR BEET ROOT MAGGOT—Apply 7-14 lb. per acre in a 5-7 inch band (based on row width of 22 in.) or 4 7/8 4 oz. per 1,000 linear feet of row in a 5-7 inch band either incorporating 2-3 inches at planting or applying above the seed after the seed is covered. Apply ahead of the press wheel. If a heavy maggot infestation is anticipated or a pest spring occurs, a second application may be made, early post-emergence, when the adult maggot (flies) first start to appear. If a second application is necessary, apply 7-10 lb. per acre in a 5-7 inch band (based on row width of 22 in.) or 4 7/8 1 oz. per 1,000 linear feet of row in a 5-7 inch band directly over the row.

Note: Diazinon should not come in direct contact with the seed or injury will occur.

WIREWORMS—Broadcast 21-28 lb. per acre just prior to planting. Work into soil 4-8 inches.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE and CAULIFLOWER: ROOT MAGGOTS—Broadcast 14-21 lb. per acre before planting or transplanting. Work into soil 3-4 inches.

RADISHES: ROOT MAGGOTS: Apply 10-14 lb. per acre (2 1/4-2 1/2 oz. per 1,000 linear feet or row) in the seed furrow at planting time.

ONIONS: ONION MAGGOTS—Apply 14-28 lb. per acre as broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by plowing, harrowing or other suitable means.

TOMATOES: VINEGAR FLIES (Drosophila spp.)—Broadcast 7 lb. per acre when flies first appear. Repeat in 7-10 day intervals as required.

OTHER VEGETABLE INSECTS

CUTWORMS (Surface and Subterranean): Broadcast, Brussels Sprouts, Cabbage, Cauliflower, Cauliflower, Celery, Collards, Cucumbers, Endive (escarole), Kale, Lettuce, Lima Beans, Madras, Parsley, Peppers, Peas, Pole Beans, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash—Broadcast 14-28 lb. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms, 3-6 inches to control subterranean cutworms.

POTATOES—Broadcast 14-28 lb. per acre before planting. Work into soil 4-8 inches.

MOLE CRICKETS: Broadcast, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Madras, Peppers, Peas, Pole Beans, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash—Apply 7 lb. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

LESSER CORNSTALK BORERS: BEANS (Lima, Pole, Snap)—Apply 7-14 lb. per acre in a 6-inch band centered over the row. Apply when plants are 1-1 1/2 inch tall.

WIREWORMS: Broadcast, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Madras, Peppers, Peas, Pole Beans, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash—Broadcast 21-28 lb. per acre just prior to planting. Work into soil 4-8 inches.

POTATOES—Broadcast 21-42 lb. per acre before planting. Work into soil 4-8 inches or apply 7-21 lb. per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

GARDEN SYMPHYLA (CENTIPEDES): Cabbage, Carrots, Lettuce, Peas, Pole Beans, Radishes, Red Beets, Snap Beans, Tomatoes, Turnips—Broadcast 70 lb. per acre just prior to planting. Work into soil 4-8 inches. This preplant treatment will also control cutworms (surface and subterranean), mole crickets, root maggots and wireworms.

LAWN INSECTS (Not for use in California)

Not for use in California. Do not apply to turf or other areas where the material is washed into the soil and ground.

Lawn Chinch Bugs: Apply 1/2-1 lb. per 1,000 sq. ft. (2 1/4-5 lb. per 5,000 sq. ft.) and brush lawn to work granules down to the soil surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control under dense growing turf such as St. Augustine grass, use 1-1 1/2 lb. per 1,000 sq. ft. (5-7 1/2 lb. per 5,000 sq. ft.).

Billbugs, White Grubs such as Japanese Beetle Larvae, European Chaffer and Southern Chaffer: Apply 1 lb. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chawed or brown grass is present, indicating damage from this insect.

In Case of Emergency,
Call CHEMTREC: (800) 424-9300

Logo TM of WILBUR ELLIS Company
IDEAS TO GROW WITH is a registered trademark of WILBUR ELLIS Company

BEST AVAILABLE COPY

COPY B

Personal Protective Equipment
Applicators and other handlers must wear:

- A) Long sleeved shirt
- B) Long pants
- C) Waterproof gloves
- D) Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

COPY D

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protective Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontaminations, notification, and emergency assistance. It also contains specific instructions, and exceptions pertaining/protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

missing language

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- A) Coveralls
- B) Waterproof gloves
- C) Shoes plus socks