

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Avoid contact with skin, eyes, or clothing. Avoid inhalation of spray mists. In case of contact, flush eyes with plenty of water. Wash exposed skin with plenty of soap and water. Harmful if swallowed.

**DO NOT STORE WITH FOODSTUFFS**

Avoid contamination of water intended for irrigation or domestic uses.

**ENVIRONMENTAL HAZARDS**

**TOXIC TO FISH:** Do not contaminate any body of water by cleaning of equipment and disposing of wastes and containers. Do not apply directly to water.

Avoid spray drifts to susceptible plants as this product may injure cotton, beans, other vegetables, certain legumes, ornamentals, etc. (Coarse sprays are less likely to drift.) Avoid use of small diameter spray nozzles.

**NOT FOR USE OR STORAGE IN OR AROUND HOME.**

**DO NOT** transport with or store near fertilizers, insecticides, fungicides, or seeds.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use Wilbur-Ellis MCPA Sodium Salt with amine salts of 2, 4 D or 2, 4, 5 T. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes or do not use same spray equipment for other purposes.

**STORAGE AND DISPOSAL**

- 1. PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2. PESTICIDE DISPOSAL:** Pesticide, spray, mixture, or rinseate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.
- 3. CONTAINER DISPOSAL:** Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.
- 4. GENERAL:** Consult Federal, State or Local Disposal Authorities for additional restrictions and approved alternatives.

**GENERAL INFORMATION**

**WILBUR-ELLIS MCPA SODIUM SALT** is selective on Broadleaf Weeds only, and has no effect on cereal grains at the recommended rates.

**WILBUR-ELLIS MCPA SODIUM SALT** is a very slow acting systemic herbicide. This property allows the chemical to translocate into the root system before the tops are killed. Other chemicals do not kill this slow, effective kill without sterilizing the soil.

MSS-2

[Wilbur-Ellis]

**MCPA**

**SODIUM SALT**

**ACTIVE INGREDIENT:**

Sodium Salt of 2-Methyl-4-Chlorophenoxyacetic Acid

**INERT INGREDIENTS**

\* Equivalent to 20.05% 2-Methyl-4-Chlorophenoxyacetic Acid, not less than 1 859 lbs. MCPA acid per gallon at 68°F.

\* Isomer specific by A.O.A.C. Method No. 6.A18-22.

E.P.A. Reg. No. 2935-416

**KEEP OUT OF REACH OF CHILDREN  
CAUTION:**

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONS

NET CONTENTS:

Distributed by

[Wilbur-Ellis]

FRESNO • GLENDALE, AZ • PORTLAND •

**SALT**

Acid\* 22.25%  
77.75%  
TOTAL 100.0%

Est. No. 39335-A1 11

**CHILDREN!**

**INARY STATEMENTS**

SEATTLE

**GENERAL INFORMATION (Continued)**

**WILBUR ELLIS MCPA SODIUM SALT** is a selective herbicide for the control of weeds in grain crops. It is effective against many broadleaf weeds and grasses. It is registered for use in the United States and Canada.

**SUSCEPTIBLE WEEDS (Most Areas)** - Broadleaf weeds, including: Horseweed, Field Bindweed, Canada Thistle, White Top, Meadow Buttercup, Field Bindweed, and others. Grasses, including: Quackgrass, and others.

**LESS SUSCEPTIBLE WEEDS** - Includes: Yellow foxtail, and others. For more information, consult the label or contact your local Wilbur-Ellis representative.

**RECOMMENDATIONS FOR USE**

**PEAS** - Apply 2 to 4 pints per acre in sufficient water per acre. For aerial application use 1/2 to 1 pint per acre. Spray or spray weeds soon after they emerge and before they are 3 inches tall. Do not apply later than 1 inch before first pod development. Use of spray pens that are stressed from lack of moisture. Do not use when temperature is over 70°F. Before using, consult State Extension Service or your county agent for general differences to these recommendations.

**SMALL GRAINS - Wheat, Oats, Barley and Rye** - Apply as a water or spray by ground sprayer or airplane. Use 1/2 to 1 pint per acre for the more susceptible weeds after crop has reached the 3 to 4 leaf stage up to boot stage. Use up to 1 pint per acre for the less susceptible weeds after crop has tilled and up to early boot stage. Do not graze or feed meat animals on treated areas within seven days of slaughter.

**WHEAT and BARLEY** - Apply 6 pints per acre when weeds are approaching bud stage, but do not spray grain in the boot to dough stage. The 6 pint (15 pounds acid equivalent) per acre application of any MCPA product can produce injury to wheat and barley. But in the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury. Treat in the spring when grain is fully tilled and 8 to 10 inches tall.

**SMALL GRAINS Underseeded with Alfalfa, Lespedeza, Red and White Clover** - For emergency control of very susceptible weeds (yellow rocket and other susceptible broadleaf weeds) apply 1/2 to 1 pint in not more than 5 to 10 gallons of water per acre. Make application after weeds are well tilled (4 to 8 inches tall) and before reaching boot stage. The use of crop and weeds should provide a protective canopy which, together with the use of low application rates, will reduce the risk of damage to the crop. Do not apply to small grains underseeded with vetch or sweet clover, which are very susceptible. The use of vetch or sweet clover is only if they are protected by a canopy.

**RECOMMENDATIONS FOR USE (Continued)**

**SEED FLAY** - Use 2 to 4 pints of MCPA Sodium Salt per acre in sufficient water to give adequate coverage. Use higher rate for white top, Canada thistle and other less easy to kill weeds. Spray perennials in early bud to full bloom stage and regrowth in the fall. Weeds in spring or fall.

**ESTABLISHED GRASSLANDS AND PASTURES** - Use 2 to 6 pints per acre in sufficient water (10 to 100 gallons) to give thorough coverage. Use higher rate for white top, Canada thistle and other less easy to kill weeds. Spray perennials in early bud to full bloom stage and regrowth in the fall. Weeds in spring or fall.

**ESTABLISHED LAWNS AND TURF** - Use 2 to 4 pints or 3 to 5 gallons of water applied uniformly over 1,000 sq. ft. On larger areas use 2 to 6 pints per acre in 10 to 100 gallons of water. Apply in spring or fall for best results. Do not mow within 2 days before or after application.

**GRASSES GROWN FOR SEED** - Use 2 to 4 pints per acre in sufficient water to give adequate coverage. Use higher rate where weed stands are heavy. In established grasses apply in spring before head comes into boot and on seedling grass after grass has tilled.

**Note** - In weed control in grasses, repeat treatment may be needed for less susceptible weeds. White clover and other legumes may be temporarily injured or killed. In some areas, Dichondra, bent, buffalo carpet and St. Augustine grasses may also be injured by the treatment.

**NON-CROP SPRAYING: Canada Thistle, White Top, Meadow Buttercup, Field Bindweed (Morning-glory)** - For spot treatment on fence rows, rights of way, equipment yards, and similar areas use 1/2 pint to 3 to 4 gallons of water, or 1 1/2 gallons of MCPA Sodium Salt per acre in sufficient water to give coverage and spray to wet weeds thoroughly when in bud to early bloom and again in fall regrowth. Do not graze or feed meat animals on treated areas within seven days of slaughter.

**NOTICE** - The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR ELLIS COMPANY nor the seller shall be held responsible in any manner for any damage, injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on those conditions.

**IN CASE OF EMERGENCY,  
CALL CHEMTREC: (800) 424-9300**