

Wilbur-Ellis
Ideas to grow with.

ACCEPTED
OCT 9 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act or extended for the pesticide registered under EPA Reg. No. 2935-427

**APRON[®]
FLOWABLE**
SEED TREATMENT FUNGICIDE

1013
Pinned
2455

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION
Harmful if swallowed. Avoid breathing vapors or spray mist. May cause eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse. This product may cause an allergic reaction in sensitive individuals.
STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by gently touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF ON SKIN OR IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists.
IF INHALED, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.
IMPORTANT: READ THE ENTIRE DIRECTIONS FOR USE AND THE CONDITIONS OF SALE AND WARRANTY BEFORE USING THIS PRODUCT.
FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

STORAGE AND DISPOSAL
Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
Storage: Do not use or store near heat or open flame.
Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.
Container Disposal: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incinerating, if allowed by state and local authorities.
General: Consult federal, state, or local disposal authorities for approved alternative procedures such as limited open burning.

NOTE: Wilbur-Ellis APRON FLOWABLE is a systemic fungicide having a specific mode of action. Wilbur-Ellis APRON FLOWABLE could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Wilbur-Ellis cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Wilbur-Ellis APRON FLOWABLE resistant strains of fungi which may occur.

Wilbur-Ellis APRON FLOWABLE may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.
Wilbur-Ellis APRON FLOWABLE is a systemic fungicide seed dressing specifically for control of Pythium and Phytophthora spp. For control of other soil-borne diseases, such as Rhizoctonia spp., Wilbur-Ellis APRON FLOWABLE should be applied in combination with other registered seed dressing fungicides.

SEED TREATMENT — DOMESTIC USE
COTTON: For Pythium seed rot and damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment to machine-defoliated or acid-defoliated cottonseed at the rate of .75-1.50 fluid ounces per 100 pounds of seed.

(Continued next page)

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS
For Use Only By Commercial Seed Treaters
ACTIVE INGREDIENTS:
Metalaxyl: N-(2,6-dimethylphenyl)-N-(methylthioacetyl) alanine methyl ester 28.35%
INERT INGREDIENTS 71.65%
TOTAL 100.00%
Contains 2.65 pounds Metalaxyl per gallon.
APRON[®] — Trademark of CIBA-GEIGY for Metalaxyl
EPA Reg. No. 2935-427-AA EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DIRECTIONS FOR USE (Continued)
FORAGE GRASSES: Grasses grown for hay grazing or silage, corn (fodder or silage), and small grain grown for hay grazing or silage.
For Pythium damping-off control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fl. oz. per 100 pounds of seed.
FORAGE LEGUMES—Alfalfa, Clover, Lespedeza, Beans (for forage), Soybeans (for forage), Soybean Hay, Peas (for forage), Pita Vine Hay, Cowpeas (for forage), Cowpea Hay, Trefoil, Vetch and Velvet Beans (for forage).
For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fluid ounces per 100 pounds of seed.
For suppression of early season Downy Mildew in Alfalfa and Soybeans: Apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.
GARDEN BEETS: For control of Pythium damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.
GRAIN CROPS—Wheat, Barley, Rye, Oats, Millet, Rice, Corn (field Corn, Sweet Corn, and Popcorn), Buckwheat and Trifoliate.
For Pythium damping-off control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of .75-1.50 fluid ounces per 100 pounds of seed.
For control of systemic Downy Mildew of Sweet Corn, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00 fl. oz. per 100 pounds of seed.
TURFGRASSES (ALL): For Pythium damping-off control on golf courses, horse trails, reclamation and erosion control projects, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of .50-1.00 fl. oz. per 100 pounds of seed.
SEED AND POD VEGETABLES: Black-eyed Peas, Cowpeas, Dry, Edible Soybeans, Field Beans, Chick Peas (Garbanzo Beans), Field Peas, Garden Peas, Green Beans, Kidney Beans, Lima Beans, Navy Beans, Onions, Peas, Pole Beans, Snap Beans, String Beans, Wax Beans, and Lentils and Lupines.
For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fluid ounces per 100 pounds of seed.
For control of Pythium damping-off and systemic Downy Mildew of peas, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.75 fl. oz. per 100 pounds of seed.
SORGHUM: For control of Pythium damping-off on all sorghum cultivars, use 1.50 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed applied as a seed treatment.
For control of systemic Downy Mildew, apply 1.50 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed on cultivars which possess sorghum Downy Mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum Downy Mildew in heavily infested areas with recent severe history, apply 3.00 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed as a seed treatment. Seed treated with the 3.00 fl. oz. rate should be planted and not carried over to the following year.
Only high quality sorghum seed (90% pure and above) should be treated with Wilbur-Ellis APRON FLOWABLE.
SUGAR BEETS: For control of Pythium damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.
SOYBEANS: For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fl. oz. per 100 pounds of seed.
SUNFLOWER: For control of systemic Downy Mildew, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00 fl. oz. per 100 pounds of seed.
(Continued on back panel)

0.75-1.50

NET CONTENTS:
In Case of Emergency, Call
CHEMTREC: (800) 424-9300

AFS-3

F-817

Distributed By: **WILBUR-ELLIS**, P.O. BOX 16458 • FRESNO, CALIFORNIA 93755

BEST AVAILABLE COPY

ACCEPTED

Wilbur-Ellis
Ideas to grow with.

OCT 9 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2935-427-AA

APRON® FLOWABLE

SEED TREATMENT FUNGICIDE

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION
Harmful if swallowed. Avoid breathing vapors or spray mist. May cause eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse. This product may cause an allergic reaction in sensitive individuals.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by gently touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF ON SKIN OR IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists.
IF INHALED, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.
IMPORTANT: READ THE ENTIRE DIRECTIONS FOR USE AND THE CONDITIONS OF SALE AND WARRANTY BEFORE USING THIS PRODUCT.
FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

STORAGE AND DISPOSAL
Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
Storage: Do not use or store near heat or open flame.
Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.
Container Disposal: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incinerating, if allowed by state and local authorities.
General: Consult federal, state, or local disposal authorities for approved alternative procedures such as limited open burning.

NOTE: Wilbur-Ellis APRON FLOWABLE is a systemic fungicide having a specific mode of action. Wilbur-Ellis APRON FLOWABLE could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Wilbur-Ellis cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Wilbur-Ellis APRON FLOWABLE resistant strains of fungi which may occur.

Wilbur-Ellis APRON FLOWABLE may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.
Wilbur-Ellis APRON FLOWABLE is a systemic fungicide seed dressing specifically for control of Pythium and Phytophthora spp. For control of other soil-borne diseases, such as Rhizoctonia spp., Wilbur-Ellis APRON FLOWABLE should be applied in combination with other registered seed dressing fungicides.

SEED TREATMENT — DOMESTIC USE
COTTON: For Pythium seed rot and damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment to machine-dressed or seed-dressed cottonseed at the rate of .75-1.50 fluid ounces per 100 pounds of seed.
(Continued next page)

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS
For Use Only By Commercial Seed Treaters

ACTIVE INGREDIENTS:
Metalaxyl-N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester 28.35%
INERT INGREDIENTS 71.65%
TOTAL 100.00%

Contains 2.65 pounds Metalaxyl per gallon.
APRON® — Trademark of CIBA-GEIGY for Metalaxyl
EPA Reg. No. 2935-427-AA EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DIRECTIONS FOR USE (Continued)
FORAGE GRASSES: Grasses grown for hay grazing or silage, corn (fodder or silage), and small grain grown for hay grazing or silage.
For Pythium damping-off control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fl. oz. per 100 pounds of seed.
FORAGE LEGUMES—Alfalfa, Clover, Lespedeza, Beans (for forage), Soybeans (for forage), Soybean Hay, Peas (for forage), Pea Vetch Hay, Cowpeas (for forage), Cowpea Hay, Trefoil, Vetch and Vetch Beans (for forage).
For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fluid ounces per 100 pounds of seed for suppression of early season Downy Mildew in Alfalfa and Soybeans. Apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.
GARDEN BEETS: For control of Pythium damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.
GRAIN CROPS—Wheat, Barley, Rye, Oats, Malt, Rice, Corn (Field Corn, Sweet Corn, and Popcorn), B. millet and Triticale.
For Pythium damping-off control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of .75-1.50 fluid ounces per 100 pounds of seed.
For control of systemic Downy Mildew of Sweet Corn, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00 fl. oz. per 100 pounds of seed.
TURFGRASSES (ALL): For Pythium damping-off control on golf courses, home lawns, recreation and erosion control projects, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 4.50-9.00 fl. oz. per 100 pounds of seed.
SEED AND POD VEGETABLES: Black-eyed Peas, Cowpeas, Dill, Edam Soybeans, Field Beans, Chick Peas (Garbanzo Beans), Field Peas, Garden Peas, Green Beans, Kidney Beans, Lima Beans, Navy Beans, Onions, Peas, Pole Beans, Snap Beans, String Beans, Wax Beans, and Lentils and Lupines.
For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fluid ounces per 100 pounds of seed.
For control of Pythium damping-off and systemic Downy Mildew of peas, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.75 fl. oz. per 100 pounds of seed.
SORGHUM: For control of Pythium damping-off on all sorghum cultivars, use 1.50 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed applied as a seed treatment.
For control of systemic Downy Mildew, apply 1.50 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed on cultivars which possess sorghum Downy Mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum Downy Mildew in heavily infested areas with recent severe history, apply 3.00 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed as a seed treatment. Seed treated with the 3.00 fl. oz. rate should be planted and not carried over to the following year.
Only high quality sorghum seed (90% germ and above) should be treated with Wilbur-Ellis APRON FLOWABLE.
SUGAR BEETS: For control of Pythium damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.
SOYBEANS: For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fl. oz. per 100 pounds of seed.
SUNFLOWER: For control of systemic Downy Mildew, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00 fl. oz. per 100 pounds of seed.
(Continued on back panel)

NET CONTENTS:
In Case of Emergency, Call
CHEMTREC: (800) 424-9300

AFS-3

F-587

Distributed By: **WILBUR-ELLIS**, P.O. BOX 16458 • FRESNO, CALIFORNIA 93755

BEST AVAILABLE COPY

DIRECTIONS FOR USE (Continued)

SEED TREATMENT — EXPORT USE

SORGHUM: For control of Pythium damping-off and systemic Downy Mildew of Sorghum, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00-10.50 fl. oz. per 100 pounds of seed.

SUNFLOWER: For control of systemic Downy Mildew, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 5.00-10.50 fl. oz. per 100 pounds of seed.

CORN: For control of Pythium damping-off and systemic Downy Mildew of Corn, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00-10.50 fl. oz. per 100 pounds of seed.

MILLET: For control of Pythium damping-off and systemic Downy Mildew of Millet, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00-10.50 fl. oz. per 100 pounds of seed.

NOTE: Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with Wilbur-Ellis APRON FLOWABLE fungicide. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs." Use with an EPA-approved dye that imparts an unnatural color to the seed.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS CO. does not make any warranty, guaranty, or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS CO. nor the seller shall be responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage, or use of this material not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage, or use and accepts and uses this material on these conditions.

Seed treated in accordance with the export use pattern must be labeled as follows: For Export Only; Not for Domestic Sale or Use.

BEST AVAILABLE COPY