

10/13/89

NOT REVISED
In Accordance with FR Notice of
Based on Draft EPA Registration

Wilbur-Ellis

Ideas to grow with.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION!

Harmful if swallowed. Avoid breathing vapors or spray mist. May cause eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse. This product may cause an allergic reaction in sensitive individuals.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by gently touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF ON SKIN OR IN EYES: Immediately flush with plenty of water. Get medical attention if irritation persists.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.
Do not plant any crop which is not registered for use with metalaxyl or metalaxyl treated soil for a period of 12 months.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
2. Storage: Do not use or store near heat or open flame.
3. Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.
4. Container Disposal: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incinerating, if allowed by State and local authorities.
5. General: Consult Federal, State, or local disposal authorities for approved alternative procedures such as limited open burning.

33786

Manufactured by U.S.A. by: WILBUR-ELLIS CO., P.O. BOX 16458, FRESNO, CA 93755

APRON[®] FLOWABLE

SEED TREATMENT FUNGICIDE

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT
AND DAMPING-OFF DISEASES OF CERTAIN CROPS
For Use Only By Commercial Seed Treaters

ACTIVE INGREDIENTS:

Metalaxyl: N-(2,6-dimethylphenyl)- N-(methoxycarbonyl)alanine methyl ester	28.35%
INERT INGREDIENTS	71.65%
Total	100.00%

Contains 2.65 pounds Metalaxyl per gallon.
APRON[®] — Trademark of CIBA-GEIGY for Metalaxyl
EPA Reg. No. 2935-427-AA

EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NOTE: Wilbur-Ellis APRON FLOWABLE is a systemic fungicide having a specific mode of action. Wilbur-Ellis APRON FLOWABLE could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Wilbur-Ellis cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Wilbur-Ellis APRON FLOWABLE resistant strains of fungi which may occur.

PM 21
C
2935-427
P. 10/13/89

DIRECTIONS FOR USE (cont.)

IMPORTANT: READ THE ENTIRE DIRECTIONS FOR USE AND THE CONDITIONS OF SALE AND WARRANTY BEFORE USING THIS PRODUCT.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

Do not apply this product through any type of irrigation system.

Wilbur-Ellis APRON FLOWABLE may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

WILBUR-ELLIS APRON FLOWABLE is a systemic fungicide seed dressing specifically for control of *Pythium* and *Phytophthora* spp. For control of other soil-borne diseases, such as *Rhizoctonia* spp., Wilbur-Ellis APRON FLOWABLE should be applied in combination with other registered seed dressing fungicides.

Federal law requires that bags containing treated seed shall be labeled with the following information: This seed has been treated with a fungicide. Do not use for feed, food or oil purposes. Store away from feeds or food stuffs, and use with an EPA-approved dye that imparts an unnatural color to the seed.

SEED TREATMENT — DOMESTIC USE

COTTON: For *Pythium* seed rot and damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment to machine-delinted or acid-delinted cottonseed at the rate of 0.75-1.50 fluid ounces per 100 pounds of seed.

FORAGE GRASSES: Grasses grown for hay grazing or silage, corn (fodder or silage), and small grain grown for hay grazing or silage.

For *Pythium* damping-off control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fl. oz. per 100 pounds of seed.

FORAGE LEGUMES — Alfalfa, Clover, Lespedeza, Beans (for forage), Soybeans (for forage), Soybean Hay, Peas (for forage), Pea Vine Hay, Cowpeas (for forage), Cowpea Hay, Trefoil, Vetch and Velvet Beans (for forage)

For *Pythium* damping-off and early season *Phytophthora* control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fluid ounces per 100 pounds of seed.

For suppression of early season Downy Mildew in Alfalfa and Soybeans: Apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.

GARDEN BEETS: For control of *Pythium* damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.

GRAIN CROPS — Wheat, Barley, Rye, Oats, Millet, Rice, Corn (Field Corn, Sweet Corn, and Popcorn), Buckwheat and Triticale

For *Pythium* damping-off control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fluid ounces per 100 pounds of seed.

For control of systemic Downy Mildew of Sweet Corn, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00 fl. oz. per 100 pounds of seed.

TURFGRASSES (ALL): For *Pythium* damping-off control on golf courses, home lawns, reclamation and erosion control projects, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50-3.00 fl. oz. per 100 pounds of seed.

SEED AND POD VEGETABLES — Black-eyed Peas, Cowpeas, Dill, Edible Soybeans, Field Beans, Chick Peas (Garbanzo Beans), Field Peas, Garden Peas, Green Beans, Kidney Beans, Lima Beans, Navy Beans, Okra, Peas, Pole Beans, Snap Beans, String Beans, Wax Beans, Lentils and Lupines

For *Pythium* damping-off and early season *Phytophthora* control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fluid ounces per 100 pounds of seed.

For control of *Pythium* damping-off and systemic Downy Mildew of Peas, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.75 fl. oz. per 100 pounds of seed.

SORGHUM: For control of *Pythium* damping-off on all sorghum cultivars, use 1.50 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed applied as a seed treatment.

For control of systemic Downy Mildew, apply 1.50 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed on cultivars which possess sorghum Downy Mildew resistance. In those cultivars which do not possess resistance to a particular race of sorghum Downy Mildew in heavily infested areas with recent severe history, apply 3.00 fl. oz. of Wilbur-Ellis APRON FLOWABLE per 100 pounds of seed as a seed treatment. Seed treated with the 3.00 fl. oz. rate should be planted and not carried over to the following year.

Only high quality sorghum seed (80% germ and above) should be treated with Wilbur-Ellis APRON FLOWABLE.

SUGAR BEETS: For control of *Pythium* damping-off, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 1.50 fl. oz. per 100 pounds of seed.

SOYBEANS: For *Pythium* damping-off and early season *Phytophthora* control, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 0.75-1.50 fluid ounces per 100 pounds of seed.

SUNFLOWER: For control of systemic Downy Mildew, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 3.00 fl. oz. per 100 pounds of seed.

SEED TREATMENT — EXPORT USE

Seed treated in accordance with the export use pattern must be labeled as follows: For Export Only, Not for Domestic Sale or Use.

SORGHUM: For control of *Pythium* damping-off and systemic Downy Mildew of Sorghum, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 10 fl. oz. per 100 pounds of seed.

SUNFLOWER: For control of systemic Downy Mildew, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 10 fl. oz. per 100 pounds of seed.

CORN: For control of *Pythium* damping-off and systemic Downy Mildew of Corn, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 10 fl. oz. per 100 pounds of seed.

MILLET: For control of *Pythium* damping-off and systemic Downy Mildew of Millet, apply Wilbur-Ellis APRON FLOWABLE as a seed treatment at the rate of 10 fl. oz. per 100 pounds of seed.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

NET CONTENTS:

In Case of Emergency, Call CHEMTREC: (800) 424-9300

F-989

pg. 2 of 2