

ACCEPTED

DEC 21 1989

Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
purpose of registered under
EPA Reg. No. 2935-428

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid if swallowed. Avoid breathing vapors or spray mist. May cause eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. **Prohibitions:** Do not contaminate water, food or feed by storage or disposal.
2. **Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or residue is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
3. **Container Disposal:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. **Storage:** Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

IMPORTANT: Read the entire Directions For Use and the Conditions of Sale and Notice before using this product.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

CONDITIONS OF SALE: DELTA-COAT AD is a systemic fungicide having a specific mode of action. DELTA-COAT AD could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Wilbur-Ellis cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible DELTA-COAT AD resistant strains of fungi which may occur.

USE THE HIGH RATE WHEN CONDITIONS FOR DISEASE EXPRESSION ARE FAVORABLE

ACCEPTED

Wilbur-Ellis

DEC 21 1989

DELTA-COAT AD

SEED TREATMENT FUNGICIDE

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purpose of registered under EPA Reg. No. 2935-428

ACTIVE INGREDIENTS:

Chloranil: 1,2-dichloro-3,3-dimethyl-4,4-dioxobenzene	39.90%
Metharyl: N-(2,6-dimethylphenyl)-N-methoxyacetamide	3.50%
Alkane methyl ester	56.50%
INERT INGREDIENTS:	66.50%
TOTAL	100.00%

Contains 2.9 lbs. a.i. Chloranil and 0.34 lbs. Metharyl per gallon.

EPA Reg. No. 2935-428

EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by gently touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on Skin or In Eyes: Immediately flush with plenty of water. Get medical attention if irritation persists.

If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NET CONTENTS: 1 GALLON

Wilbur-Ellis

P.O. Box 16458 • Fresno, CA 93755

DIRECTIONS FOR

Beets: String beets, black-eyed peas, kidney beans, lima beans, navy beans, and lupins.

For Pythium and Rhizoctonia disease treatment at the rate of 5.1 seed.

Cotton: For Pythium and Rhizoctonia machine defoliated cotton seed, treat at the rate of 5.75-11.75 lb.

Soybeans: For early season Phytophthora damping-off, apply DELTA-COAT 5.5-7.0 fluid ounces per 100 lbs.

Sugar Beets: For Pythium and Rhizoctonia seed treatment at the rate of 5.1 pounds of seed.

COMMERCIAL USE

DELTA-COAT AD may be applied as a wet seed treatment fungicide and fungicide if commercial seed treatment equipment.

NOTE: Federal Law requires that bags contain the following information: "This seed has been treated with a fungicide. Do not use treated seed for food. Stay away from feeds and feed stuffs."

Use with an EPA-approved dye that reports on a

HOPPER-BOX TRE

The following procedure is suggested to those seed before planting. Fill hopper box with half amount of DELTA-COAT AD and mix thoroughly on the seed in the hopper box and add the rest of the seed, again mixing thoroughly. Do not use excess

NOTICE: The statements made on this label or label because of conditions of use which are the COMPANY does not make, nor does it warrant make, any warranty, guarantee or representation by this material or the use thereof, except in the label. Neither WILBUR-ELLIS COMPANY is liable in any manner for any personal injury or to the buyer or to the other person from handling not in accordance with directions. The liability resulting from improper handling, storage or material on these conditions.

In Case of Emer
Call CHEMTREC: (800