

SPECIMEN LABEL

ACCEPTED
DEC 30 1992
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2935-428

NU-GRO

DELTA-COAT AD

SEED TREATMENT FUNGICIDE

ACTIVE INGREDIENTS:

Chloranil, 1,4-dichloro-2,5-dimethylbenzene	38.89%
Metalaxyl, 2-(2,6-dimethylphenyl)-N-methylacetamide	1.99%
INERT INGREDIENTS	59.12%
TOTAL	100.00%

Contains 2.9 lbs. a.i. Chloranil and 0.34 lb. Metalaxyl per gallon.

EPA Reg. No. 2935-428

EPA Est. No. 2935-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by gently touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on Skin or in Eyes: Immediately flush with plenty of water. Get medical attention if irritation persists.

If Inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. May cause eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes and pastures). Do not discharge water when disposing of equipment washwaters.

Do not plant any crop which is not registered for use with metalaxyl in metalaxyl treated soil for a period of 12 months.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- Prohibition:** Do not contaminate water, food or feed by storage or disposal.
- Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or waste is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
- Container Disposal:** This container (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
- Storage:** Store in original container only and keep sealed. Store in closed storage areas. Use caution when re-opening, closing or peeling.

IMPORTANT: Read the entire Directions For Use and the Conditions of Sale and Notice before using this product.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

CONDITIONS OF SALE: DELTA-COAT AD is a systemic fungicide having a specific mode of action. DELTA-COAT AD could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Wilbur-Ellis cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible DELTA-COAT AD resistant strains of fungi which may occur.

USE THE HIGH RATE WHEN CONDITIONS FOR DISEASE EXPRESSION ARE FAVORABLE

NET CONTENTS: 1 GALLON

DIRECTIONS FOR USE (cont.)

- Beans:** String beans, black-eyed peas, cowpeas, field beans, green beans, kidney beans, lima beans, navy beans, pole beans, snap beans, soy beans, and lupins.
- For Pythium and Rhizoctonia damping-off, apply DELTA-COAT AD as a seed treatment at the rate of 5.5-7.8 fluid ounces per 100 pounds of seed.
- Cotton:** For Pythium and Rhizoctonia seed rot and damping-off of seed and machine defoliated cotton seed, apply DELTA-COAT AD as a seed treatment at the rate of 5.75-11.75 fluid ounces per 100 pounds of seed.
- Soybeans:** For early season Phytophthora control and Pythium and Rhizoctonia damping-off, apply DELTA-COAT AD as a seed treatment at the rate of 5.5-7.0 fluid ounces per 100 pounds of seed.
- Sugar Beets:** For Pythium and Rhizoctonia seed rot and damping-off, apply DELTA-COAT AD as a seed treatment at the rate of 10.75 fluid ounces per 100 pounds of seed.

HOPPER-BOX TREATMENT

The following procedure is suggested to thoroughly mix DELTA-COAT AD with the seed before planting. Fill planter box only half-full of seed, add half the required amount of DELTA-COAT AD and mix thoroughly with a stick. Then add the remainder of the seed to the planter box and add the rest of the required amount of product to the seed, again mixing thoroughly. Do not use excess seed for food, feed or oil purposes.

"Copy A"

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, expressed or implied, concerning this material or its use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

3288

F-888

In Case of Emergency, Call CHEMTREC: (800) 424-9300

Wilbur-Ellis

P.O. Box 16458 • Fresno, CA 93755

BEST AVAILABLE COPY

SEED PROTECTANTS Hopper Box Products

Copy A

Always apply Nu-Gro Delta Coat AD to the seed prior to using any plantability aid for vacuum planters.

