

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Causes eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes, or clothing. Remove and wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by gently touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN OR IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists.

IF INHALED, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. When disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: READ THE ENTIRE DIRECTIONS FOR USE AND THE CONDITIONS OF SALE AND WARRANTY BEFORE USING THIS PRODUCT.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Storage: Do not use or store near heat or open flame.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local authorities.

Container Disposal: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incinerating, if allowed by State and local authorities.

General: Consult manufacturer, State, or disposal authorities for approved alternative procedures such as limited open burning.

NOTE: Wilbur-Ellis APRON TL is a systemic fungicide having a specific mode of action. Wilbur-Ellis APRON TL could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted, therefore, Wilbur-Ellis cannot assure liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Wilbur-Ellis APRON TL resistant strains of fungi which may occur.

Wilbur-Ellis APRON TL may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

Do Not Apply This Product Through Any Wilbur-Ellis APRON TL is a systemic fungicide type of seed dressing specifically for control of ~~irrigation systems~~ Pythium and Phytophthora spp. For control of other soil-borne diseases, such as Phizoctonia spp., Wilbur-Ellis APRON TL should be applied in combination with other registered seed dressing fungicides.

NOTE: Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with Wilbur-Ellis APRON TL fungicide. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs." Use with an EPA-approved dye that imparts an unnatural color to the seed.

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS CO. does not make any warranty, guaranty, or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS CO. nor the seller shall be responsible in any matter for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage, or use of this material not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage, or use and accepts and uses this material on these conditions.

**APRON® TL
SEED TREATMENT FUNGICIDE**

A seed treatment chemical for control of systemic downy mildew, seed rot and damping-off diseases of certain crops.

FOR HOPPER BOX APPLICATION AND USE BY
COMMERCIAL SEED TREATERS

ACTIVE INGREDIENT:

Metalaxyl:N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester... 11.5%

INERT INGREDIENTS..... 88.5%

TOTAL..... 100.0%


Contains 1.0 pounds Metalaxyl per gallon.

APRON®-Trademark of CIBA-GEIGY for Metalaxyl.

EPA Reg. No. 2935- EPA Est. No. 2935-CA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY
STATEMENTS**



NET CONTENTS:

In Case of Emergency, Call
QUESTREC: (800) 224-9800

MANUFACTURED IN U.S.A. BY: WILBUR-ELLIS COMPANY, P.O. BOX 16458, FRESNO, CALIF. 93755

BEST AVAILABLE COPY

P 100 1

FORAGE LEGUMES: Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).

For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 4.0 fl. oz. per 100 pounds of seed.

For suppression of early season downy mildew in alfalfa and soybeans: Apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 4.0 fl. oz. per 100 pounds of seed.

GARDEN BEETS: For control of Pythium damping-off, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 4.0 fl. oz. per 100 pounds of seed.

GRAIN CROPS: Wheat, barley, rye, oats, millet, rice, corn (field corn, sweet corn, and popcorn), buckwheat and triticale.

For Pythium damping-off control, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 2.0-4.0 fl. oz. per 100 pounds of seed.

For control of systemic downy mildew of sweet corn, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 8.0 fl. oz. per 100 pounds of seed.

TURFGRASSES (ALL): For Pythium damping-off control on golf courses, home lawns, reclamation and erosion control projects, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of ~~4.0-8.0~~ 2.0 to 4.0 fl. oz. per 100 pounds of seed.

SEED & POD VEGETABLES: Black-eyed peas, cowpeas, dill, edible soybeans, field beans, chick peas (garbanzo beans), field peas, garden peas, green beans, kidney beans, lima beans, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, lentils and lupines.

For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 4.0 fl. oz. per 100 pounds of seed.

For control of Pythium damping-off and systemic downy mildew of peas, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 8.67 fl. oz. per 100 pounds of seed.

SORGHUM: For control of Pythium damping-off on all sorghum cultivars. use 4.0 fl. oz. of Wilbur-Ellis Apron TL per 100 pounds of seed applied as a seed treatment.

For control of systemic downy mildew, apply 4.0 fl. oz. of Wilbur-Ellis

Apron TL per 100 pounds of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 8.0 fl. oz. of Wilbur-Ellis Apron TL per 100 pounds of seed as a seed treatment. Seed treated with the 8.0 fl. oz. rate should be planted and not carried over to the following year.

Only high quality sorghum seed (90% germ and above) should be treated with Wilbur-Ellis Apron TL.

SUGAR BEETS: For control of Pythium damping-off, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 4.0 fl. oz. per 100 pounds of seed.

SOYBEANS: For Pythium damping-off and early season Phytophthora control, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 2.0-4.0 fl. oz. per 100 pounds of seed.

SUNFLOWER: For control of systemic downy mildew, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 8.0 fl. oz. per 100 pounds of seed.

SEED TREATMENT-EXPORT USE - FOR EXPORT ONLY, NOT FOR DOMESTIC SALE OR USE.

SORGHUM: For control of Pythium damping-off and systemic downy mildew of sorghum, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 8.0-28.0 fl. oz. per 100 pounds of seed.

SUNFLOWER: For control of systemic downy mildew, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 16.0-28.0 fl. oz. per 100 pounds of seed.

CORN: For control of Pythium damping-off and systemic downy mildew of corn, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 8.0-28.0 fl. oz. per 100 pounds of seed.

MILLET: For control of Pythium damping-off and systemic downy mildew of millet, apply Wilbur-Ellis Apron TL as a seed treatment at the rate of 8.0-28.0 fl. oz. per 100 pounds of seed.

Addendum To Report # 702784

AEROSOL PARTICLE ANALYSIS

CASCADE IMPACTOR SYSTEM

STAGE	MICRON SIZE	CALIBRATION RUN	+ 60 MINUTES	% OF TOTAL	+ 120 MINUTES	% OF TOTAL
1	>10.0	0	0.36000	1.6%	0.12000	0.7%
2	<10.0 > 6.4	0	0.18600	0.8%	0.09300	0.5%
3	< 6.4 > 3.1	0	4.68000	21.3%	3.02400	17.2%
4	< 3.1 > 1.7	0	10.48800	47.8%	8.76300	49.9%
5	< 1.7 > 0.26	0	6.07200	27.7%	5.41200	30.8%
6	< 0.26 > 0.05	0	0.16200	0.7%	0.16200	0.9%

STAGE	MICRON SIZE	+ 160 MINUTES	% OF TOTAL
1	>10.0	0.12000	0.9%
2	<10.0 > 6.4	0.18600	1.3%
3	< 6.4 > 3.1	3.74400	27.0%
4	< 3.1 > 1.7	6.76200	48.8%
5	< 1.7 > 0.26	2.97000	21.4%
6	< 0.26 > 0.05	0.08100	0.6%