ACCEPTED

GRANOL® N-M

RECEIVED

NOV 12 1990

WECO/ASCO

INSECTICIDE AND FUNGICIDE SEED TREATMENT

A DUAL PURPOSE ready-to-use formulation for drill or planter box application to control smuts, seed decay and seedling blight and for temporary protection of seeds and seedlings against wireworms.

WHEAT-GATSEBARLEY RYES FLAX

ACTIVE INGREDIENTS

MANEB (Manganese ethylenebisoithio arbanate) ... LINDANE (Gamma isomer of Berzene Berachlonde)

50.00 18.75

INSRT INGREDIENTS

TOTAL

KEEP OUT OF REACH OF CHILDREN

WARNING

See side panel for additional precautionary information.

EPA Repland 2935 6. GRANOL is a registered trace marked to

S. Santa

BEST AVAILABLE COPY

Werning

Avoid contact with GRANOL N-M It may be fatal if inhaled, swallowed or absorbed through the skin, Do not breathe dust. Do not get in eyes, on skin or on clothing. Protective clothing (long pants, long sleeve shirt, gloves, hat, boots and a disposable, paper dust mask which covers at least one-third of the face) must be worn during mixing and loading. Wear goggles, a mask over mouth and nose and rubber gloves while handling. Wash thoroughly after handling, and before smoking or eating. Treat outdoors or in a thoroughly ventilated place.

Do not treat seed indoors unless there is thorough ventilation.

Avoid contamination of feed and foodstuffs. Clean up and bury any untreated seed or spillage immediately.

Treated seed should be labelled: Treated seed, do not use for food, feed or oil purposes.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning of equipment of disposal of wastes. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose

It excess treated seeds by burial away from busies of water.

STATEMENT OF PRACTICAL TREATMENT If swallowed, cail a physician and/or the Poison Control Centre immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. If inhaled, remove victim to fresh air. Apply artificial respiration if indicated. If on skin, remove contaminated clothing and wash affected areas with soap and water. If in eyes, flush eyes for at least 15 minutes with water. Call a physician immediately.

PRODUCT INFORMATION

GRANOL N-M is a special seed treatment formulation of lindane and maneb which, when used at the correct dosage rates, protects crops against seedhe diseases such as Stinking Smut (Bunt) of Jat, Oat Smuts, Covered and False Loose Smuts of Barley and Bunt of Rye. It also protects cereal crops against certain seed and soil-borne diseases such as Root Rot and Seedling Blight and for temporary protection of seeds and seedlings against wireworms. GRANOL N-M is especially designed for on-the-farm treatment for drill or planter box application to be mixed directly with the seed in the box, either as it is run into the box or by spreading the treatment in a layer over the leveked seed and thoroughly mixing the treatment into the seed with a paddle.

If maneb becomes wet in storage, its fungicidal effectiveness may be reduced and it may also produce vapors which may be flammable. Keep away from flames or sparks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHEAT, OATS, BARLEY, RYE -

Use 2 oz. per Bushel
FLAX Use 4 oz. per Bushel

Use only recommended rates - lower affounts may not give the desired control. Higher amounts may cause seed injury. Seed should be well cured, dry and cleaned before treatment. For maximum benefits avoid deep seeding and exceptionally early sowing under poor growing conditions.

- Mix GRANOL N-M thoroughly into the seed until
 the powder uniformly coats the seed. Use an
 automatic seed treater that meters the praper
 amount of GRANOL N-M and uniformly mixes
 it as seed is augered from the grain box into the
 planter or drill box.
- Place and level one-half of the seed in the drill or planter box, sprinkle one-half of the total GRANOL N-M required for the full planter box uniformly over the seed. Mix thoroughly with a stick or paddle, then fill the box with seed and sprinkle the remaining one-half of the powder, over the seed and again mix until the seed is uniformly coated.
- Dribble required amount of GRANOL No. 100 each bushel of cereal seed as it is poured into the drill box, and mix until seed is uniformly coated.

DO NOT MIX WITH HANDS.

Wear a property designed dust mask and goggles and change filters frequently while treating seed. Keep away or upwind from any dust that may arise. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store treated seeds beyond planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container tightly closed when not in use. Store away from water, food and feed. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the mearest EPA Regional office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

BEST AVAILABLE COPY

Distributed by
Wilbur-Ellis Company
P.O. Box 16458
Fresno, CA 93755

272